

# FRACTIONAL POROUS MEDIA EQUATIONS: EXISTENCE AND UNIQUENESS OF WEAK SOLUTIONS WITH MEASURE DATA

GABRIELE GRILLO, MATTEO MURATORI, FABIO PUNZO

ABSTRACT. We prove existence and uniqueness of solutions to a class of porous media equations driven by the fractional Laplacian when the initial data are positive finite measures on the Euclidean space  $\mathbb{R}^d$ . For given solutions without a prescribed initial condition, the problem of existence and uniqueness of the initial trace is also addressed. By the same methods we can also treat weighted fractional porous media equations, with a weight that can be singular at the origin, and must have a sufficiently slow decay at infinity (power-like). In particular, we show that the Barenblatt-type solutions exist and are unique. Such a result has a crucial role in [26], where the asymptotic behavior of solutions is investigated. Our uniqueness result solves a problem left open, even in the non-weighted case, in [44].

## 1. INTRODUCTION

The main goal of this note is to prove existence and uniqueness of solutions to the following problem:

$$\begin{cases} \rho(x)u_t + (-\Delta)^s(u^m) = 0 & \text{in } \mathbb{R}^d \times \mathbb{R}^+, \\ \rho(x)u = \mu & \text{on } \mathbb{R}^d \times \{0\}, \end{cases} \quad (1.1)$$

where we assume that  $s \in (0, 1)$ ,  $d > 2s$ ,  $m > 1$ ,  $\mu$  is a positive finite Radon measure on  $\mathbb{R}^d$  (so that  $u \geq 0$ ) and that the (Lebesgue) measurable weight  $\rho$  satisfies

$$c|x|^{-\gamma_0} \leq \rho(x) \leq C|x|^{-\gamma_0} \quad \text{a.e. in } B_1 \quad \text{and} \quad c|x|^{-\gamma} \leq \rho(x) \leq C|x|^{-\gamma} \quad \text{a.e. in } B_1^c \quad (1.2)$$

for some  $\gamma \in [0, 2s)$ ,  $\gamma_0 \in [0, \gamma]$  and  $0 < c < C$ , where  $B_r = B_r(0)$ . Furthermore, for any given solution to the differential equation in (1.1), namely without a prescribed initial datum, we also prove that there exists a unique initial trace which is a finite measure (see Theorem 3.3). Observe that this result suggests that is quite natural to consider a finite measure  $\mu$  as the initial condition in (1.1). We stress that the results concerning uniqueness are new even for  $\rho \equiv 1$ , which obviously fulfills (1.2), thus solving an open problem posed in [44] where such a problem is addressed for initial data given by Dirac deltas, namely for *Barenblatt solutions*. In this case, the problem is known as *fractional porous media equation* and has been thoroughly analysed in [18, 19] for initial data in  $L^1(\mathbb{R}^d)$ . More in general, in view of various applications well outlined in the literature (see e.g. [28]), we also consider the weight  $\rho(x)$  since the same methods of proof work in this case as well. In this respect, observe that even if  $\rho \in C(\mathbb{R}^d)$  has a suitable decay at infinity, and  $\mu = u_0 \in L^1_\rho(\mathbb{R}^d)$ , then the asymptotics of *any* solution can be determined by referring to the *Barenblatt solution* (i.e. the solution to problem (1.1) with  $\mu = \delta$ ) for the problem with *singular, homogeneous* weight  $\rho(x) = |x|^{-\gamma}$ , which makes the problem scale-invariant. Also for this reason we treat weights  $\rho$  that satisfy (1.2), thus being allowed to be singular at  $x = 0$ . However, some further restrictions on  $s$ ,  $d$  and  $\gamma$  will be required and clarified later, see Theorems 3.2 and 3.4. Let us mention that our results entailing the existence and uniqueness of Barenblatt solutions for singular weights are used in a crucial way in [26] to obtain the asymptotic behavior recalled above.

The analysis of the evolutions addressed here poses significant difficulties especially as concerns uniqueness, as can be guessed even when considering their linear analogues. In fact, the first issue

---

*Key words and phrases.* Weighted porous media equation; weighted Sobolev inequalities; nonlinear diffusion equations; smoothing effect; asymptotics of solutions.

we have to deal with is the essential self-adjointness of the operator formally defined as  $\rho^{-1}(-\Delta)^s$  on test functions, and the validity of the Markov property for the associated linear evolution. This will be crucial in the uniqueness proof and holds true only if  $\gamma$  is not too large. For larger  $\gamma$  one expects that suitable conditions at infinity should be required to recover self-adjointness.

Notice that the study of weighted linear differential operators of second order has a long story, see for example [14, Section 4.7] or [34]. Recently, the analysis of the spectral properties of operators which are modeled on the critical operator formally given by  $|x|^2\Delta$  has been performed in [15].

As for nonlinear evolutions, the study of porous media and fast diffusion equations with measure data can be tracked back to the pioneering papers [3, 8, 36, 12]. See [45, Section 13] for details and additional references. The fast diffusion case, which will not be dealt with here, is investigated in [9, 10]: notice that for such evolutions the Dirac delta may not be smoothed into a regular solution, so that different techniques must be used, see the recent paper [37] for a general approach. In [18, 19], the fractional porous media and fast diffusion equations have been introduced and thoroughly studied for initial data which are integrable functions. The construction of Barenblatt solutions and the analysis of their role as asymptotic attractors for general integrable data is performed in [44]. Existence and uniqueness of solutions in the fractional, weighted case is studied in [39, 40]: however, the weight there cannot be singular and data cannot be measures.

Semilinear heat equations with measure data have a long history as well and have recently been studied also in the fractional case, see e.g. [31, 11] and references quoted. We remark that the terminology “measure data” is sometimes used in different contexts in which a measure appears as a source term in certain evolution equations: see e.g. [32].

There is a huge literature on the weighted porous media equation: see for example [16, 17, 21, 22, 24, 25, 27, 28, 29, 38, 41, 42, 43] and references quoted therein. It should be pointed out that the possible singularity of the weight, and the fact that we consider measure data as well, makes our problem significantly different both from the unweighted, fractional case and from the weighted, non-fractional case: straightforward modifications of the strategies used to tackle such problems turn out not to be applicable here.

Finally, notice that fractional porous media equations are being used as a model in several applied contexts, see e.g. [6, Appendix B] and references quoted for details.

**Outline of the paper.** The paper is organized as follows. Section 2 briefly collects some preliminary tools on measure theory, fractional Laplacians and fractional Sobolev spaces. In Section 3 we state our main results. In Section 4 we prove existence of weak solutions and the result concerning existence and uniqueness of the initial trace, whereas in Section 5 uniqueness, which is by far the most delicate issue, is addressed: notice that, although we do not state this explicitly, the proofs work also in the case  $s = 1$  and the corresponding results are new in this context as well for the weighted case. In proving uniqueness, we use a “duality method”, following the same line of reasoning introduced by M. Pierre in [36]. This entails serious new difficulties due to the presence of the fractional diffusion and of the weight  $\rho$ . In Appendix A we recall some technical results on the fractional Laplacian, which are exploited in several approximating procedures developed in the proofs below. In Appendix B we sketch the proof of the main properties of the linear operator formally given by  $\rho^{-1}(-\Delta)^s$ . Such properties are of independent interest but are also crucial in order to establish uniqueness.

## 2. PRELIMINARY TOOLS

In this section we outline some basic notation, definitions and properties that we shall make use of later, which concern weighted Lebesgue spaces, measures, fractional Laplacians, fractional Sobolev spaces and Riesz potentials of measures.

**Weighted Lebesgue spaces.** For a given measurable function  $\rho : \mathbb{R}^d \rightarrow \mathbb{R}^+$  (that is, a weight), we denote as  $L^p_\rho(\mathbb{R}^d)$  (let  $p \in [1, \infty)$ ) the Banach space constituted by all (classes of equivalence of)

measurable functions  $f : \mathbb{R}^d \rightarrow \mathbb{R}$  such that

$$\|f\|_{p,\rho} := \left( \int_{\mathbb{R}^d} |f(x)|^p \rho(x) dx \right)^{1/p} < \infty.$$

In the special case  $\rho(x) = |x|^\alpha$  (let  $\alpha \in \mathbb{R}$ ) we simplify notation and replace  $L_\rho^p(\mathbb{R}^d)$  by  $L_\alpha^p(\mathbb{R}^d)$  and  $\|f\|_{p,\rho}$  by  $\|f\|_{p,\alpha}$ . For the usual unweighted Lebesgue spaces we keep the symbol  $L^p(\mathbb{R}^d)$ , denoting the corresponding norms as  $\|f\|_p$  or  $f\|_{L^p(\mathbb{R}^d)}$ .

**Positive finite measures on  $\mathbb{R}^d$ .** Since in (1.1) we deal with positive, finite measures  $\mu$  on  $\mathbb{R}^d$ , we recall some basic properties enjoyed by the set of such measures, which we denote as  $\mathcal{M}(\mathbb{R}^d)$  (with a slight abuse of notation: this is the usual symbol for the space of *signed* measures on  $\mathbb{R}^d$ ). To begin with, consider a sequence  $\{\mu_n\} \subset \mathcal{M}(\mathbb{R}^d)$ . Following the notation of [36], we say that  $\{\mu_n\}$  converges to  $\mu \in \mathcal{M}(\mathbb{R}^d)$  in  $\sigma(\mathcal{M}(\mathbb{R}^d), C_c(\mathbb{R}^d))$  if there holds

$$\lim_{n \rightarrow \infty} \int_{\mathbb{R}^d} \phi d\mu_n = \int_{\mathbb{R}^d} \phi d\mu \quad \forall \phi \in C_c(\mathbb{R}^d), \quad (2.1)$$

where  $C_c(\mathbb{R}^d)$  is the space of continuous, compactly supported functions on  $\mathbb{R}^d$ . This is usually referred to as *local weak\* convergence* (see [2, Definition 1.58]). A classical theorem in measure theory asserts that if

$$\sup_n \mu_n(\mathbb{R}^d) < \infty \quad (2.2)$$

then there exists  $\mu \in \mathcal{M}(\mathbb{R}^d)$  such that  $\{\mu_n\}$  converges to  $\mu$  in  $\sigma(\mathcal{M}(\mathbb{R}^d), C_c(\mathbb{R}^d))$  up to subsequences (see [2, Theorem 1.59]). The same holds true if we replace  $C_c(\mathbb{R}^d)$  with  $C_0(\mathbb{R}^d)$ , the latter being the closure of the former w.r.t.  $\|\cdot\|_\infty$ . A stronger notion of convergence is the following. A sequence  $\{\mu_n\} \subset \mathcal{M}(\mathbb{R}^d)$  is said to converge to  $\mu \in \mathcal{M}(\mathbb{R}^d)$  in  $\sigma(\mathcal{M}(\mathbb{R}^d), C_b(\mathbb{R}^d))$  if

$$\lim_{n \rightarrow \infty} \int_{\mathbb{R}^d} \phi d\mu_n = \int_{\mathbb{R}^d} \phi d\mu \quad \forall \phi \in C_b(\mathbb{R}^d), \quad (2.3)$$

where  $C_b(\mathbb{R}^d)$  is the space of continuous, bounded functions on  $\mathbb{R}^d$ . Trivially, (2.3) implies (2.1). The opposite holds true under a further hypothesis. That is, if  $\{\mu_n\}$  converges to  $\mu$  in  $\sigma(\mathcal{M}(\mathbb{R}^d), C_c(\mathbb{R}^d))$  and  $\lim_{n \rightarrow \infty} \mu_n(\mathbb{R}^d) = \mu(\mathbb{R}^d)$ , then  $\{\mu_n\}$  converges to  $\mu$  also in  $\sigma(\mathcal{M}(\mathbb{R}^d), C_b(\mathbb{R}^d))$  (see [2, Proposition 1.80]). Notice that if  $\{\mu_n\}$  converges to  $\mu$  in  $\sigma(\mathcal{M}(\mathbb{R}^d), C_c(\mathbb{R}^d))$  and (2.2) holds, a priori one only has a weak\* lower semi-continuity property:

$$\mu(\mathbb{R}^d) \leq \liminf_{n \rightarrow \infty} \mu_n(\mathbb{R}^d)$$

(see again [2, Theorem. 1.59]).

**Fractional Laplacians and fractional Sobolev spaces.** The fractional  $s$ -Laplacian operator which appears in (1.1) is defined, at least for any  $\phi \in \mathcal{D}(\mathbb{R}^d) := C_c^\infty(\mathbb{R}^d)$ , as

$$(-\Delta)^s(\phi)(x) := p.v. \int_{\mathbb{R}^d} \frac{\phi(x) - \phi(y)}{|x - y|^{d+2s}} dy \quad \forall x \in \mathbb{R}^d, \quad (2.4)$$

where  $C_{d,s}$  is a suitable positive constant depending only on  $d$  and  $s$ . However, since a priori we have no clue about the regularity of solutions to (1.1), it is necessary to reformulate the problem in a suitable weak sense, see Definition 3.1 below. Before doing it, we need to introduce some fractional Sobolev spaces. Here we shall mainly deal with  $\dot{H}^s(\mathbb{R}^d)$ , that is the closure of  $\mathcal{D}(\mathbb{R}^d)$  w.r.t. the norm

$$\|\phi\|_{\dot{H}^s}^2 := C_{d,s} \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \frac{(\phi(x) - \phi(y))^2}{|x - y|^{d+2s}} dx dy \quad \forall \phi \in \mathcal{D}(\mathbb{R}^d).$$

Notice that the space usually denoted as  $H^s(\mathbb{R}^d)$  is just  $L^2(\mathbb{R}^d) \cap \dot{H}^s(\mathbb{R}^d)$ . For definitions and properties of the general fractional Sobolev spaces  $W^{r,p}(\mathbb{R}^d)$  we refer the reader e.g. to [20].

The link between the  $s$ -Laplacian and the space  $\dot{H}^s(\mathbb{R}^d)$  can be seen by means of the identity

$$\begin{aligned} C_{d,s} \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \frac{(\phi(x) - \phi(y))(\psi(x) - \psi(y))}{|x - y|^{d+2s}} dx dy &= \int_{\mathbb{R}^d} (-\Delta)^{\frac{s}{2}}(\phi)(x) (-\Delta)^{\frac{s}{2}}(\psi)(x) dx \\ &= \int_{\mathbb{R}^d} \phi(x) (-\Delta)^s(\psi)(x) dx \end{aligned} \quad (2.5)$$

for all  $\phi, \psi \in \mathcal{D}(\mathbb{R}^d)$ , see [20, Section 3]. In particular,  $\|\phi\|_{\dot{H}^s}^2 = \|(-\Delta)^{\frac{s}{2}}(\phi)\|_{L^2}^2$  for all  $\phi \in \mathcal{D}(\mathbb{R}^d)$ . Notice that (2.5) can be shown to hold, by approximation, also when  $\phi \in \mathcal{D}(\mathbb{R}^d)$  is replaced by any  $v \in \dot{H}^s(\mathbb{R}^d)$ , where  $(-\Delta)^{\frac{s}{2}}(v)$  is meant in the sense of distributions. By a further approximation procedure one then gets

$$C_{d,s} \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \frac{(v(x) - v(y))(w(x) - w(y))}{|x - y|^{d+2s}} dx dy = \int_{\mathbb{R}^d} (-\Delta)^{\frac{s}{2}}(v)(x) (-\Delta)^{\frac{s}{2}}(w)(x) dx \quad \forall v, w \in \dot{H}^s(\mathbb{R}^d). \quad (2.6)$$

If we set  $v = w$  in (2.6) we deduce that  $\|v\|_{\dot{H}^s}^2 = \|(-\Delta)^{\frac{s}{2}}(v)\|_{L^2}^2$  also for any  $v \in \dot{H}^s(\mathbb{R}^d)$ . In Sections 4 and 5 (and in Appendix B) we shall deal with functions which belong to  $\dot{H}^s(\mathbb{R}^d)$  and to weighted Lebesgue spaces.

**Riesz potentials.** Another mathematical object deeply linked with the  $s$ -Laplacian is its Riesz kernel, namely the function

$$I_{2s}(x) := \frac{k_{d,s}}{|x|^{d-2s}},$$

where  $k_{d,s}$  is again a positive constant depending only on  $d$  and  $s$ . For a given (possibly signed) finite measure  $\nu$ , one can show that the convolution

$$U^\nu := I_{2s} * \nu$$

yields an  $L^1_{loc}(\mathbb{R}^d)$  function referred to as the *Riesz potential* of  $\nu$ , which formally satisfies

$$(-\Delta)^s(U^\nu) = \nu.$$

That is, still at a formal level, the convolution against  $I_{2s}$  coincides with the operator  $(-\Delta)^{-s}$ . One of the most important and classical references for Riesz potentials is the monograph [30] by N. S. Landkof. In the proof of Theorem 3.2 and throughout Section 5 we shall exploit some crucial properties of Riesz potentials collected in [30], along with their connections with the  $s$ -Laplacian.

### 3. STATEMENTS OF THE MAIN RESULTS

We start by introducing a suitable notion of weak solution to (1.1), in the spirit of [19] and [40].

**Definition 3.1.** *Given a finite positive measure  $\mu$ , by a weak solution to problem (1.1) we mean a nonnegative function  $u$  such that*

$$u \in L^\infty((0, \infty); L^1_\rho(\mathbb{R}^d)) \cap L^\infty(\mathbb{R}^d \times (\tau, \infty)) \quad \forall \tau > 0, \quad (3.1)$$

$$u \in L^2_{loc}((0, \infty); \dot{H}^s(\mathbb{R}^d)), \quad (3.2)$$

$$\begin{aligned} - \int_0^\infty \int_{\mathbb{R}^d} u(x, t) \varphi_t(x, t) \rho(x) dx dt + \int_0^\infty \int_{\mathbb{R}^d} (-\Delta)^{\frac{s}{2}}(u^m)(x, t) (-\Delta)^{\frac{s}{2}}(\varphi)(x, t) dx dt &= 0 \\ \forall \varphi \in C_c^\infty(\mathbb{R}^d \times (0, \infty)) \end{aligned} \quad (3.3)$$

and

$$\operatorname{ess\,lim}_{t \rightarrow 0} \rho(\cdot) u(\cdot, t) = \mu \quad \text{in } \sigma(\mathcal{M}(\mathbb{R}^d), C_b(\mathbb{R}^d)). \quad (3.4)$$

Our first result concerns existence.

**Theorem 3.2.** *Let  $d > 2s$  and assume that  $\rho$  satisfies (1.2) for some  $\gamma \in [0, 2s) \cap [0, d - 2s]$  and  $\gamma_0 \in [0, \gamma]$ . Let  $\mu$  be a positive finite measure. Then there exists a weak solution  $u$  to (1.1) in the sense of Definition 3.1, which conserves the mass in the sense that  $\mu(\mathbb{R}^d) = \int_{\mathbb{R}^d} u(x, t) \rho(x) dx$  for all  $t > 0$ , and satisfies the smoothing effect*

$$\|u(t)\|_{\infty} \leq K t^{-\alpha} \mu(\mathbb{R}^d)^{\beta} \quad \forall t > 0, \quad (3.5)$$

where  $K$  depends only on  $m, \gamma, s, d$  and on  $C$  appearing in (1.2), and

$$\alpha := \frac{d - \gamma}{(m - 1)(d - \gamma) + 2s - \gamma}, \quad \beta := \frac{2s - \gamma}{(m - 1)(d - \gamma) + 2s - \gamma}.$$

In particular,  $u(\cdot, t) \in L^p_{\rho}(\mathbb{R}^d)$  for all  $t > 0$  and  $p \in [1, \infty]$ . The solution satisfies the energy estimates

$$\int_{t_1}^{t_2} \int_{\mathbb{R}^d} |(-\Delta)^{\frac{s}{2}}(u^m)(x, t)|^2 dx dt + \frac{1}{m+1} \int_{\mathbb{R}^d} u^{m+1}(x, t_2) \rho(x) dx = \frac{1}{m+1} \int_{\mathbb{R}^d} u^{m+1}(x, t_1) \rho(x) dx \quad (3.6)$$

and

$$\int_{t_1}^{t_2} \int_{\mathbb{R}^d} |z_t(x, t)|^2 \rho(x) dx dt \leq C \int_{\mathbb{R}^d} u^{m+1}(x, t_1/2) \rho(x) dx \quad (3.7)$$

for all  $t_2 > t_1 > 0$ , where  $z := u^{\frac{m+1}{2}}$  and  $C$  depends on  $m, t_1$  and  $t_2$ .

The method of proof of Theorem 3.2 allows to prove the following result on existence and uniqueness of the initial trace, in the spirit of [6, Section 7] and [4].

**Theorem 3.3.** *Let  $d > 2s$  and assume that  $\rho$  satisfies (1.2) for some  $\gamma \in [0, 2s) \cap [0, d - 2s]$  and  $\gamma_0 \in [0, \gamma]$ . Consider a weak solution  $u$  to  $\rho(x)u_t + (-\Delta)^s(u^m) = 0$  in the sense that  $u$  satisfies (3.1), (3.2) and (3.3). Then there exists a unique positive finite measure  $\mu$  which is the initial trace of  $u$  in the sense of (3.4). The same result holds true if the condition  $u \in L^{\infty}(\mathbb{R}^d \times (\tau, \infty))$  in (3.1) is replaced by the weaker condition  $\int_{t_1}^{t_2} u^m(\cdot, \tau) d\tau \in L^1_{\rho}(\mathbb{R}^d)$  for all  $t_2 > t_1 > 0$ . In particular,  $\mu(\mathbb{R}^d) = \int_{\mathbb{R}^d} u(x, t) \rho(x) dx$  for all  $t > 0$ .*

As for uniqueness of weak solutions we have the next result.

**Theorem 3.4.** *Let  $d > 2s$  and assume that  $\rho$  satisfies (1.2) for some  $\gamma \in [0, 2s) \cap [0, d - 2s]$  and  $\gamma_0 \in [0, \gamma]$ . Let  $u_1, u_2$  be two weak solutions to (1.1) in the sense of Definition 3.1. Suppose that they assume as initial datum the same positive finite measure  $\mu$ , in the sense of (3.4). Then  $u_1 = u_2$ .*

**Remark 3.5.** Notice that, if  $d \geq 4s$ , then the assumptions on  $\gamma$  reduce to  $\gamma \in [0, 2s)$ .

Let us stress that, in order to prove Theorem 3.4, we shall exploit crucially the properties of the operator  $A = \rho^{-1}(-\Delta)^s$  contained in Theorem 3.7 and Proposition B.1 below. Such results are of independent interest; their proofs will be just sketched, to keep the paper in a reasonable length, in Appendix B. Some further details and extensions are given in [33].

**Definition 3.6.** *Let  $d > 2s$  and assume that  $\rho$  satisfies (1.2) for some  $\gamma \in [0, 2s)$  and  $\gamma_0 \in [0, d]$ . We denote as  $X_{s, \rho}$  the Hilbert space of all functions  $v \in L^2_{\rho}(\mathbb{R}^d)$  such that  $(-\Delta)^s(v)$  (as a distribution) belongs to  $L^2_{\rho^{-1}}(\mathbb{R}^d)$ , equipped with the norm*

$$\|v\|_{X_{s, \rho}}^2 := \|v\|_{2, \rho}^2 + \|(-\Delta)^s(v)\|_{2, \rho^{-1}}^2 \quad \forall v \in X_{s, \rho}.$$

**Theorem 3.7.** *Let  $d > 2s$  and assume that  $\rho$  satisfies (1.2) for some  $\gamma \in [0, 2s)$  and  $\gamma_0 \in [0, d]$ . Let  $A : D(A) := X_{s, \rho} \subset L^2_{\rho}(\mathbb{R}^d) \rightarrow L^2_{\rho}(\mathbb{R}^d)$  be the operator*

$$A(v) := \rho^{-1}(-\Delta)^s(v) \quad \forall v \in X_{s, \rho}.$$

Then  $A$  is densely defined, positive and self-adjoint on  $L^2_{\rho}(\mathbb{R}^d)$ , and the quadratic form associated to it is

$$Q(v, v) := \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \frac{(v(x) - v(y))^2}{|x - y|^{d+2s}} dx dy$$

with domain  $D(Q) := L^2_\rho(\mathbb{R}^d) \cap \dot{H}^s(\mathbb{R}^d)$ . Moreover,  $Q$  is a Dirichlet form on  $L^2_\rho(\mathbb{R}^d)$  and  $A$  generates a Markov semigroup  $S_2(t)$  on  $L^2_\rho(\mathbb{R}^d)$ . In particular, for all  $p \in [1, \infty]$  there exists a contraction semigroup  $S_p(t)$  on  $L^p_\rho(\mathbb{R}^d)$ , consistent with  $S_2(t)$  on  $L^2_\rho(\mathbb{R}^d) \cap L^p_\rho(\mathbb{R}^d)$ , which is furthermore analytic with a suitable angle  $\theta_p > 0$  for  $p \in (1, \infty)$ .

#### 4. EXISTENCE OF WEAK SOLUTIONS

We start showing a direct consequence of Definition 3.1, namely the conservation in time of the “mass”  $\int_{\mathbb{R}^d} u(x, t) \rho(x) dx$  (recall that we are considering nonnegative solutions).

**Proposition 4.1.** Let  $d > 2s$  and assume that  $\rho$  satisfies (1.2) for some  $\gamma \in [0, 2s) \cap [0, d - 2s]$  and  $\gamma_0 \in [0, \gamma]$ . Let  $u$  be a weak solution to (1.1) according to Definition 3.1. Then

$$\|u(t)\|_{1, \rho} = \int_{\mathbb{R}^d} u(x, t) \rho(x) dx = \mu(\mathbb{R}^d) \quad \text{for a.e. } t > 0, \quad (4.1)$$

namely we have *conservation of mass*.

*Proof.* We plug into (3.3) the test function  $\varphi_R(x, t) := \vartheta(t)\xi_R(x)$ , where  $\xi_R$  is the same cut-off function as in Lemma A.3 and  $\vartheta$  is a suitable positive, regular and compactly supported approximation of  $\chi_{[t_1, t_2]}$  (let  $t_2 > t_1 > 0$ ). Using (2.6), Lemma A.1, Lemma A.3 and letting  $\vartheta \rightarrow \chi_{[t_1, t_2]}$  in (3.3), it is straightforward to obtain the following estimate:

$$\begin{aligned} & \left| \int_{\mathbb{R}^d} u(x, t_2) \xi_R(x) \rho(x) dx - \int_{\mathbb{R}^d} u(x, t_1) \xi_R(x) \rho(x) dx \right| \\ & \leq c^{-1} \left( \frac{1}{R^{2s}} + \frac{1}{R^{2s-\gamma}} \right) \|(1 + |x|^\gamma)(-\Delta)^s(\xi)\|_\infty \int_{t_1}^{t_2} \int_{\mathbb{R}^d} u^m(x, t) \rho(x) dx dt, \end{aligned} \quad (4.2)$$

where on the r.h.s. we exploited the inequality  $\rho^{-1}(x) \leq c^{-1}(1 + |x|^\gamma)$  for all  $x \in \mathbb{R}^d$ , direct consequence of (1.2). Letting  $R \rightarrow \infty$  in (4.2) and recalling (3.4) we get the conclusion.  $\square$

The proof of existence of weak solutions to (1.1) is based on an approximation procedure, that is on picking a sequence of initial data in  $L^1_\rho(\mathbb{R}^d) \cap L^\infty(\mathbb{R}^d)$  which suitably converges to  $\mu$ . An additional approximation will be needed to deal with the possible singularity of the weight at the origin. The corresponding approximate problems are addressed in the next subsection. Since the procedure is in principle standard although technically delicate, we underline the main points only.

**4.1. Approximate problems with initial data in  $L^1_\rho(\mathbb{R}^d) \cap L^\infty(\mathbb{R}^d)$ .** We are concerned with existence of solutions to the following problem:

$$\begin{cases} \rho(x)u_t + (-\Delta)^s(u^m) = 0 & \text{in } \mathbb{R}^d \times \mathbb{R}^+, \\ u = u_0 & \text{on } \mathbb{R}^d \times \{0\}. \end{cases} \quad (4.3)$$

Such solutions are meant in the sense of Definition 3.1 with  $\mu$  replaced by  $\rho u_0$ .

**Lemma 4.2.** Let  $d > 2s$  and assume that  $\rho$  satisfies (1.2) for some  $\gamma \in [0, 2s) \cap [0, d - 2s]$  and  $\gamma_0 \in [0, \gamma]$ . Let  $u_0 \in L^1_\rho(\mathbb{R}^d) \cap L^\infty(\mathbb{R}^d)$ , with  $u_0 \geq 0$ . Then there exists a weak solution  $u$  to (4.3) which satisfies the energy estimates (3.6), (3.7) with a constant  $C$  depending only on  $m$ ,  $t_1$  and  $t_2$ .

Let us outline the strategy of the proof. We further approximate the problem (4.3) by regularizing the weight  $\rho(x)$  in a neighbourhood of  $x = 0$  (where it can be singular). More precisely, we introduce for any  $\eta > 0$  the following problem (4.3):

$$\begin{cases} \rho_\eta(x)(u_\eta)_t + (-\Delta)^s(u_\eta^m) = 0 & \text{in } \mathbb{R}^d \times \mathbb{R}^+, \\ u_\eta = u_0 & \text{on } \mathbb{R}^d \times \{0\}, \end{cases} \quad (4.4)$$

where  $\{\rho_\eta\} \subset C(\mathbb{R}^d)$  is a family of strictly positive weights which behave like  $|x|^{-\gamma}$  at infinity and approximate  $\rho(x)$  monotonically from below, as  $\eta \rightarrow 0$ . Existence (and uniqueness) of weak solutions to (4.4) for such weights and initial data have been established in [40, Theorem 3.1]. We

get suitable a priori estimates (namely (3.6) applied to  $u_\eta$ , which will be proved later, and (4.7) below), that enable us to pass to the limit as  $\eta \rightarrow 0$ , and obtain a solution to problem (4.3), by standard compactness arguments.

*Proof.* For any  $\eta > 0$  let  $u_\eta$  be the unique solution to problem (4.4). The solutions  $u_\eta$  belong to  $C([0, \infty); L^1_{\rho_\eta}(\mathbb{R}^d))$  and satisfy the bound  $\|u_\eta\|_{L^\infty(\mathbb{R}^d \times (0, \infty))} \leq \|u_0\|_{L^\infty(\mathbb{R}^d)}$ . Exploiting these properties one can show that each  $u_\eta$  satisfies a weak formulation which is slightly stronger than the one of Definition 3.1:

$$\begin{aligned} & - \int_0^T \int_{\mathbb{R}^d} u_\eta(x, t) \varphi_t(x, t) \rho_\eta(x) dx dt + \int_0^T \int_{\mathbb{R}^d} (-\Delta)^{\frac{s}{2}}(u_\eta^m)(x, t) (-\Delta)^{\frac{s}{2}}(\varphi)(x, t) dx dt \\ & = \int_{\mathbb{R}^d} u_0(x) \varphi(x, 0) \rho_\eta(x) dx \end{aligned} \quad (4.5)$$

for all  $T > 0$  and  $\varphi \in C_c^\infty(\mathbb{R}^d \times [0, T])$  such that  $\varphi(\cdot, T) = 0$ , where  $u_\eta^m \in L^2((0, \infty); \dot{H}^s(\mathbb{R}^d))$ . The latter property follows from the validity of the energy identity (3.6) for  $u_\eta$  for all  $t_2 > t_1 \geq 0$ . Formally, (3.6) can be proved by plugging the test function  $\varphi(x, t) := \vartheta(t) u_\eta^m(x, t)$  into the weak formulation (4.5) and letting  $\vartheta$  tend to  $\chi_{[t_1, t_2]}$  as in the proof of Proposition 4.1. In order to justify the validity of (3.6) for  $u_\eta$  rigorously one must proceed as in [19, Section 8]. A crucial point concerns the fact that our solutions are *strong*, which follows by techniques analogous to the ones used in [19, Section 8.1]. We refer the reader to Section 4.5 below for more details. Now:

$$\int_{t_1}^{t_2} \int_{\mathbb{R}^d} |(z_\eta)_t(x, t)|^2 \rho_\eta(x) dx dt \leq C \int_{\mathbb{R}^d} u_\eta^{m+1}(x, t_1/2) \rho(x) dx \quad \forall t_2 > t_1 > 0, \quad (4.6)$$

where  $z_\eta := u_\eta^{\frac{m+1}{2}}$  and  $C$  depends only on  $m$ ,  $t_1$  and  $t_2$ . Formula (4.6) follows as in [19, Lemma 8.1]. Since

$$(u_\eta^m)_t = c_m z_\eta^{\frac{m-1}{2}} (z_\eta)_t \quad \text{and} \quad \|z_\eta\|_{L^\infty(\mathbb{R}^d \times (0, \infty))} = \|u_\eta\|_{L^\infty(\mathbb{R}^d \times (0, \infty))}^{\frac{m+1}{2}} \leq \|u_0\|_{L^\infty(\mathbb{R}^d)}^{\frac{m+1}{2}}.$$

From (4.6) we deduce that

$$\int_{t_1}^{t_2} \int_{\mathbb{R}^d} \left| (u_\eta^m)_t(x, t) \right|^2 \rho_\eta(x) dx dt \leq k \|u_0\|_\infty^{m-1} \quad \forall t_2 > t_1 > 0 \quad (4.7)$$

for a suitable  $k > 0$  independent of  $\eta$ . Moreover, the validity of  $\int_{t_1}^{t_2} \int_{\mathbb{R}^d} |u_\eta^m(x, t)|^2 \rho_\eta(x) dx dt \leq C'$  for all  $t_2 > t_1 \geq 0$  and for another suitable positive constant  $C'$  that depends only on  $m$ ,  $t_1$ ,  $t_2$  and  $u_0$  is ensured by the conservation of mass (4.1) (with  $\rho = \rho_\eta$ ) and by the uniform bound on  $\|u_\eta\|_{L^\infty(\mathbb{R}^d \times (0, \infty))}$ . Using (A.4) and the fact that  $H^s(\mathbb{R}^{d+1})$  is compactly embedded in  $L^2_{loc}(\mathbb{R}^{d+1})$  (see e.g. [20, Theorem 7.1]), one can pass to the limit as  $\eta \rightarrow 0$  in (4.5) and get that the weak limit  $u$  of  $\{u_\eta\}$  satisfies

$$\begin{aligned} & - \int_0^T \int_{\mathbb{R}^d} u(x, t) \varphi_t(x, t) \rho(x) dx dt + \int_0^T \int_{\mathbb{R}^d} (-\Delta)^{\frac{s}{2}}(u^m)(x, t) (-\Delta)^{\frac{s}{2}}(\varphi)(x, t) dx dt \\ & = \int_{\mathbb{R}^d} u_0(x) \varphi(x, 0) \rho(x) dx \end{aligned} \quad (4.8)$$

for all  $T > 0$  and  $\varphi \in C_c^\infty(\mathbb{R}^d \times [0, T])$  such that  $\varphi(\cdot, T) = 0$ . The validity of (3.4) follows by plugging into (4.8) the test function  $\varphi(x, t) := \vartheta(t) \xi_R(x)$ , where  $\xi_R$  is a cut-off function as in Lemma A.3 and  $\vartheta$  is a regular approximation of  $\chi_{[0, t_2]}$ . One then lets  $t_2 \rightarrow 0$  and  $R \rightarrow \infty$ .

The energy estimates (3.6) and (3.7) for  $u$  can be obtained reasoning exactly as above (one uses again the fact that solutions are strong).  $\square$

**4.2. Stroock-Varopoulos inequality and smoothing estimate.** Having at our disposal an existence result for problem (4.3), we can now let  $\rho u_0$  approximate  $\mu$ . In order to show that the corresponding solutions converge to a solution of (1.1), we need first some technical results. We begin with a modification of the classical Stroock-Varopoulos inequality: it is proved here for  $v \in L^\infty(\mathbb{R}^d) \cap \dot{H}^s(\mathbb{R}^d)$  with  $(-\Delta)^s(v) \in L^1(\mathbb{R}^d)$ . Observe that, under the hypothesis that  $v \in L^q(\mathbb{R}^d) \cap \dot{H}^s(\mathbb{R}^d)$  with  $(-\Delta)^s(v) \in L^q(\mathbb{R}^d)$ , for  $q > 1$ , such an inequality can be found, e.g., in [19, Section 5] or [5]. See also [14, formula (2.2.7)] for a similar inequality involving general Dirichlet forms. The present result seems to be new, in view of its functional framework, therefore its proof is given in some detail.

**Lemma 4.3.** *Let  $d > 2s$ . For all nonnegative  $v \in L^\infty(\mathbb{R}^d) \cap \dot{H}^s(\mathbb{R}^d)$  such that  $(-\Delta)^s(v) \in L^1(\mathbb{R}^d)$ , the inequality*

$$\int_{\mathbb{R}^d} v^{q-1}(x)(-\Delta)^s(v)(x) dx \geq \frac{4(q-1)}{q^2} \int_{\mathbb{R}^d} \left| (-\Delta)^{\frac{s}{2}}(v^{\frac{q}{2}})(x) \right|^2 dx \quad (4.9)$$

holds true for any  $q > 1$ .

*Proof.* We shall assume, with no loss of generality, that  $v$  is a regular function. Indeed, by standard mollification arguments, one can always pick a sequence  $\{v_n\} \subset C^\infty(\mathbb{R}^d) \cap L^\infty(\mathbb{R}^d) \cap \dot{H}^s(\mathbb{R}^d)$  such that  $\{v_n\}$  converges pointwise to  $v$ ,  $\|v_n\|_\infty \leq \|v\|_\infty$  and  $\{(-\Delta)^s(v_n)\}$  converges to  $(-\Delta)^s(v)$  in  $L^1(\mathbb{R}^d)$ . This is enough to pass to the limit as  $n \rightarrow \infty$  on the l.h.s. of (4.9), while on the r.h.s. one exploits the weak lower semi-continuity of the  $L^2$  norm.

Consider now the following sequences of functions:

$$\begin{aligned} \psi_n(x) &:= \int_0^{x \wedge \frac{1}{n}} y^{\frac{4s}{d-2s}} dy + (q-1) \int_{\frac{1}{n}}^{x \vee \frac{1}{n}} y^{q-2} dy \quad \forall x \in \mathbb{R}^+, \\ \Psi_n(x) &:= \int_0^{x \wedge \frac{1}{n}} y^{\frac{2s}{d-2s}} dy + (q-1)^{\frac{1}{2}} \int_{\frac{1}{n}}^{x \vee \frac{1}{n}} y^{\frac{q}{2}-1} dy \quad \forall x \in \mathbb{R}^+. \end{aligned}$$

It is plain that  $\psi_n$  and  $\Psi_n$  are absolutely continuous, monotone increasing functions such that  $\psi'_n(x) = [\Psi'_n(x)]^2$  for all  $x \in \mathbb{R}^+$ . For any  $R > 0$ , take a cut-off function  $\xi_R$  as in Lemma A.3. To the function  $\xi_R v$  one can apply Lemma 5.2 of [19] with the choices  $\psi = \psi_n$  and  $\Psi = \Psi_n$ , which yields

$$\int_{\mathbb{R}^d} \psi_n(\xi_R v)(x) (-\Delta)^s(\xi_R v)(x) dx \geq \int_{\mathbb{R}^d} \left| (-\Delta)^{\frac{s}{2}}(\Psi_n(\xi_R v))(x) \right|^2 dx. \quad (4.10)$$

Expanding the  $s$ -Laplacian of the product of two functions, we get that the l.h.s. of (4.10) equals

$$\begin{aligned} &\int_{\mathbb{R}^d} \psi_n(\xi_R v)(x) \xi_R(x) (-\Delta)^s(v)(x) dx + \int_{\mathbb{R}^d} \psi_n(\xi_R v)(x) (-\Delta)^s(\xi_R)(x) v(x) dx \\ &+ 2C_{d,s} \int_{\mathbb{R}^d} \psi_n(\xi_R v)(x) \int_{\mathbb{R}^d} \frac{(\xi_R(x) - \xi_R(y))(v(x) - v(y))}{|x-y|^{d+2s}} dy dx. \end{aligned} \quad (4.11)$$

By dominated convergence,

$$\lim_{R \rightarrow \infty} \int_{\mathbb{R}^d} \psi_n(\xi_R v)(x) \xi_R(x) (-\Delta)^s(v)(x) dx = \int_{\mathbb{R}^d} \psi_n(v)(x) (-\Delta)^s(v)(x) dx.$$

Our aim is to show that the other two integrals in (4.11) go to zero as  $R \rightarrow \infty$ . We have:

$$\begin{aligned} &\left| \int_{\mathbb{R}^d} \psi_n(\xi_R v)(x) (-\Delta)^s(\xi_R)(x) v(x) dx \right| \\ &\leq \|(-\Delta)^s(\xi_R)\|_\infty \left( \frac{d-2s}{d+2s} \int_{\{v \leq \frac{1}{n}\}} v^{\frac{2d}{d-2s}}(x) dx + \psi_n(\|v\|_\infty) \|v\|_\infty \int_{\{v > \frac{1}{n}\}} dx \right) \end{aligned} \quad (4.12)$$



and

$$\begin{aligned}
& \left| \int_{\mathbb{R}^d} \psi_n(\xi_R v)(x) \int_{\mathbb{R}^d} \frac{(\xi_R(x) - \xi_R(y))(v(x) - v(y))}{|x - y|^{d+2s}} dy dx \right| \\
& \leq \|v\|_{\dot{H}^s} \left( \int_{\mathbb{R}^d} [\psi_n(\xi_R v)(x)]^2 \int_{\mathbb{R}^d} \frac{(\xi_R(x) - \xi_R(y))^2}{|x - y|^{d+2s}} dy dx \right)^{\frac{1}{2}} \\
& \leq \|v\|_{\dot{H}^s} \|l_s(\xi_R)\|_{\infty}^{\frac{1}{2}} \left( \left[ \frac{d-2s}{d+2s} \right]^2 \int_{\{v \leq \frac{1}{n}\}} v^{2\frac{d+2s}{d-2s}}(x) dx + [\psi_n(\|v\|_{\infty})]^2 \int_{\{v > \frac{1}{n}\}} dx \right)^{\frac{1}{2}},
\end{aligned} \tag{4.13}$$

where  $l_s$  is defined in Lemma A.2. Thanks to the scaling properties of both  $(-\Delta)^s(\xi_R)$  and  $l_s(\xi_R)$  (Lemma A.3), it is immediate to check that  $\lim_{R \rightarrow \infty} \|(-\Delta)^s(\xi_R)\|_{\infty} = \lim_{R \rightarrow \infty} \|l_s(\xi_R)\|_{\infty} = 0$ . Moreover, notice that  $v \in L^{\frac{2d}{d-2s}}(\mathbb{R}^d) \cap L^{\infty}(\mathbb{R}^d)$  (see [20, Section 6] or Lemma 4.4 below). In particular,  $v$  also belongs to  $L^{2\frac{d+2s}{d-2s}}(\mathbb{R}^d)$ . Thus, letting  $R \rightarrow \infty$  in (4.12) and (4.13), we deduce that the last two integrals in (4.11) vanish, so that we can pass to the limit on the l.h.s. of (4.10). On the r.h.s. we just use the fact that  $(-\Delta)^{\frac{s}{2}}(\Psi_n(\xi_R v))$  converges to  $(-\Delta)^{\frac{s}{2}}(\Psi_n(v))$  weakly in  $L^2(\mathbb{R}^d)$ . This proves the validity of

$$\int_{\mathbb{R}^d} \psi_n(v)(x) (-\Delta)^s(v)(x) dx \geq \int_{\mathbb{R}^d} |(-\Delta)^{\frac{s}{2}}(\Psi_n(v))(x)|^2 dx. \tag{4.14}$$

The final step is to let  $n \rightarrow \infty$  in (4.14). It is clear that the sequence  $\{\psi_n(x)\}$  converges locally uniformly to the function  $x^{q-1}$ , while  $\{\Psi_n(x)\}$  converges locally uniformly to  $2(q-1)^{\frac{1}{2}}x^{\frac{q}{2}}/q$ . Hence,  $\{\psi_n(v)\}$  and  $\{\Psi_n(v)\}$  converge in  $L^{\infty}(\mathbb{R}^d)$  to  $v^{q-1}$  and  $2(q-1)^{\frac{1}{2}}v^{\frac{q}{2}}/q$ , respectively. This is enough in order to pass to the limit in (4.14) and obtain (4.9).  $\square$

**Lemma 4.4.** *Let  $d > 2s$  and assume that  $\rho$  satisfies (1.2) for some  $\gamma \in [0, 2s) \cap [0, d-2s]$  and  $\gamma_0 \in [0, \gamma]$ . There exists a positive constant  $C_{CKN} = C_{CKN}(C, \gamma, s, d)$  such that the Caffarelli-Kohn-Nirenberg type inequalities*

$$\|v\|_{q,\rho} \leq C_{CKN} \|(-\Delta)^{\frac{s}{2}}(v)\|_2^{\frac{1}{\alpha+1}} \|v\|_{p,\rho}^{\frac{\alpha}{\alpha+1}} \quad \forall v \in L^p_{\rho}(\mathbb{R}^d) \cap \dot{H}^s(\mathbb{R}^d) \tag{4.15}$$

hold true for any  $\alpha \geq 0$ ,  $p \geq 1$  and  $q = 2(d-\gamma)(\alpha+1)/[(d-\gamma)^{\frac{\alpha}{p}} + d-2s]$ .

*Proof.* See e.g. [13, Theorem 1.8], where one considers the Sobolev inequality corresponding to  $\alpha = 0$  here, and then uses an elementary interpolation.  $\square$

Lemmas 4.3 and 4.4 provide us with some functional inequalities which are crucial to prove the following *smoothing effect* for solutions to (4.3).

**Proposition 4.5.** *Let  $d > 2s$  and assume that  $\rho$  satisfies (1.2) for some  $\gamma \in [0, 2s) \cap [0, d-2s]$  and  $\gamma_0 \in [0, \gamma]$ . There exists a constant  $K > 0$  depending only on  $m, \gamma, s, d$  and  $C$  such that, for all nonnegative initial datum  $u_0 \in L^1_{\rho}(\mathbb{R}^d) \cap L^{\infty}(\mathbb{R}^d)$  and the corresponding weak solution  $u$  to (4.3) constructed in Lemma 4.2, the following  $L^{p_0}$ - $L^{\infty}$  smoothing effect holds true for any  $p_0 \in [1, \infty)$ :*

$$\|u(t)\|_{\infty} \leq K t^{-\alpha_0} \|u_0\|_{p_0,\rho}^{\beta_0} \quad \forall t > 0, \tag{4.16}$$

where

$$\alpha_0 := \frac{d-\gamma}{(m-1)(d-\gamma) + (2s-\gamma)p_0}, \quad \beta_0 := \frac{(2s-\gamma)p_0}{(m-1)(d-\gamma) + (2s-\gamma)p_0}. \tag{4.17}$$

*Proof.* We omit the details, since the claim follows as in [19, Theorem 8.2] by means of a standard parabolic Moser iteration. Nevertheless, notice that the proof relies on the Stroock-Varopoulos inequality (which has to hold for the precise set of functions stated in Lemma 4.3), the Caffarelli-Kohn-Nirenberg type inequalities provided by Lemma 4.4 and the fact that the  $L^p_{\rho}$  norms do not increase along the evolution (see Section 4.5).  $\square$

**4.3. Proof of the existence result.** We outline the main steps of this proof. Suppose first that  $\mu$  is compactly supported. Consider the family  $\{u_\varepsilon\}$  of weak solutions to (1.1) that take on the regular initial data  $\mu_\varepsilon := \psi_\varepsilon * \mu$  (let  $\varepsilon > 0$ ), where  $\psi_\varepsilon := \frac{1}{\varepsilon^d} \psi\left(\frac{x}{\varepsilon}\right)$  with  $\psi \in \mathcal{D}_+(\mathbb{R}^d)$  and  $\|\psi\|_1 = 1$ . The existence of such family is ensured by Lemma 4.2, upon setting  $u_0 = \rho^{-1}\mu_\varepsilon$ . In view of certain a priori estimates (see (4.18), (4.19) and (4.20) below), we prove that  $\{u_\varepsilon\}$  converges (up to subsequences), as  $\varepsilon \rightarrow 0$ , to a function  $u$  which satisfies (3.1), (3.2) and (3.3). Afterwards we shall deal with (3.4). To do this, we exploit some results in potential theory, following [36] or [44], using the Riesz potential  $U_\varepsilon(\cdot, t)$  of  $\rho(\cdot)u_\varepsilon(\cdot, t)$ . Then we let  $\varepsilon \rightarrow 0$ ; in doing this, a uniform estimate w.r.t.  $\varepsilon$  for the potentials (see (4.26) below) will be crucial. Finally, we consider general positive finite measures  $\mu$ , by a further approximation.

*Proof of Theorem 3.2.* For any  $\varepsilon > 0$ , let  $u_\varepsilon$  as above. Combining the smoothing effect (4.16) with the fact that  $\|\mu_\varepsilon\|_1 = \mu(\mathbb{R}^d)$  and with the conservation of mass (4.1), we obtain:

$$\int_{\mathbb{R}^d} u_\varepsilon^{m+1}(x, t) \rho(x) dx \leq \|u_\varepsilon(t)\|_\infty^m \|\mu_\varepsilon\|_1 \leq K^m t^{-\alpha m} \mu(\mathbb{R}^d)^{1+\beta m} \quad (4.18)$$

for all  $t > 0$ . Hence, using (3.6), (3.7) and (4.18) we get:

$$\int_{t_1}^{t_2} \int_{\mathbb{R}^d} |(-\Delta)^{\frac{s}{2}}(u_\varepsilon^m)(x, t)|^2 dx dt + \int_{\mathbb{R}^d} u_\varepsilon^{m+1}(x, t_2) \rho(x) dx \leq K^m t_1^{-\alpha m} \mu(\mathbb{R}^d)^{1+\beta m}, \quad (4.19)$$

$$\int_{t_1}^{t_2} \int_{\mathbb{R}^d} |(z_\varepsilon)_t(x, t)|^2 \rho(x) dx dt \leq C \int_{\mathbb{R}^d} u_\varepsilon^{m+1}(x, t_1/2) \rho(x) dx \quad (4.20)$$

for all  $t_2 > t_1 > 0$ , where  $z_\varepsilon := u_\varepsilon^{\frac{m+1}{2}}$  and  $C$  is a positive constant that depends on  $m, t_1, t_2$  but is independent of  $\varepsilon$ . Thanks to (4.19), (4.20), the conservation of mass and the smoothing effect (which, in particular, bounds  $\{u_\varepsilon\}$  in  $L^\infty(\mathbb{R}^d \times (\tau, \infty))$  for all  $\tau > 0$  independently of  $\varepsilon$ ), we are allowed to proceed exactly as in the proof of Lemma 4.2. That is, we obtain that the pointwise limit  $u$  of  $\{u_\varepsilon\}$ , up to subsequences, satisfies (3.1), (3.2) and (3.3).

Let us now introduce the Riesz potential  $U_\varepsilon(\cdot, t)$  of  $\rho(\cdot)u_\varepsilon(\cdot, t)$ . The equation solved by  $u_\varepsilon$  is

$$\rho(x)(u_\varepsilon)_t(x, t) = -(-\Delta)^s(u_\varepsilon^m)(x, t) \quad \forall (x, t) \in \mathbb{R}^d \times \mathbb{R}^+. \quad (4.21)$$

Applying to both sides of (4.21) the operator  $(-\Delta)^{-s}$ , namely the convolution against the Riesz kernel  $I_{2s}$  (recall the discussion in Section 2), formally yields

$$(U_\varepsilon)_t(x, t) = -u_\varepsilon^m(x, t) \quad \forall (x, t) \in \mathbb{R}^d \times \mathbb{R}^+. \quad (4.22)$$

To prove rigorously (4.22), we plug into (3.3) (with  $u = u_\varepsilon$ ) the test function  $\varphi(y, t) := \vartheta(t)\phi(y)$ , where  $\vartheta$  is a smooth and compactly supported approximation of  $\chi_{[t_1, t_2]}$  and  $\phi \in \mathcal{D}(\mathbb{R}^d)$ . Integrating by parts (in space), letting  $\vartheta$  tend to  $\chi_{[t_1, t_2]}$  and replacing the function  $\phi(y)$  by  $\phi(y+x)$ , with  $x$  fixed, we get:

$$\begin{aligned} & \int_{\mathbb{R}^d} u_\varepsilon(y, t_2) \phi(y+x) \rho(y) dy - \int_{\mathbb{R}^d} u_\varepsilon(y, t_1) \phi(y+x) \rho(y) dy \\ &= - \int_{\mathbb{R}^d} \left( \int_{t_1}^{t_2} u_\varepsilon^m(y, t) dt \right) (-\Delta)^s(\phi)(y+x) dy. \end{aligned} \quad (4.23)$$

Integrating (4.23) against the Riesz kernel  $I_{2s}(x)$  and applying Fubini's Theorem gives (let  $z = y+x$ )

$$\begin{aligned} & \int_{\mathbb{R}^d} U_\varepsilon(z, t_2) \phi(z) dz - \int_{\mathbb{R}^d} U_\varepsilon(z, t_1) \phi(z) dz \\ &= - \int_{\mathbb{R}^d} \left( \int_{t_1}^{t_2} u_\varepsilon^m(y, t) dt \right) \left( \int_{\mathbb{R}^d} (-\Delta)^s(\phi)(y+x) I_{2s}(x) dx \right) dy = - \int_{\mathbb{R}^d} \left( \int_{t_1}^{t_2} u_\varepsilon^m(y, t) dt \right) \phi(y) dy. \end{aligned} \quad (4.24)$$

The applicability of Fubini's Theorem is justified thanks to Lemma A.5, Lemma A.1 (recall that  $d - 2s \geq \gamma$  by assumption) and to the fact that  $\int_{t_1}^{t_2} u_\varepsilon^m(\cdot, t) dt$  belongs to  $L^1_\rho(\mathbb{R}^d) \cap L^\infty(\mathbb{R}^d)$  by (3.1).

By Lemma 4.2 and Definition 3.1, we know that  $\rho u_\varepsilon(t)$  converges to  $\mu_\varepsilon$  in  $\sigma(\mathcal{M}(\mathbb{R}^d), C_b(\mathbb{R}^d))$  as  $t \rightarrow 0$ . Hence, letting  $t_1 \rightarrow 0$  in (4.24), we find that

$$\int_{\mathbb{R}^d} U_\varepsilon(x, t_2) \phi(x) \, dx - \int_{\mathbb{R}^d} U^{\mu_\varepsilon}(x) \phi(x) \, dx = - \int_{\mathbb{R}^d} \left( \int_0^{t_2} u_\varepsilon^m(x, t) \, dt \right) \phi(x) \, dx \quad (4.25)$$

for all  $t_2 > 0$  and  $\phi \in \mathcal{D}(\mathbb{R}^d)$ . In fact,

$$\begin{aligned} \int_{\mathbb{R}^d} U_\varepsilon(x, t_1) \phi(x) \, dx &= \int_{\mathbb{R}^d} \left( \int_{\mathbb{R}^d} I_{2s}(x-y) \rho(y) u_\varepsilon(y, t_1) \, dy \right) \phi(x) \, dx \\ &= \int_{\mathbb{R}^d} \underbrace{\left( \int_{\mathbb{R}^d} I_{2s}(y-x) \phi(x) \, dx \right)}_{U^\phi(y)} \rho(y) u_\varepsilon(y, t_1) \, dy, \end{aligned}$$

and in view of Lemma A.5 we know that, in particular,  $U^\phi \in C_0(\mathbb{R}^d)$ , which allows to pass to the limit in the integral as  $t_1 \rightarrow 0$ . Thanks to the smoothing effect, the conservation of mass and the hypotheses on  $\rho$ , we can provide the following bound for (4.25):

$$\left| \int_{\mathbb{R}^d} U_\varepsilon(x, t_2) \phi(x) \, dx - \int_{\mathbb{R}^d} U^{\mu_\varepsilon}(x) \phi(x) \, dx \right| \leq \|\rho^{-1} \phi\|_\infty K^{m-1} \mu(\mathbb{R}^d)^{1+\beta(m-1)} \int_0^{t_2} t^{-\alpha(m-1)} \, dt. \quad (4.26)$$

Note that the time integral in the r.h.s. is finite since  $\alpha(m-1) < 1$  (recall (4.17) for  $p_0 = 1$ ). We proved above that  $\{u_\varepsilon\}$  converges pointwise a.e. (up to subsequences) to a function  $u$  which satisfies (3.1), (3.2) and (3.3). If we exploit once again the smoothing effect and the conservation of mass, we easily infer that such convergence also takes place in  $\sigma(\mathcal{M}(\mathbb{R}^d), C_0(\mathbb{R}^d))$ :

$$\lim_{\varepsilon \rightarrow 0} \rho u_\varepsilon(t) = \rho u(t) \quad \text{in } \sigma(\mathcal{M}(\mathbb{R}^d), C_0(\mathbb{R}^d)), \text{ for a.e. } t > 0. \quad (4.27)$$

Using (4.27), the fact that  $\mu_\varepsilon \rightarrow \mu$  in  $\sigma(\mathcal{M}(\mathbb{R}^d), C_b(\mathbb{R}^d))$  and proceeding exactly as we did in the proof of (4.25), we can let  $\varepsilon \rightarrow 0$  in (4.26) to get

$$\left| \int_{\mathbb{R}^d} U(x, t_2) \phi(x) \, dx - \int_{\mathbb{R}^d} U^\mu(x) \phi(x) \, dx \right| \leq \|\rho^{-1} \phi\|_\infty K^{m-1} \mu(\mathbb{R}^d)^{1+\beta(m-1)} \int_0^{t_2} t^{-\alpha(m-1)} \, dt \quad (4.28)$$

for a.e.  $t_2 > 0$  and  $\phi \in \mathcal{D}(\mathbb{R}^d)$ , where we denote as  $U$  the potential of  $\rho u$ . Note that, passing to the limit in (4.25) for any nonnegative  $\phi \in \mathcal{D}(\mathbb{R}^d)$ , we deduce in particular that  $U(x, t)$  is nonincreasing in  $t$ . Moreover, (4.28) implies that  $U(t)$  converges to  $U^\mu$  in  $L^1_{\text{loc}}(\mathbb{R}^d)$ , whence

$$\lim_{t \rightarrow 0} U(x, t) = U^\mu(x) \quad \text{for a.e. } x \in \mathbb{R}^d. \quad (4.29)$$

Letting  $\varepsilon \rightarrow 0$  in the conservation of mass (4.1) (applied to  $u = u_\varepsilon$  and  $\mu = \mu_\varepsilon$ ), by means *e.g.* of Fatou's Lemma we obtain

$$\|u(t)\|_{1,\rho} \leq \mu(\mathbb{R}^d) \quad \text{for a.e. } t > 0. \quad (4.30)$$

Due to the compactness results recalled in Section 2, from (4.30) we infer that (almost) every sequence  $t_n \rightarrow 0$  admits a subsequence  $\{t_{n_k}\}$  such that  $\{\rho u(t_{n_k})\}$  converges to a certain positive finite Radon measure  $\nu$  in  $\sigma(\mathcal{M}(\mathbb{R}^d), C_c(\mathbb{R}^d))$ . Thanks to (4.29) and [30, Theorem 3.8] we have that  $U^\nu(x) = U^\mu(x)$  almost everywhere. Alternatively, such identity can be proved by passing to the limit in  $\int_{\mathbb{R}^d} U(x, t_{n_k}) \phi(x) \, dx$ , recalling that  $U(t_{n_k}) \rightarrow U^\mu$  in  $L^1_{\text{loc}}(\mathbb{R}^d)$  as  $k \rightarrow \infty$ . Theorem 1.12 of [30] then ensures that two positive finite Radon measures whose potentials are equal almost everywhere must coincide. Hence,  $\nu = \mu$  and the limit measure does not depend on the particular subsequence, so that

$$\lim_{t \rightarrow 0} \rho u(t) = \mu \quad \text{in } \sigma(\mathcal{M}(\mathbb{R}^d), C_c(\mathbb{R}^d)).$$

In order to show that convergence also takes place in  $\sigma(\mathcal{M}(\mathbb{R}^d), C_b(\mathbb{R}^d))$ , it is enough to establish that

$$\lim_{t \rightarrow 0} \|u(t)\|_{1,\rho} = \mu(\mathbb{R}^d). \quad (4.31)$$

Since  $\rho u(t)$  converges to  $\mu$  in  $\sigma(\mathcal{M}(\mathbb{R}^d), C_c(\mathbb{R}^d))$  as  $t \rightarrow 0$ , we know that

$$\mu(\mathbb{R}^d) \leq \liminf_{t \rightarrow 0} \|u(t)\|_{1,\rho}, \quad (4.32)$$

see again Section 2. But (4.32) and (4.30) entail (4.31).

Finally, the validity of the smoothing estimate (3.5) is just a consequence of passing to the limit in (4.16) (applied to  $u_\varepsilon$  and  $p_0 = 1$ ) as  $\varepsilon \rightarrow 0$  (recall that  $\{u_\varepsilon\}$  converges pointwise to  $u$ ).

At the beginning of the proof we required  $\mu$  to be compactly supported. Otherwise, take a sequence of compactly supported measures  $\{\mu_n\}$  converging to  $\mu$  in  $\sigma(\mathcal{M}(\mathbb{R}^d), C_b(\mathbb{R}^d))$  and consider the corresponding sequence of solutions  $\{u_n\}$  to (1.1). Estimates (4.19) and (4.20), as well as the conservation of mass and the smoothing effect, are clearly stable as  $\varepsilon \rightarrow 0$ , thus they also hold true upon replacing  $u_\varepsilon$  with  $u_n$  and  $\mu_\varepsilon$  with  $\mu_n$ . Hence, using the same techniques as above, one proves that  $\{u_n\}$  converges to a solution  $u$  of (1.1) starting from  $\mu$ .  $\square$

**4.4. Existence and uniqueness of initial traces.** We use the next preliminary result.

**Lemma 4.6.** *Let  $\nu$  be a signed finite measure such that  $U^\nu \geq 0$  almost everywhere. Then  $\nu(\mathbb{R}^d) \geq 0$ .*

*Proof.* From the assumptions on  $U^\nu$  and thanks to Fubini's Theorem, there holds

$$\int_{\mathbb{R}^d} \chi_{B_n}(y) U^\nu(y) dy = \int_{\mathbb{R}^d} (I_{2s} * \chi_{B_n})(x) d\nu = k_{d,s} \int_{\mathbb{R}^d} \left( \int_{B_n} |x-y|^{-d+2s} dy \right) d\nu \geq 0 \quad \forall n \in \mathbb{N}. \quad (4.33)$$

Performing the change of variable  $z = y/n$ , the last inequality in (4.33) reads

$$\int_{\mathbb{R}^d} \left( \int_{B_1} |x/n - z|^{-d+2s} dz \right) d\nu \geq 0 \quad \forall n \in \mathbb{N}. \quad (4.34)$$

It is plain that for every  $x \in \mathbb{R}^d$  the sequence  $\{\int_{B_1} |x/n - z|^{-d+2s} dz\}$  converges to the positive constant  $\int_{B_1} |z|^{-d+2s} dz$  and it is dominated by the latter. Passing to the limit as  $n \rightarrow \infty$  in (4.34), we get the assertion by dominated convergence (recall that  $\nu$  is finite).  $\square$

*Proof of Theorem 3.3.* Consider a function  $u$  satisfying (3.1), (3.2) and (3.3). Monotonicity in time of the associated potential is proved as we did after (4.22): notice that, for such an argument to work, the running assumptions on  $\gamma$  are required. The same proof holds if, instead of  $u \in L^\infty(\mathbb{R}^d \times (\tau, \infty))$ ,  $u$  is only supposed to satisfy  $\int_{t_1}^{t_2} u^m(\cdot, \tau) d\tau \in L_\rho^1(\mathbb{R}^d)$  for all  $t_2 > t_1 > 0$ . Existence of an initial trace  $\mu$ , meant as convergence in  $\sigma(\mathcal{M}(\mathbb{R}^d), C_c(\mathbb{R}^d))$  along subsequences of a given sequence of times tending to  $t = 0$ , follows by compactness, since we are assuming that solutions belong to  $L^\infty((0, \infty); L_\rho^1(\mathbb{R}^d))$ . Uniqueness of such a trace is established proceeding as we did after (4.29), using the monotonicity of potentials and the results of [30].

We are left with proving that convergence to  $\mu$  takes places also in  $\sigma(\mathcal{M}(\mathbb{R}^d), C_b(\mathbb{R}^d))$ , namely that  $\text{ess lim}_{t \rightarrow 0} \int_{\mathbb{R}^d} u(x, t) \rho(x) dx = \mu(\mathbb{R}^d)$ . By weak\* lower semi-continuity, it is then enough to show that  $\text{ess lim sup}_{t \rightarrow 0} \int_{\mathbb{R}^d} u(x, t) \rho(x) dx \leq \mu(\mathbb{R}^d)$ . Let  $U(\cdot, t)$  be the potential of  $\{\rho(\cdot)u(\cdot, t)\}$ . Again, the monotonicity in time of  $U(\cdot, t)$  and the first part of the proof ensure that  $U^\mu - U(\cdot, t) \geq 0$  almost everywhere. Therefore, Lemma 4.6 applied to the signed finite measure  $d\nu = d\mu - u(x, t)\rho(x)dx$  entails  $\mu(\mathbb{R}^d) \geq \int_{\mathbb{R}^d} u(x, t) \rho(x) dx$ . Letting  $t \rightarrow 0$  concludes the proof.  $\square$

**4.5. Strong solutions and decrease of the norms.** In order to justify rigorously some of the above computations, it is essential to show that the weak solutions constructed in Lemma 4.2 are strong. By a ‘‘strong solution’’, following [19, Section 6.2], we mean a weak solution  $u$  such that  $u_t \in L^\infty((\tau, \infty), L_\rho^1(\mathbb{R}^d))$  for all  $\tau > 0$ . The fact that our solutions are indeed strong could be proved as in [19, Section 8.1]. The first step consists in showing that  $\rho(\cdot)u_t(\cdot, t)$  is a bounded Radon measure which satisfies the estimate

$$\|\rho(\cdot)u_t(\cdot, t)\|_{\mathcal{M}(\mathbb{R}^d)} \leq \frac{2\|u_0\|_{1,\rho}}{(m-1)t} \quad \forall t > 0, \quad (4.35)$$

where now, by  $\mathcal{M}(\mathbb{R}^d)$  we mean the Banach space of *signed* measures on  $\mathbb{R}^d$ , equipped with the usual norm of the variation. As in [45, Lemma 8.5], this follows using

$$\int_{\mathbb{R}^d} [u(x, t) - \tilde{u}(x, t)]_+ \rho(x) dx \leq \int_{\mathbb{R}^d} [u_0(x) - \tilde{u}_0(x)]_+ \rho(x) dx \quad \forall t > 0, \quad (4.36)$$

where  $u$  and  $\tilde{u}$  are the solutions to (4.3) *constructed in Lemma 4.2* corresponding to the initial data  $u_0$  and  $\tilde{u}_0$ , respectively. Such principle does hold for the approximate solutions  $u_\eta$  and  $\tilde{u}_\eta$  used in the proof of Lemma 4.2 (see [40, Proposition 3.4]), whence (4.36) follows by passing to the limit. Afterwards, as [19, Lemma 8.1], one proves that  $z := u^{\frac{m+1}{2}}$  fulfills (3.7). In particular,

$$z_t \in L_{loc}^2((0, \infty); L_\rho^2(\mathbb{R}^d)). \quad (4.37)$$

Thanks to (4.35) and (4.37), the abstract result contained in [7, Theorem 1.1] ensures that  $u_t \in L_{loc}^1((0, \infty); L_\rho^1(\mathbb{R}^d))$ . In particular, (4.35) holds true with  $\|\rho(\cdot)u_t(\cdot, t)\|_{\mathcal{M}(\mathbb{R}^d)}$  replaced by  $\|u_t(t)\|_{1, \rho}$ , whence the assertion.

An important consequence of the fact that the solutions constructed in Lemma 4.2 are strong is the *decrease of their  $L_\rho^p$  norms* for any  $p \in [1, \infty]$ . Indeed, by definition of strong solution, for  $p \in (1, \infty)$ , we are allowed to multiply the differential equation in (4.3) by  $u^{p-1}$  and integrate in  $\mathbb{R}^d \times [t_1, t_2]$ . By Stroock-Varopoulos inequality (4.9) (let  $v = u^m$  and  $q = (p + m - 1)/m$ ), we get

$$\int_{\mathbb{R}^d} u^p(x, t_2) \rho(x) dx - \int_{\mathbb{R}^d} u^p(x, t_1) \rho(x) dx = -p \int_{t_1}^{t_2} \int_{\mathbb{R}^d} u^{p-1}(x, t) (-\Delta)^s (u^m)(x, t) dx dt \leq 0 \quad (4.38)$$

for all  $t_2 > t_1 > 0$ . The validity of (4.38) down to  $t_1 = 0$  follows using the approximate solutions  $\{u_\eta\}$  from the proof of Lemma 4.2 and letting  $\eta \rightarrow 0$ . The case  $p = \infty$  can be handled by approximation.

## 5. UNIQUENESS OF WEAK SOLUTIONS

Prior to the proof of Theorem 3.4, we need some technical lemmas. Hereafter, by “weak solution” to (1.1), we shall mean a solution in the sense of Definition 3.1.

**Lemma 5.1.** *Let  $d > 2s$  and assume that  $\rho$  satisfies (1.2) for some  $\gamma \in [0, 2s) \cap [0, d - 2s]$  and  $\gamma_0 \in [0, \gamma]$ . Let  $u$  be a weak solution to (1.1). Then the potential  $U(\cdot, t)$  of  $\rho(\cdot)u(\cdot, t)$  admits an absolutely continuous version (in  $L_{loc}^1(\mathbb{R}^d)$ ) which is nonincreasing in  $t$ .*

*Proof.* One proceeds as in the first part of the proof of Theorem 3.2, using the same technique we exploited to prove (4.22) rigorously.  $\square$

**Lemma 5.2.** *Let  $d > 2s$  and assume that  $\rho$  satisfies (1.2) for some  $\gamma \in [0, 2s) \cap [0, d - 2s]$  and  $\gamma_0 \in [0, \gamma]$ . Let  $u$  be a weak solution to (1.1), starting from the initial datum  $\mu$  whose potential is  $U^\mu$ . Then there holds*

$$\lim_{t \downarrow 0} U(x, t) = U^\mu(x) \quad \forall x \in \mathbb{R}^d. \quad (5.1)$$

*Proof.* It is a direct application of Theorem 3.9 of [30] but, for the reader’s convenience, we give some details.

Thanks to Theorem 3.8 of [30] and to the monotonicity ensured by Lemma 5.1, we have that the limit in (5.1) is taken at least for a.e.  $x \in \mathbb{R}^d$ . However, for what follows it will be crucial to prove that it is taken *for every*  $x \in \mathbb{R}^d$ . To this end we make use again of the monotonicity property provided by Lemma 5.1. In fact, Lemma 1.12 of [30] shows that, as a consequence of the monotonicity of potentials, there exists a positive measure  $\nu$ , whose potential is denoted by  $U^\nu$ , and a constant  $A \geq 0$  such that

$$\lim_{t \downarrow 0} U(x, t) = U^\nu(x) + A \quad \forall x \in \mathbb{R}^d.$$

Since (5.1) holds almost everywhere,

$$U^\mu(x) = U^\nu(x) + A \quad \text{for a.e. } x \in \mathbb{R}^d. \quad (5.2)$$

But using the corollary at page 129 of [30], from (5.2) we deduce that necessarily  $A = 0$ . Hence, (5.2) implies that  $U^\nu = U^\mu$  almost everywhere, and from Theorem 1.12 of [30] we know that two potentials coinciding a.e. in fact coincide everywhere, whence (5.1) follows.  $\square$

**5.1. Main ideas in the proof of uniqueness.** Since the proof of Theorem 3.4 is rather delicate, we point out its main ingredients. We should note that from a general viewpoint it is based on a “duality method”, and in particular it is modeled on the uniqueness proof given by M. Pierre in [36]. We comment again that our uniqueness result seems to be new even if  $s = 1$ , in the weighted case, or if  $\rho \equiv 1$  when  $s \in (0, 1)$ .

Let  $u_1$  and  $u_2$  be two weak solutions to (1.1) such that they both take a common positive, finite measure  $\mu$  as initial datum. We denote as  $U_1(\cdot, t)$  and  $U_2(\cdot, t)$  the potentials of  $\rho(\cdot)u_1(\cdot, t)$  and  $\rho(\cdot)u_2(\cdot, t)$ , respectively. Fix once for all the parameters  $h, T > 0$  and consider the function

$$g(x, t) := U_2(x, t + h) - U_1(x, t) \quad \forall (x, t) \in \mathbb{R}^d \times (0, T]. \quad (5.3)$$

Proceeding again as in the proof of Theorem 3.2 (under the hypothesis  $\gamma \leq d - 2s$ , see the proof of (4.22)), we get that  $g(\cdot, t)$  is an absolutely continuous curve (for instance in  $L^1_{loc}(\mathbb{R}^d)$ ) satisfying

$$\rho(x)g_t(x, t) = \rho(x)(u_1^m(x, t) - u_2^m(x, t + h)) = -a(x, t)(-\Delta)^s(g)(x, t) \quad (5.4)$$

for a.e.  $(x, t) \in \mathbb{R}^d \times (0, T)$ , where we define the function  $a$  as

$$a(x, t) := \begin{cases} \frac{u_1^m(x, t) - u_2^m(x, t + h)}{u_1(x, t) - u_2(x, t + h)} & \text{if } u_1(x, t) \neq u_2(x, t + h), \\ 0 & \text{if } u_1(x, t) = u_2(x, t + h), \end{cases} \quad (5.5)$$

and we used the fact that, thanks to the properties of Riesz potentials,

$$(-\Delta)^s(g)(x, t) = \rho(x)u_2(x, t + h) - \rho(x)u_1(x, t).$$

Note that, since  $m > 1$  and  $u_1, u_2 \in L^\infty(\mathbb{R}^d \times (\tau, \infty))$  for all  $\tau > 0$ ,  $a$  is a nonnegative function belonging to  $L^\infty(\mathbb{R}^d \times (\tau, \infty))$  for all  $\tau > 0$ .

Hence  $g$  is a solution to the *linear fractional* equation (5.4). Moreover, by Lemmas 5.1 and 5.2,  $g(x, 0) \leq 0$  for a.e.  $x \in \mathbb{R}^d$ . If we could apply the maximum principle, then we would get  $g \leq 0$  in  $\mathbb{R}^d \times (0, \infty)$ . This would imply  $u_1 \leq u_2$  and, by swapping the roles of  $u_1, u_2$ ,  $u_1 \equiv u_2$ . However, a priori a maximum principle is not available for solutions to (5.4). We then consider the “dual” problem

$$\begin{cases} \rho(x)\varphi_t = (-\Delta)^s(a\varphi) & \text{in } \mathbb{R}^d \times (0, T), \\ \varphi(x, T) = \psi(x) & \text{on } \mathbb{R}^d \times \{T\}, \end{cases} \quad (5.6)$$

for any  $\psi \in \mathcal{D}_+(\mathbb{R}^d)$ . Suppose for a moment that it admits a unique smooth solution  $\varphi$ . Multiplying (5.4) by  $\varphi$  and integrating by parts we formally obtain

$$\int_{\mathbb{R}^d} g(x, T)\rho(x)\psi(x) dx = \int_{\mathbb{R}^d} \varphi(x, 0)g(x, 0) dx. \quad (5.7)$$

The conclusion would again follow should a maximum principle for (5.7) hold, and in order to justify its applicability rigorously a further approximation is necessary. In fact, for every  $n \in \mathbb{N}$  and  $\varepsilon > 0$ , we consider a family  $\{\psi_{n,\varepsilon}\}$  which solve, in a sense which will be clarified later, the problem

$$\begin{cases} \rho(x)(\psi_{n,\varepsilon})_t = (-\Delta)^s[(a_n + \varepsilon)\psi_{n,\varepsilon}] & \text{in } \mathbb{R}^d \times (0, T), \\ \psi_{n,\varepsilon} = \psi & \text{on } \mathbb{R}^d \times \{T\}, \end{cases} \quad (5.8)$$

where  $\psi \in \mathcal{D}_+(\mathbb{R}^d)$ . The sequence  $\{a_n\}$  is a suitable approximation of the function  $a$  defined in (5.5). In particular we suppose that, for every  $n \in \mathbb{N}$ ,  $a_n(x, t)$  is a piecewise constant function of  $t$  (regular in  $x$ ) on the time intervals  $(T - (k + 1)T/n, T - kT/n]$ , for any  $k \in \{0, \dots, n - 1\}$ . Thanks to Theorem 3.7 and to Proposition B.1 below, we are then able to treat problem (5.8) by means of standard semigroup theory. Here the Markov property for the linear semigroup associated to the operator  $A = \rho^{-1}(-\Delta)^s$  will have a crucial role. Let us mention that in [36, Theorem 1], where  $s = 1, \rho \equiv 1$ , in view of standard parabolic theory, it was not necessary to approximate the function

$a$  by a piecewise constant function of  $t$ . Using the family  $\psi_{n,\varepsilon}$  and passing to the limit as  $n \rightarrow \infty$  and then as  $\varepsilon \rightarrow 0$  we get the next crucial identity:

$$\int_{\mathbb{R}^d} g(x, T) \psi(x) \rho(x) dx = \int_{\mathbb{R}^d} g(x, t) d\nu(t) \quad \text{for a.e. } t \in (0, T), \quad (5.9)$$

where  $\{\nu(t)\}$  is a specific family of positive finite measures. More precisely,  $\nu(t)$  is the limit in  $\sigma(\mathcal{M}(\mathbb{R}^d), C_b(\mathbb{R}^d))$  as  $\varepsilon \rightarrow 0$  of  $\{\rho(\cdot) \psi_\varepsilon(\cdot, t)\}$ , where  $\psi_\varepsilon$  is in turn the weak limit in  $L^2_\rho(\mathbb{R}^d \times (\tau, T))$  (for all  $\tau \in (0, T)$ ) as  $n \rightarrow \infty$  of  $\{\psi_{n,\varepsilon}\}$ . Note that, roughly speaking, (5.9) corresponds to identity (5.7) in the previous formal argument. Finally, we shall rigorously prove that the right-hand side of (5.9) has a nonpositive limit as  $t \rightarrow 0$ , whence the conclusion follows.

**5.2. Construction and properties of the family  $\{\psi_{n,\varepsilon}\}$ .** We begin our proof by introducing the functions  $\psi_{n,\varepsilon}$ , which formally solve (5.8).

**Lemma 5.3.** *Let  $d > 2s$  and assume that  $\rho$  satisfies (1.2) for some  $\gamma \in (0, 2s)$ . Let  $\{a_n\}$  be a sequence of functions converging a.e. to the function  $a$  as in (5.5) such that:*

- for any  $n \in \mathbb{N}$  and  $t > 0$ ,  $a_n(x, t)$  is a regular function of  $x$ ;
- for any  $n \in \mathbb{N}$  and  $x \in \mathbb{R}^d$ ,  $a_n(x, t)$  is a piecewise constant function of  $t$  on the time intervals  $(T - (k+1)T/n, T - kT/n]$ , for any  $k \in \{0, \dots, n-1\}$ ;
- $\{\|a_n\|_{L^\infty(\mathbb{R}^d \times (\tau, \infty))}\}$  is uniformly bounded in  $n$  for any  $\tau > 0$ .

Then, for any  $\varepsilon > 0$  and any  $\psi \in \mathcal{D}_+(\mathbb{R}^d)$ , there exists a nonnegative solution  $\psi_{n,\varepsilon}$  to problem (5.8), in the sense that  $\psi_{n,\varepsilon}(\cdot, t)$  is a continuous curve in  $L^p_\rho(\mathbb{R}^d)$  (for all  $p \in (1, \infty)$ ) satisfying  $\psi_{n,\varepsilon}(\cdot, 0) = \psi(\cdot, 0)$  and it is absolutely continuous on  $(T - (k+1)T/n, T - kT/n)$  for all  $k \in \{0, \dots, n-1\}$ , so that the identity

$$\begin{aligned} \psi_{n,\varepsilon}(\cdot, t_2) - \psi_{n,\varepsilon}(\cdot, t_1) &= \int_{t_1}^{t_2} \rho^{-1}(\cdot) (-\Delta)^s [(a_n + \varepsilon) \psi_{n,\varepsilon}](\cdot, \tau) d\tau \\ \forall t_1, t_2 &\in \left( T - \frac{(k+1)T}{n}, T - \frac{kT}{n} \right), \quad \forall k \in \{0, \dots, n-1\} \end{aligned} \quad (5.10)$$

holds true in  $L^p_\rho(\mathbb{R}^d)$  for all  $p \in (1, \infty)$ . Moreover,

$$\psi_{n,\varepsilon} \in L^\infty((0, T); L^p_\rho(\mathbb{R}^d)) \quad \forall p \in [1, \infty] \quad \text{and} \quad \|\psi_{n,\varepsilon}(t)\|_{1,\rho} \leq \|\psi\|_{1,\rho} \quad \forall t \in [0, T]. \quad (5.11)$$

*Proof.* To construct  $\psi_{n,\varepsilon}$  as in the statement, we first define  $\zeta_1$  as the solution of

$$\begin{cases} \rho(x) (\zeta_1)_t = (-\Delta)^s [(a_n(T) + \varepsilon) \zeta_1] & \text{in } \mathbb{R}^d \times (T - \frac{T}{n}, T), \\ \zeta_1 = \psi & \text{on } \mathbb{R}^d \times \{T\}. \end{cases} \quad (5.12)$$

To obtain  $\zeta_1$ , one can for instance exploit the change of variable

$$\phi_1(x, t) := (a_n(x, T) + \varepsilon) \zeta_1(x, t) \quad (5.13)$$

and consider  $\phi_1$  as the solution of

$$\begin{cases} (\phi_1)_t = (a_n(T) + \varepsilon) \rho^{-1}(-\Delta)^s(\phi_1) & \text{in } \mathbb{R}^d \times (T - \frac{T}{n}, T), \\ \phi_1 = (a_n(T) + \varepsilon) \psi & \text{on } \mathbb{R}^d \times \{T\}. \end{cases} \quad (5.14)$$

Problem (5.14) is indeed solvable by standard semigroup theory. In fact, consider the operator  $A_1 := \rho_1^{-1}(-\Delta)^s$  where we have set  $\rho_1(x) := (a_n(x, T) + \varepsilon)^{-1} \rho(x)$ , and the domain of  $L_1$  is  $X_{s,\rho_1} = X_{s,\rho}$  (see Definition 3.6).  $L_1$  is positive, self-adjoint and generates a Markov semigroup on  $L^2_{\rho_1}(\mathbb{R}^d)$ . These properties follow from Theorem 3.7. Our initial datum  $\phi_1$  belongs to  $L^p_{\rho_1}(\mathbb{R}^d)$  for all  $p \in [1, \infty]$ , and this is enough in order to have a solution to (5.14) which is continuous up to  $t = T$  and absolutely continuous in  $(T - \frac{T}{n}, T)$  in  $L^p_{\rho_1}(\mathbb{R}^d)$  for all  $p \in (1, \infty)$ . In fact, as recalled, the semigroup associated to  $A_1$  enjoys the Markov property and therefore, as a consequence of [14, Theorems 1.4.1, 1.4.2], it is extendible to a contraction semigroup on  $L^p_{\rho_1}(\mathbb{R}^d)$  (consistent with the original semigroup on  $L^2_{\rho_1}(\mathbb{R}^d) \cap L^p_{\rho_1}(\mathbb{R}^d)$ ) for all  $p \in [1, \infty]$ , which is analytic with a suitable angle  $\theta_p > 0$  if  $p \in (1, \infty)$ . By

classical results (see e.g. [35, Theorem 5.2 at p. 61]) the latter property ensures in particular that problem (5.14) is solved by a *differentiable* curve  $\phi_1(\cdot, t)$  in  $L^p_{\rho_1}(\mathbb{R}^d)$  for all  $p \in (1, \infty)$ . Going back to the original variable  $\zeta_1$  through (5.13), we deduce that it solves (5.12) in the same sense in which  $\phi_1$  solves (5.14). Having at our disposal such a  $\zeta_1$ , we can then solve the problem

$$\begin{cases} \rho(x) (\zeta_2)_t = (-\Delta)^s [(a_n(T - \frac{T}{n}) + \varepsilon) \zeta_2] & \text{in } \mathbb{R}^d \times (T - \frac{2T}{n}, T - \frac{T}{n}), \\ \zeta_2 = (a_n(x, T) + \varepsilon)^{-1} \phi_1 & \text{on } \mathbb{R}^d \times \{T - \frac{T}{n}\}, \end{cases}$$

just by proceeding as above. That is, we perform the change of variable

$$\phi_2(x, t) := \left( a_n \left( x, T - \frac{T}{n} \right) + \varepsilon \right) \zeta_2(x, t)$$

and take  $\phi_2$  as the solution of

$$\begin{cases} (\phi_2)_t = (a_n(T - \frac{T}{n}) + \varepsilon) \rho^{-1}(-\Delta)^s(\phi_2) & \text{in } \mathbb{R}^d \times (T - \frac{2T}{n}, T - \frac{T}{n}), \\ \phi_2 = (a_n(T - \frac{T}{n}) + \varepsilon) \zeta_1 = \frac{(a_n(T - \frac{T}{n}) + \varepsilon)}{(a_n(T) + \varepsilon)} \phi_1 & \text{on } \mathbb{R}^d \times \{T - \frac{T}{n}\}. \end{cases}$$

It is clear how the procedure goes on and allows to obtain a solution  $\psi_{n,\varepsilon}$  to (5.8) in the sense of the statement, just by defining it as

$$\psi_{n,\varepsilon}(\cdot, t) := \zeta_{k+1}(\cdot, t) \quad \forall t \in \left( T - \frac{(k+1)T}{n}, T - \frac{kT}{n} \right], \quad \forall k \in \{0, \dots, n-1\}.$$

Finally, since

$$\rho_{k+1}^{-1}(-\Delta)^s$$

generates a contraction semigroup on  $L^p_{\rho_{k+1}}(\mathbb{R}^d)$  for all  $p \in [1, \infty]$ , where

$$\rho_{k+1}(x) := \left( a_n \left( x, T - \frac{kT}{n} \right) + \varepsilon \right)^{-1} \rho(x), \quad (5.15)$$

the inequalities

$$\begin{aligned} \|\phi_{k+1}(t)\|_{p, \rho_{k+1}} &\leq \left\| \frac{(a_n(T - \frac{kT}{n}) + \varepsilon)}{(a_n(T - \frac{(k-1)T}{n}) + \varepsilon)} \phi_k \left( T - \frac{kT}{n} \right) \right\|_{p, \rho_{k+1}} \\ &\forall t \in \left( T - \frac{(k+1)T}{n}, T - \frac{kT}{n} \right], \quad \forall p \in [1, \infty] \end{aligned} \quad (5.16)$$

hold true for any  $k \in \{0, \dots, n-1\}$  (on the r.h.s. of (5.16) for  $k=0$  we conventionally set  $\phi_0 = \psi$  and  $a_n(T + T/n) + \varepsilon = 1$ ). Going back to the variables  $\zeta_{k+1}$  and recalling (5.15), from (5.16) one deduces (5.11): in fact, for  $p=1$  it is easy to see that the terms containing  $a_n$  cancel out and give the corresponding inequality, while for  $p > 1$  such terms remain and one obtains an inequality of the type  $\|\psi_{n,\varepsilon}(t)\|_{p,\rho} \leq C(n,\varepsilon)\|\psi\|_{p,\rho}$ , where  $C(n,\varepsilon)$  is a positive constant depending on  $n,\varepsilon$ .  $\square$

**Lemma 5.4.** *Let  $d > 2s$  and assume that  $\rho$  satisfies (1.2) for some  $\gamma \in (0, 2s) \cap (0, d - 2s]$ . Let  $g$  be as in (5.3),  $a$  as in (5.5) and  $a_n, \psi_{n,\varepsilon}, \psi$  as in Lemma 5.3. Then the identity*

$$\begin{aligned} &\int_{\mathbb{R}^d} g(x, T) \psi(x) \rho(x) dx - \int_{\mathbb{R}^d} g(x, t) \psi_{n,\varepsilon}(x, t) \rho(x) dx \\ &= \int_t^T \int_{\mathbb{R}^d} (a_n(x, \tau) + \varepsilon - a(x, \tau)) (-\Delta)^s(g)(x, \tau) \psi_{n,\varepsilon}(x, \tau) dx d\tau \end{aligned} \quad (5.17)$$

holds true for all  $t \in (0, T]$ .



*Proof.* To begin with, let us set  $t_k := T(n-k)/n$  for all  $k \in \{0, \dots, n\}$ . Recall that, from Lemma 5.3,  $\psi_{n,\varepsilon}(\cdot, t)$  is a continuous curve in  $L_\rho^p(\mathbb{R}^d)$  on  $(0, T]$ , absolutely continuous on any interval  $(t_{k+1}, t_k)$  for  $k \in \{0, \dots, n-1\}$  and satisfying the differential equation in (5.8) on such intervals, for all  $p \in (1, \infty)$ . Moreover,  $g(\cdot, t)$  is an absolutely continuous curve in  $L_\rho^p(\mathbb{R}^d)$  on  $(0, T]$  for all  $p$  such that

$$p \in \left( \frac{d-\gamma}{d-2s}, \infty \right). \quad (5.18)$$

Since  $g(x, t)$  is a continuous function of  $x$  (recall Lemma A.6) and the weight  $\rho(x)$  is locally integrable, in order to prove that  $g(\cdot, t) \in L_\rho^p(\mathbb{R}^d)$  for all  $p$  as in (5.18) it suffices to show that  $g(\cdot, t) \in L_\rho^p(B_1^c)$ . To this end, still Lemma A.6 ensures that  $g(\cdot, t) \in L^p(\mathbb{R}^d)$  for all  $p$  satisfying (A.11): the latter property and Hölder inequality imply that  $g(\cdot, t) \in L_\rho^p(B_1^c)$  for all  $p$  as in (5.18).

The fact that  $g(\cdot, t)$  is also absolutely continuous in  $L_\rho^p(\mathbb{R}^d)$  on the time interval  $(0, T]$  is then a consequence of (5.4) and of the integrability properties of  $u_1, u_2$ . Hence, due to Lemma 5.3, we get that

$$t \rightarrow \int_{\mathbb{R}^d} g(x, t) \psi_{n,\varepsilon}(x, t) \rho(x) dx \quad (5.19)$$

is a continuous function on  $(0, T]$ , absolutely continuous on each interval  $(t_{k+1}, t_k)$  and satisfies

$$\begin{aligned} & \frac{d}{dt} \int_{\mathbb{R}^d} g(x, t) \psi_{n,\varepsilon}(x, t) \rho(x) dx \\ &= \int_{\mathbb{R}^d} \{-a(x, t)(-\Delta)^s(g)(x, t) \psi_{n,\varepsilon}(x, t) + g(x, t)(-\Delta)^s[(a_n + \varepsilon) \psi_{n,\varepsilon}](x, t)\} dx \end{aligned} \quad (5.20)$$

there. As we have just seen,  $g(\cdot, t) \in L_\rho^p(\mathbb{R}^d)$  for all  $p$  satisfying (5.18) and  $\rho^{-1}(\cdot)(-\Delta)^s(g)(\cdot, t) \in L_\rho^p(\mathbb{R}^d)$  for all  $p \in [1, \infty]$ . Moreover, as a consequence of Lemma 5.3, we have that  $(a_n(\cdot, t) + \varepsilon) \psi_{n,\varepsilon}(\cdot, t) \in L_\rho^p(\mathbb{R}^d)$  for all  $p \in [1, \infty]$  and  $\rho^{-1}(\cdot)(-\Delta)^s[(a_n(\cdot, t) + \varepsilon) \psi_{n,\varepsilon}(\cdot, t)] \in L_\rho^p(\mathbb{R}^d)$  for all  $p \in (1, \infty)$ . We are therefore in position to apply Proposition B.1 to the r.h.s. of (5.20) (note that the interval  $((d-\gamma)/(d-2s), \infty) \cap [2, 2(d-\gamma)/(d-2s))$  is not empty) to get that

$$\frac{d}{dt} \int_{\mathbb{R}^d} g(x, t) \psi_{n,\varepsilon}(x, t) \rho(x) dx = \int_{\mathbb{R}^d} (a_n(x, t) + \varepsilon - a(x, t)) (-\Delta)^s(g)(x, t) \psi_{n,\varepsilon}(x, t) dx. \quad (5.21)$$

But the r.h.s. of (5.21) is in  $L^1((\tau, T))$  for any  $\tau \in (0, T)$ , from which (5.19) is absolutely continuous on the whole of  $(0, T]$  and not only on  $(t_{k+1}, t_k)$ . Integrating (5.21) between  $t$  and  $T$  then yields (5.17).  $\square$

Now we prove a key ‘‘conservation of mass’’ property for  $\psi_{n,\varepsilon}$ .

**Lemma 5.5.** *Let  $d > 2s$  and assume that  $\rho$  satisfies (1.2) for some  $\gamma \in (0, 2s)$ . Let  $\psi_{n,\varepsilon}$  and  $\psi$  be as in Lemma 5.3. Then the  $L_\rho^1$  norm of  $\psi_{n,\varepsilon}(\cdot, t)$  is preserved, that is*

$$\int_{\mathbb{R}^d} \psi_{n,\varepsilon}(x, t) \rho(x) dx = \int_{\mathbb{R}^d} \psi(x) \rho(x) dx \quad \forall t \in (0, T]. \quad (5.22)$$

*Proof.* Multiplying (5.10) by any  $\varphi \in \mathcal{D}(\mathbb{R}^d)$  and integrating in  $\mathbb{R}^d$ , we obtain:

$$\begin{aligned} & \int_{\mathbb{R}^d} \psi_{n,\varepsilon}(x, t^*) \varphi(x) \rho(x) dx - \int_{\mathbb{R}^d} \psi_{n,\varepsilon}(x, t_*) \varphi(x) \rho(x) dx \\ &= \int_{\mathbb{R}^d} (-\Delta)^s(\varphi)(x) \left( \int_{t_*}^{t^*} (a_n(x, \tau) + \varepsilon) \psi_{n,\varepsilon}(x, \tau) d\tau \right) dx \end{aligned} \quad (5.23)$$

for all  $t_*, t^* \in (t_{k+1}, t_k)$ . Since the  $L_\rho^1$  norm of  $\psi_{n,\varepsilon}(\cdot, t)$  is controlled by the  $L_\rho^1$  norm of the final datum  $\psi$  (recall (5.11)), from (5.23) we get:

$$\left| \int_{\mathbb{R}^d} \psi_{n,\varepsilon}(x, t^*) \varphi(x) \rho(x) dx - \int_{\mathbb{R}^d} \psi_{n,\varepsilon}(x, t_*) \varphi(x) \rho(x) dx \right| \leq C |t^* - t_*| \|\psi\|_{1,\rho} \|\rho^{-1}(-\Delta)^s(\varphi)\|_\infty, \quad (5.24)$$

where  $C := \|a_n + \varepsilon\|_{L^\infty(\mathbb{R}^d \times (t_* \wedge t^*, T))}$  is a positive constant independent of  $n$  and  $\varepsilon$ . Replacing  $\varphi$  with the cut-off function  $\xi_R$  (defined in Lemma A.3) and estimating the r.h.s. of (5.24) as in the proof of Proposition 4.1 yields

$$\begin{aligned} & \left| \int_{\mathbb{R}^d} \psi_{n,\varepsilon}(x, t^*) \xi_R(x) \rho(x) dx - \int_{\mathbb{R}^d} \psi_{n,\varepsilon}(x, t_*) \xi_R(x) \rho(x) dx \right| \\ & \leq C |t^* - t_*| \|\psi\|_{1,\rho} c^{-1} (R^{-2s} + R^{-2s+\gamma}) \|(1 + |x|^\gamma)(-\Delta)^s(\xi)\|_\infty \end{aligned} \quad (5.25)$$

for all  $R > 0$  and  $t_*, t^* \in (t_{k+1}, t_k)$ ,  $c$  being as in (1.2). Recalling that  $\psi_{n,\varepsilon}(\cdot, t)$  is a continuous curve (for instance in  $L^2_\rho(\mathbb{R}^d)$ ) on  $(0, T]$ , we can extend the validity of (5.25) (and (5.24)) to any  $t_*, t^* \in (0, T]$ . By choosing  $t^* = T$  and letting  $R \rightarrow \infty$  in (5.25) we finally get (5.22).  $\square$

In the next lemma we introduce the Riesz potential of  $\rho(\cdot)\psi_{n,\varepsilon}(\cdot, t)$ , which will play a fundamental role below.

**Lemma 5.6.** *Let  $d > 2s$  and assume that  $\rho$  satisfies (1.2) for some  $\gamma \in (0, 2s) \cap (0, d - 2s]$ . Let  $a_n, \psi_{n,\varepsilon}$  and  $\psi$  be as in Lemma 5.3. We denote as  $H_{n,\varepsilon}(\cdot, t)$  the Riesz potential of  $\rho(\cdot)\psi_{n,\varepsilon}(\cdot, t)$ , that is*

$$H_{n,\varepsilon}(x, t) := [I_{2s} * (\rho(\cdot)\psi_{n,\varepsilon}(\cdot, t))](x) \quad \forall (x, t) \in \mathbb{R}^d \times (0, T].$$

Then  $H_{n,\varepsilon}(\cdot, t) \in \dot{H}^s(\mathbb{R}^d)$  and the identity

$$\|I_{2s} * (\rho\psi)\|_{\dot{H}^s}^2 = \|H_{n,\varepsilon}(t)\|_{\dot{H}^s}^2 + 2 \int_t^T \int_{\mathbb{R}^d} (a_n(x, \tau) + \varepsilon) \psi_{n,\varepsilon}^2(x, \tau) \rho(x) dx d\tau \quad (5.26)$$

holds true for all  $t \in (0, T]$ .

*Proof.* First notice that  $\rho^{-1}(\cdot)(-\Delta)^s(H_{n,\varepsilon})(\cdot, t) = \psi_{n,\varepsilon}(\cdot, t) \in L^p_\rho(\mathbb{R}^d)$  for all  $p \in [1, \infty]$  (recall (5.11)) and  $H_{n,\varepsilon}(\cdot, t) \in L^p_\rho(\mathbb{R}^d)$  for all  $p$  satisfying (5.18) (this can be proved exploiting Lemma A.6 exactly as in the proof of Lemma 5.4). Again, since the interval  $((d - \gamma)/(d - 2s), \infty) \cap [2, 2(d - \gamma)/(d - 2s)]$  is not empty, applying Proposition B.1 we get that  $H_{n,\varepsilon}(\cdot, t) \in \dot{H}^s(\mathbb{R}^d)$  and the identity

$$\|H_{n,\varepsilon}(t)\|_{\dot{H}^s}^2 = \int_{\mathbb{R}^d} H_{n,\varepsilon}(x, t) (-\Delta)^s(H_{n,\varepsilon})(x, t) dx = \int_{\mathbb{R}^d} H_{n,\varepsilon}(x, t) \psi_{n,\varepsilon}(x, t) \rho(x) dx \quad (5.27)$$

holds true. Thanks to the validity of the differential equation

$$(H_{n,\varepsilon})_t(x, t) = (a_n(x, t) + \varepsilon) \psi_{n,\varepsilon}(x, t) \quad \text{for a.e. } (x, t) \in \mathbb{R}^d \times (0, T), \quad (5.28)$$

which can be justified as we did for (5.4), taking the time derivative of (5.27) in the intervals  $(t_{k+1}, t_k)$ , using (5.28), (5.8) and again Proposition B.1, we obtain:

$$\frac{d}{dt} \|H_{n,\varepsilon}(t)\|_{\dot{H}^s}^2 = 2 \int_{\mathbb{R}^d} (a_n(x, t) + \varepsilon) \psi_{n,\varepsilon}^2(x, t) \rho(x) dx. \quad (5.29)$$

A priori, from (5.27), we have that  $\|H_{n,\varepsilon}(t)\|_{\dot{H}^s}^2$  is continuous on  $(0, T]$  and absolutely continuous only on  $(t_{k+1}, t_k)$ . However, the r.h.s. of (5.29) is in  $L^1((\tau, T))$  for any  $\tau \in (0, T)$ . Hence, (5.26) just follows by integrating (5.29) from  $t$  to  $T$ .  $\square$

**5.3. Passing to the limit as  $n \rightarrow \infty$ .** The goal of the next lemma is to show that, as  $n \rightarrow \infty$ ,  $\{\psi_{n,\varepsilon}\}$  suitably converges to a limit function  $\psi_\varepsilon$  that enjoys some crucial properties.

**Lemma 5.7.** *Let  $d > 2s$  and assume that  $\rho$  satisfies (1.2) for some  $\gamma \in (0, 2s) \cap (0, d - 2s]$ . Let  $u_1$  and  $u_2$  be two weak solutions to problem (1.1), taking the common positive finite measure  $\mu$  as initial datum. Let  $g$  be as in (5.3),  $a$  as in (5.5) and  $\psi_{n,\varepsilon}, \psi$  as in Lemma 5.3. Then, up to subsequences,  $\{\psi_{n,\varepsilon}\}$  converges weakly in  $L^2_\rho(\mathbb{R}^d \times (\tau, T))$  (for all  $\tau \in (0, T)$ ) to a suitable nonnegative function  $\psi_\varepsilon$  and  $\{\rho(\cdot)\psi_{n,\varepsilon}(\cdot, t)\}$  converges to  $\rho(\cdot)\psi_\varepsilon(\cdot, t)$  in  $\sigma(\mathcal{M}(\mathbb{R}^d), C_b(\mathbb{R}^d))$  for a.e.  $t \in (0, T)$ . Moreover,  $\psi_\varepsilon$  enjoys the following properties:*

$$\int_{\mathbb{R}^d} \psi_\varepsilon(x, t) \rho(x) dx = \int_{\mathbb{R}^d} \psi(x) \rho(x) dx, \quad (5.30)$$

$$\begin{aligned} & \int_{\mathbb{R}^d} \psi(x) \varphi(x) \rho(x) dx - \int_{\mathbb{R}^d} \psi_\varepsilon(x, t) \varphi(x) \rho(x) dx \\ &= \int_{\mathbb{R}^d} (-\Delta)^s(\varphi)(x) \left( \int_t^T (a(x, \tau) + \varepsilon) \psi_\varepsilon(x, \tau) d\tau \right) dx, \end{aligned} \quad (5.31)$$

$$\begin{aligned} & \left| \int_{\mathbb{R}^d} g(x, T) \psi(x) \rho(x) dx - \int_{\mathbb{R}^d} g(x, t) \psi_\varepsilon(x, t) \rho(x) dx \right| \\ & \leq \varepsilon (T - t) \|\psi\|_{1, \rho} \|u_2(\tau + h) - u_1(\tau)\|_{L^\infty(\mathbb{R}^d \times (t, T))} \end{aligned} \quad (5.32)$$

for a.e.  $t \in (0, T)$ , for any  $\varphi \in \mathcal{D}(\mathbb{R}^d)$ .

*Proof.* From (5.26) one gets that, up to subsequences,  $\{\psi_{n, \varepsilon}\}$  converges weakly in  $L^2_\rho(\mathbb{R}^d \times (\tau, T))$  (for all  $\tau \in (0, T)$ ) to a suitable  $\psi_\varepsilon$ . Moreover, thanks to the uniform boundedness of  $\{\rho(\cdot) \psi_{n, \varepsilon}(\cdot, t)\}$  in  $L^1(\mathbb{R}^d)$  (see (5.11)), for every  $t \in (0, T)$  there exists a subsequence (which a priori may depend on  $t$ ) such that  $\{\rho(\cdot) \psi_{n, \varepsilon}(\cdot, t)\}$  converges in  $\sigma(\mathcal{M}(\mathbb{R}^d), C_c(\mathbb{R}^d))$  to some positive, finite measure  $\nu(t)$  (recall the preliminary results of Section 2). We aim at identifying (at least for almost every  $t \in (0, T)$ )  $\nu(t)$  with  $\rho(\cdot) \psi_\varepsilon(\cdot, t)$ , so that a posteriori the subsequence does not depend on  $t$ . In order to do that, let  $t \in (0, T)$  be a Lebesgue point of  $\psi_\varepsilon(\cdot, t)$  (as a curve in  $L^1((\tau, T); L^2_\rho(\mathbb{R}^d))$ ). Taking any  $\varphi \in \mathcal{D}(\mathbb{R}^d)$  and using (5.24), we obtain:

$$\begin{aligned} & \left| \int_t^{t+\delta} \int_{\mathbb{R}^d} \psi_{n, \varepsilon}(x, \tau) \varphi(x) \rho(x) dx d\tau - \int_t^{t+\delta} \int_{\mathbb{R}^d} \psi_{n, \varepsilon}(x, t) \varphi(x) \rho(x) dx d\tau \right| \\ & \leq \int_t^{t+\delta} \left| \int_{\mathbb{R}^d} \psi_{n, \varepsilon}(x, \tau) \varphi(x) \rho(x) dx - \int_{\mathbb{R}^d} \psi_{n, \varepsilon}(x, t) \varphi(x) \rho(x) dx \right| d\tau \\ & \leq \int_t^{t+\delta} C(\tau - t) \|\psi\|_{1, \rho} \|\rho^{-1}(-\Delta)^s(\varphi)\|_\infty d\tau = \frac{\delta^2}{2} C \|\psi\|_{1, \rho} \|\rho^{-1}(-\Delta)^s(\varphi)\|_\infty \end{aligned} \quad (5.33)$$

for all  $\delta$  sufficiently small. Letting  $n \rightarrow \infty$  (up to subsequences) in (5.33) yields

$$\left| \int_t^{t+\delta} \int_{\mathbb{R}^d} \psi_\varepsilon(x, \tau) \varphi(x) \rho(x) dx d\tau - \delta \int_{\mathbb{R}^d} \varphi(x) d\nu(t) \right| \leq \frac{\delta^2}{2} C \|\psi\|_{1, \rho} \|\rho^{-1}(-\Delta)^s(\varphi)\|_\infty. \quad (5.34)$$

Dividing (5.34) by  $\delta$  and letting  $\delta \rightarrow 0$  one deduces that (recall that  $t$  is a Lebesgue point for  $\{\psi_\varepsilon(\cdot, t)\}$ )

$$\int_{\mathbb{R}^d} \psi_\varepsilon(x, t) \varphi(x) \rho(x) dx = \int_{\mathbb{R}^d} \varphi(x) d\nu(t),$$

which is valid for any  $\varphi \in \mathcal{D}(\mathbb{R}^d)$ , whence  $\psi_\varepsilon(x, t) \rho(x) dx = d\nu(t)$ .

We now prove the claimed properties of  $\psi_\varepsilon$ . Letting  $n \rightarrow \infty$  in (5.25) (with  $t^* = T$  and  $t_* = t$ ) and using the just proved convergence of  $\{\rho(\cdot) \psi_{n, \varepsilon}(\cdot, t)\}$  to  $\rho(\cdot) \psi_\varepsilon(\cdot, t)$  in  $\sigma(\mathcal{M}(\mathbb{R}^d), C_c(\mathbb{R}^d))$ , we get

$$\begin{aligned} & \left| \int_{\mathbb{R}^d} \psi(x) \xi_R(x) \rho(x) dx - \int_{\mathbb{R}^d} \psi_\varepsilon(x, t) \xi_R(x) \rho(x) dx \right| \\ & \leq C(T - t) \|\psi\|_{1, \rho} c^{-1} (R^{-2s} + R^{-2s+\gamma}) \|(1 + |x|^\gamma)(-\Delta)^s(\xi)\|_\infty \end{aligned} \quad (5.35)$$

for a.e.  $t \in (0, T)$ ,  $c$  being as in (1.2). Letting  $R \rightarrow \infty$  in (5.35) we deduce (5.30). Thanks to (5.22) and (5.30) we infer in particular that

$$\lim_{n \rightarrow \infty} \|\psi_{n, \varepsilon}(t)\|_{1, \rho} = \|\psi_\varepsilon(t)\|_{1, \rho},$$

so that the convergence of  $\{\rho(\cdot)\psi_{n,\varepsilon}(\cdot, t)\}$  to  $\rho(\cdot)\psi_\varepsilon(\cdot, t)$  takes place also in  $\sigma(\mathcal{M}(\mathbb{R}^d), C_b(\mathbb{R}^d))$ . Recalling that  $g(\cdot, t)$  belongs to  $C_b(\mathbb{R}^d)$  (Lemma A.6), we can let  $n \rightarrow \infty$  in (5.17) to obtain

$$\begin{aligned} & \int_{\mathbb{R}^d} g(x, T)\psi(x)\rho(x)dx - \int_{\mathbb{R}^d} g(x, t)\psi_\varepsilon(x, t)\rho(x)dx \\ &= \lim_{n \rightarrow \infty} \left( \int_t^T \int_{\mathbb{R}^d} (a_n(x, \tau) + \varepsilon - a(x, \tau)) (-\Delta)^s(g)(x, \tau) \psi_{n,\varepsilon}(x, \tau) dx d\tau \right) \\ &= \lim_{n \rightarrow \infty} \left( \int_t^T \int_{\mathbb{R}^d} (a_n(x, \tau) + \varepsilon - a(x, \tau)) (u_2(x, \tau + h) - u_1(x, \tau)) \psi_{n,\varepsilon}(x, \tau) \rho(x) dx d\tau \right) \\ &= \varepsilon \int_t^T \int_{\mathbb{R}^d} (u_2(x, \tau + h) - u_1(x, \tau)) \psi_\varepsilon(x, \tau) \rho(x) dx d\tau \quad \text{for a.e. } t \in (0, T), \end{aligned} \tag{5.36}$$

where in the last integral we can pass to the limit since  $\{\psi_{n,\varepsilon}\}$  tends to  $\psi_\varepsilon$  weakly in  $L^2_\rho(\mathbb{R}^d \times (t, T))$ ,  $\{a_n\}$  tends to  $a$  in  $L^\infty(\mathbb{R}^d \times (t, T))$  and  $u_1, u_2$  belong to  $L^p_\rho(\mathbb{R}^d \times (t, T + h))$  for all  $p \in [1, \infty]$ . In particular, from (5.36) and (5.30) we get (5.32). Notice that, in a similarly way, we can pass to the limit in (5.23) (which actually holds for any  $t_*, t^* \in (0, T)$ ) and obtain (5.31).  $\square$

**5.4. Passing to the limit as  $\varepsilon \rightarrow 0$  and proof of Theorem 3.4.** We are now in position to prove Theorem 3.4, using the strategy of [36]: we give some detail for the reader's convenience.

*Proof of Theorem 3.4.* To begin with, we introduce the Riesz potential  $H_\varepsilon(\cdot, t)$  of  $\rho(\cdot)\psi_\varepsilon(\cdot, t)$ . Since we only know that  $\rho(\cdot)\psi_\varepsilon(\cdot, t) \in L^1(\mathbb{R}^d)$ , we have no information over the integrability of  $H_\varepsilon(\cdot, t)$  other than  $L^1_{loc}(\mathbb{R}^d)$  (by classical results, see e.g. [30, p. 61]). However, exploiting (5.31) and proceeding once again as in the proof of (4.22), we obtain

$$I_{2s} * (\rho\psi) - H_\varepsilon(\cdot, t) = \int_t^T (a(\cdot, \tau) + \varepsilon) \psi_\varepsilon(\cdot, \tau) d\tau \geq 0 \quad \text{for a.e. } t \in (0, T),$$

whence, in particular,

$$0 \leq H_\varepsilon(x, t_1) \leq H_\varepsilon(x, t_2) \leq H_\varepsilon(x, T) = I_{2s} * (\rho\psi)(x) \tag{5.37}$$

for a.e.  $0 < t_1 \leq t_2 \leq T$  and a.e.  $x \in \mathbb{R}^d$ . The above inequality shows that  $H_\varepsilon(\cdot, t)$  belongs to  $L^p(\mathbb{R}^d)$  at least for the same  $p$  for which  $H_\varepsilon(\cdot, T)$  does, namely for any  $p \in (d/(d-2), \infty]$ .

Our next goal is to let  $\varepsilon \rightarrow 0$  (along a fixed sequence whose index for the moment we omit, in order to simplify readability). Thanks to the boundedness of  $\{\rho(\cdot)\psi_\varepsilon(\cdot, t)\}$  in  $L^1(\mathbb{R}^d)$  (trivial consequence of (5.30)), for a.e.  $t \in (0, T)$  there exists a subsequence  $\{\varepsilon_n\}$  (a priori depending on  $t$ ) such that  $\{\rho(\cdot)\psi_\varepsilon(\cdot, t)\}$  converges to a positive finite measure  $\nu(t)$  in  $\sigma(\mathcal{M}(\mathbb{R}^d), C_c(\mathbb{R}^d))$ . In order to overcome the possible dependence of  $\{\varepsilon_n\}$  on  $t$ , we exploit the properties of  $\{H_\varepsilon\}$ . First notice that (5.37) ensures the uniform boundedness of  $\{H_\varepsilon\}$  in  $L^p(\mathbb{R}^d \times (0, T))$  for any  $p \in (d/(d-2), \infty]$ . This entails the existence of a decreasing subsequence  $\{\varepsilon_m\}$  such that  $\{H_{\varepsilon_m}\}$  converges weakly in  $L^p(\mathbb{R}^d \times (0, T))$  to a suitable limit  $H$ . Mazur's Lemma implies that there exists a sequence  $\{H_k\}$  of convex combinations of  $\{H_{\varepsilon_m}\}$  that converges *strongly* to  $H$  in  $L^p(\mathbb{R}^d \times (0, T))$ . By definition, the sequence  $\{H_k\}$  is of the form

$$H_k = \sum_{m=1}^{M_k} \lambda_{m,k} H_{\varepsilon_m}, \quad \sum_{m=1}^{M_k} \lambda_{m,k} = 1$$

for some sequence  $\{M_k\} \subset \mathbb{N}$  and a suitable choice of the coefficients  $\lambda_{m,k} \in [0, 1]$ . With no loss of generality we shall also assume that

$$\lim_{k \rightarrow \infty} \left( \sum_{m=1}^{M_k} \varepsilon_m \lambda_{m,k} \right) = 0.$$

This can be justified by applying iteratively Mazur's Lemma on suitable subsequences of  $\{H_{\varepsilon_m}\}$ . Now notice that the function whose Riesz potential is  $H_k$  is

$$f_k(x, t) = \sum_{m=1}^{M_k} \lambda_{m,k} \rho(x) \psi_{\varepsilon_m}(x, t).$$

Multiplying (5.32) (with  $\varepsilon = \varepsilon_m$ ) by  $\lambda_{m,k}$  and summing over  $k$ , one gets that  $f_k$  satisfies

$$\begin{aligned} & \left| \int_{\mathbb{R}^d} g(x, T) \psi(x) \rho(x) dx - \int_{\mathbb{R}^d} g(x, t) f_k(x, t) dx \right| \\ & \leq \left( \sum_{m=1}^{M_k} \varepsilon_m \lambda_{m,k} \right) (T - t) \|\psi\|_{1,\rho} \|u_2(\tau + h) - u_1(\tau)\|_{L^\infty(\mathbb{R}^d \times (t, T))} \end{aligned} \quad (5.38)$$

for a.e.  $t \in (0, T)$ , whereas from (5.30) and (5.35) we infer that

$$\begin{aligned} & \left| \int_{\mathbb{R}^d} \psi(x) \xi_R(x) \rho(x) dx - \int_{\mathbb{R}^d} f_k(x, t) \xi_R(x) dx \right| \\ & \leq C(T - t) \|\psi\|_{1,\rho} c^{-1} (R^{-2s} + R^{-2s+\gamma}) \|(1 + |x|^\gamma)(-\Delta)^s(\xi)\|_\infty \end{aligned} \quad (5.39)$$

for a.e.  $t \in (0, T)$  and

$$\int_{\mathbb{R}^d} \psi(x) \rho(x) dx = \int_{\mathbb{R}^d} f_k(x, t) dx \quad \text{for a.e. } t \in (0, T). \quad (5.40)$$

Letting  $k \rightarrow \infty$  we find that, for a.e.  $t \in (0, T)$ , there exists a subsequence of  $\{f_k(\cdot, t)\}$  (a priori depending on  $t$ ) that converges in  $\sigma(\mathcal{M}(\mathbb{R}^d), C_c(\mathbb{R}^d))$  to a positive, finite measure  $\nu(t)$ . But the fact that  $\{H_k\}$  converges strongly in  $L^p(\mathbb{R}^d \times (0, T))$  to  $H$  forces the potential of  $\nu(t)$  to coincide a.e. with  $H(\cdot, t)$ . This is a consequence of [30, Theorem 3.8]. By [30, Theorem 1.12] we therefore deduce that the limit  $\nu(t)$  is uniquely determined by its potential  $H(\cdot, t)$ . This identification allows to assert that for a.e.  $t \in (0, T)$  the *whole* sequence  $\{f_k(\cdot, t)\}$  converges to  $\nu(t)$  in  $\sigma(\mathcal{M}(\mathbb{R}^d), C_c(\mathbb{R}^d))$ .

Passing to the limit in (5.37) (after having set  $\varepsilon = \varepsilon_m$ , multiplied by  $\lambda_{m,k}$  and summed over  $k$ ) we deduce that also the potentials  $H(\cdot, t)$  of  $\nu(t)$  are ordered and bounded above by  $I_{2s} * (\rho\psi)$ :

$$0 \leq H(x, t_1) \leq H(x, t_2) \leq I_{2s} * (\rho\psi)(x) \quad \text{for a.e. } 0 < t_1 \leq t_2 \leq T, \text{ for a.e. } x \in \mathbb{R}^d. \quad (5.41)$$

Letting  $k \rightarrow \infty$  in (5.39) yields

$$\begin{aligned} & \left| \int_{\mathbb{R}^d} \psi(x) \xi_R(x) \rho(x) dx - \int_{\mathbb{R}^d} \xi_R(x) d\nu(t) \right| \\ & \leq C(T - t) \|\psi\|_{1,\rho} c^{-1} (R^{-2s} + R^{-2s+\gamma}) \|(1 + |x|^\gamma)(-\Delta)^s(\xi)\|_\infty \end{aligned} \quad (5.42)$$

for a.e.  $t \in (0, T)$ , whence, letting  $R \rightarrow \infty$  in (5.42), we obtain:

$$\int_{\mathbb{R}^d} \psi(x) \rho(x) dx = \int_{\mathbb{R}^d} d\nu(t) \quad \text{for a.e. } t \in (0, T). \quad (5.43)$$

Gathering (5.40) and (5.43) we infer that  $\{f_k(\cdot, t)\}$  converges to  $\nu(t)$  also in  $\sigma(\mathcal{M}(\mathbb{R}^d), C_b(\mathbb{R}^d))$ : this allows to pass to the limit in (5.38) to get the identity (by exploiting (5.4) as well) (5.9). As a consequence of the monotonicity given by (5.41) and thanks to (5.42)-(5.9), the curve  $\nu(t)$  can be extended to *every*  $t \in (0, T]$  so that it still satisfies (5.41)-(5.9) (one uses again [30, Theorem 3.8]). Recalling that  $g(x, t) = U_2(x, t + h) - U_1(x, t)$  and that potentials do not increase in time (Lemma 5.1), we have that  $g(x, t) \leq U_2(x, h) - U_1(x, t_0)$  holds for all  $x \in \mathbb{R}^d$  and all  $t_0 > t$ . Because  $\nu(t)$  is a positive finite measure, this fact and (5.9) imply that

$$\int_{\mathbb{R}^d} g(x, T) \psi(x) \rho(x) dx \leq \int_{\mathbb{R}^d} (U_2(x, h) - U_1(x, t_0)) d\nu(t) \quad \forall t_0 > t. \quad (5.44)$$

Our next goal is to let  $t$  tend to zero in (5.44). Since the mass of  $\nu(t)$  is constant (formula (5.43)), up to subsequences  $\nu(t)$  converges to a suitable positive finite measure  $\nu$  in  $\sigma(\mathcal{M}(\mathbb{R}^d), C_c(\mathbb{R}^d))$ . Moreover, by (5.41), we know that the potentials  $H(\cdot, t)$  of  $\nu(t)$  are nondecreasing in  $t$  (for a.e.

$x$ ): in particular,  $H(\cdot, t)$  admits a pointwise limit almost everywhere  $H_0$  as  $t \downarrow 0$ . Theorem 3.8 of [30] ensures that  $H_0$  coincides almost everywhere with the potential of the limit measure  $\nu$  (which therefore does not depend on the subsequence). We can then pass to the limit in the integral

$$\int_{\mathbb{R}^d} U_1(x, t_0) d\nu(t). \quad (5.45)$$

Indeed, by Fubini's Theorem (5.45) is equal to

$$\int_{\mathbb{R}^d} u_1(x, t_0) H(x, t) \rho(x) dx. \quad (5.46)$$

Passing to the limit in (5.46) as  $t \downarrow 0$  we get that

$$\lim_{t \downarrow 0} \int_{\mathbb{R}^d} u_1(x, t_0) H(x, t) \rho(x) dx = \int_{\mathbb{R}^d} u_1(x, t_0) H_0(x) \rho(x) dx \quad (5.47)$$

by dominated convergence. Recalling that  $H_0$  is the potential of  $\nu$ , and using again Fubini's Theorem, (5.47) can be rewritten as

$$\lim_{t \downarrow 0} \int_{\mathbb{R}^d} U_1(x, t_0) d\nu(t) = \int_{\mathbb{R}^d} U_1(x, t_0) d\nu.$$

One proceeds similarly for the integral

$$\int_{\mathbb{R}^d} U_2(x, h) d\nu(t).$$

Hence, passing to the limit as  $t \downarrow 0$  in (5.44) yields

$$\int_{\mathbb{R}^d} g(x, T) \psi(x) \rho(x) dx \leq \int_{\mathbb{R}^d} (U_2(x, h) - U_1(x, t_0)) d\nu \quad \forall t_0 > 0. \quad (5.48)$$

Now we let  $t_0 \downarrow 0$  in (5.48). By monotone convergence (Lemmas 5.1 and 5.2) we obtain

$$\int_{\mathbb{R}^d} g(x, T) \psi(x) \rho(x) dx \leq \int_{\mathbb{R}^d} (U_2(x, h) - U^\mu(x)) d\nu \quad (5.49)$$

In this step it is crucial that the limit of  $U_1(x, t_0)$  to  $U^\mu(x)$  is taken *for every*  $x \in \mathbb{R}^d$  (Lemma 5.2), because we have no information over  $\nu$  besides the fact that it is a positive finite measure. Still by monotonicity we have that  $U_2(x, h) \leq U^\mu(x)$  for every  $x \in \mathbb{R}^d$ . Thus, from (5.49) it follows that

$$\int_{\mathbb{R}^d} g(x, T) \psi(x) \rho(x) dx \leq 0. \quad (5.50)$$

Since (5.50) holds true for any  $h, T > 0$  and any  $\psi \in \mathcal{D}_+(\mathbb{R}^d)$ , we infer that  $U_2 \leq U_1$ . Interchanging the role of  $u_1$  and  $u_2$  we get that  $U_1 \leq U_2$ , whence  $U_1 = U_2$  and  $u_1 = u_2$ .  $\square$

## APPENDIX A

We recall here some basic properties of the fractional Laplacian (and of a similar nonlocal, non-linear operator) of functions in  $\mathcal{D}(\mathbb{R}^d)$ . We omit the proofs of the first two lemmas, since they follow by exploiting the same strategy as in [6, Lemma 2.1].

**Lemma A.1.** *The  $s$ -Laplacian  $(-\Delta)^s(\phi)(x)$  of any  $\phi \in \mathcal{D}(\mathbb{R}^d)$  is a regular function which decays (together with its derivatives) at least like  $|x|^{-d-2s}$  as  $|x| \rightarrow \infty$ .*

**Lemma A.2.** *For any  $\phi \in \mathcal{D}(\mathbb{R}^d)$ , the function*

$$I_s(\phi)(x) := \int_{\mathbb{R}^d} \frac{(\phi(x) - \phi(y))^2}{|x - y|^{d+2s}} dy \quad \forall x \in \mathbb{R}^d$$

*is regular and decays (together with its derivatives) at least like  $|x|^{-d-2s}$  as  $|x| \rightarrow \infty$ .*

**Lemma A.3.** For any  $R > 0$ , let  $\xi_R$  be the cut-off function

$$\xi_R(x) := \xi\left(\frac{x}{R}\right) \quad \forall x \in \mathbb{R}^d,$$

where  $\xi(x)$  is a positive, regular function such that  $\|\xi\|_\infty \leq 1$ ,  $\xi = 1$  in  $B_1$  and  $\xi = 0$  in  $B_2^c$ . Then,  $(-\Delta)^s(\xi_R)$  and  $l_s(\xi_R)$  enjoy the following property:

$$(-\Delta)^s(\xi_R)(x) = \frac{1}{R^{2s}}(-\Delta)^s(\xi)\left(\frac{x}{R}\right), \quad l_s(\xi_R)(x) = \frac{1}{R^{2s}}l_s(\xi)\left(\frac{x}{R}\right) \quad \forall x \in \mathbb{R}^d.$$

*Proof.* We only prove the result for  $l_s(\xi_R)$ , since the proof for  $(-\Delta)^s(\xi_R)$  is identical. Letting  $\tilde{y} = y/R$ , one has:

$$l_s(\xi_R)(x) = \int_{\mathbb{R}^d} \frac{(\xi_R(x) - \xi_R(y))^2}{|x - y|^{d+2s}} dy = \frac{1}{R^{2s}} \int_{\mathbb{R}^d} \frac{(\xi(x/R) - \xi(\tilde{y}))^2}{|x/R - \tilde{y}|^{d+2s}} d\tilde{y} = \frac{1}{R^{2s}}l_s(\xi)\left(\frac{x}{R}\right).$$

□

The next lemmas contain technical ingredients concerning fractional Sobolev spaces and Riesz potentials, which we need in the proofs of our existence and uniqueness results.

**Lemma A.4.** Let  $d > 2s$  and assume that  $\rho$  satisfies (1.2) for some  $\gamma \in (0, d + 2s]$ . Consider a function  $v \in L^2_{loc}((0, \infty); \dot{H}^s(\mathbb{R}^d))$  such that, for all  $t_2 > t_1 > 0$ ,

$$\int_{t_1}^{t_2} \int_{\mathbb{R}^d} |v(x, t)|^2 \rho(x) dx dt \leq C, \quad (\text{A.1})$$

$$\int_{t_1}^{t_2} \int_{\mathbb{R}^d} |(-\Delta)^{\frac{s}{2}}(v)(x, t)|^2 dx dt \leq C \quad (\text{A.2})$$

and

$$\int_{t_1}^{t_2} \int_{\mathbb{R}^d} |v_t(x, t)|^2 \rho(x) dx dt \leq C, \quad (\text{A.3})$$

where  $C$  is a positive constant depending only on  $t_1$  and  $t_2$ . Take any cut-off functions  $\xi_1 \in C_c^\infty(\mathbb{R}^d)$ ,  $\xi_2 \in C_c^\infty((0, \infty))$  and define  $v_c : \mathbb{R}^d \rightarrow \mathbb{R}$  as follows:

$$v_c(x, t) := \xi_1(x)\xi_2(t)v(x, t) \quad \forall (x, t) \in \mathbb{R}^d \times \mathbb{R},$$

where we implicitly assume  $\xi_2$  and  $v$  to be zero for  $t < 0$ . Then

$$\|v_c\|_{\dot{H}^s(\mathbb{R}^{d+1})}^2 = \|v_c\|_{L^2(\mathbb{R}^{d+1})}^2 + \|v_c\|_{\dot{H}^s(\mathbb{R}^{d+1})}^2 \leq C' \quad (\text{A.4})$$

for a positive constant  $C'$  that depends only on  $\xi_1$  and  $\xi_2$  (also through  $C$ ).

*Proof.* The validity of

$$\|v_c\|_{L^2(\mathbb{R}^{d+1})}^2 \leq C' \quad (\text{A.5})$$

is an immediate consequence of (A.1) and of the fact that  $\rho$  is bounded away from zero on compact sets (from now on  $C'$  will be a constant as in the statement that we shall not relabel). Moreover, since  $(v_c)_t = \xi_1\xi_2'v + \xi_1\xi_2v_t$ , by (A.1), (A.3) and again the fact that  $\rho$  is bounded away from zero on compact sets we deduce that

$$\|(v_c)_t\|_{L^2(\mathbb{R}^{d+1})}^2 \leq C'. \quad (\text{A.6})$$

Now we have to handle the spatial regularity of  $v_c$ . Straightforward computations show that

$$\begin{aligned} \|v_c(t)\|_{\dot{H}^s(\mathbb{R}^d)}^2 &= C_{d,s} \xi_2^2(t) \int_{\mathbb{R}^d} \xi_1^2(x) \left( \int_{\mathbb{R}^d} \frac{(v(x, t) - v(y, t))^2}{|x - y|^{d+2s}} dy \right) dx \\ &\quad + C_{d,s} \xi_2^2(t) \int_{\mathbb{R}^d} |v(y, t)|^2 \left( \int_{\mathbb{R}^d} \frac{(\xi_1(x) - \xi_1(y))^2}{|x - y|^{d+2s}} dx \right) dy \\ &\quad + 2C_{d,s} \xi_2^2(t) \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \xi_1(x)v(y, t) \frac{(v(x, t) - v(y, t))(\xi_1(x) - \xi_1(y))}{|x - y|^{d+2s}} dx dy. \end{aligned} \quad (\text{A.7})$$

The Cauchy-Schwarz inequality allows to bound the third integral on the r.h.s. of (A.7) by the first two integrals. As concerns the first one, we have:

$$C_{d,s} \xi_2^2(t) \int_{\mathbb{R}^d} \xi_1^2(x) \left( \int_{\mathbb{R}^d} \frac{(v(x,t) - v(y,t))^2}{|x-y|^{d+2s}} dy \right) dx \leq \chi_{\text{supp } \xi_2}(t) \|\xi_2\|_\infty^2 \|\xi_1\|_\infty^2 \|v(t)\|_{\dot{H}^s(\mathbb{R}^d)}^2. \quad (\text{A.8})$$

In order to bound the second integral, it is important to recall that the function  $l_s(\xi_1)(y)$  is regular and decays at least like  $|y|^{-d-2s}$  as  $|y| \rightarrow \infty$  (for the definition and properties of  $l_s$  see Lemmas A.2 and A.3). Hence, thanks to the assumptions on  $\rho$  and  $\gamma$ , we infer that

$$\xi_2^2(t) \int_{\mathbb{R}^d} |v(y,t)|^2 \left( \int_{\mathbb{R}^d} \frac{(\xi_1(x) - \xi_1(y))^2}{|x-y|^{d+2s}} dx \right) dy \leq c' \chi_{\text{supp } \xi_2}(t) \|\xi_2\|_\infty^2 \int_{\mathbb{R}^d} |v(y,t)|^2 \rho(y) dy \quad (\text{A.9})$$

for a suitable positive constant  $c'$ . Integrating in time (A.7), using (A.8), (A.9), (A.1), (A.2) and recalling the identity  $\|(-\Delta)^{\frac{s}{2}}(v_c)(t)\|_{L^2(\mathbb{R}^d)}^2 = \|v_c(t)\|_{\dot{H}^s(\mathbb{R}^d)}^2$ , we then get

$$\|(-\Delta)^{\frac{s}{2}}(v_c)\|_{L^2(\mathbb{R}^{d+1})}^2 \leq C'. \quad (\text{A.10})$$

By exploiting (A.5), (A.6) and (A.10) one deduces (A.4) e.g. using Fourier transform methods.  $\square$

**Lemma A.5.** *Let  $d > 2s$  and  $\phi : \mathbb{R}^d \rightarrow \mathbb{R}$  be a continuous function which belongs to  $L^1(\mathbb{R}^d)$  and decays at least like  $|x|^{-d}$  as  $|x| \rightarrow \infty$ . Then, the convolution  $I_{2s} * \phi$  (namely, the Riesz potential of  $\phi$ ) is also a continuous function, decaying at least like  $|x|^{-d+2s}$  as  $|x| \rightarrow \infty$ .*

*Proof.* The idea of the proof is to split the convolution  $(I_{2s} * \phi)(x)$  in the three regions  $B_{2|x|}^c(0)$ ,  $B_{|x|/2}(x)$ ,  $B_{2|x|}(0) \setminus B_{|x|/2}(x)$  and use there the decay and integrability properties of  $\phi$  and  $I_{2s}$ . We omit the details.  $\square$

**Lemma A.6.** *Let  $d > 2s$  and assume that  $\rho$  satisfies (1.2) for some  $\gamma \in (0, 2s)$ . Let  $v \in L_\rho^1(\mathbb{R}^d) \cap L^\infty(\mathbb{R}^d)$  and  $U_\rho^v$  be the Riesz potential of  $\rho v$ . Then  $U_\rho^v$  belongs to  $C(\mathbb{R}^d) \cap L^p(\mathbb{R}^d)$  for all  $p$  such that*

$$p \in \left( \frac{d}{d-2s}, \infty \right]. \quad (\text{A.11})$$

*Proof.* In order to prove that  $U_\rho^v$  belongs to  $C(\mathbb{R}^d) \cap L^p(\mathbb{R}^d)$  for all  $p$  satisfying (A.11), we proceed as follows:

$$U_\rho^v(x) = \underbrace{\int_{B_1(0)} \rho(y) v(y) I_{2s}(x-y) dy}_{U_{\rho,1}^v(x)} + \underbrace{\int_{\mathbb{R}^d} \chi_{B_1^c(0)}(y) \rho(y) v(y) I_{2s}(x-y) dy}_{U_{\rho,2}^v(x)}.$$

Exploiting the fact that  $v \in L^\infty(\mathbb{R}^d)$  and  $\gamma < 2s$  (so that  $|y|^{-d+2s} \rho(y)$  is locally integrable), it is easily seen that  $U_{\rho,1}^v(x)$  is a continuous function which decays at least like  $|x|^{-d+2s}$  as  $|x| \rightarrow \infty$ . In particular, it belongs to  $L^p(\mathbb{R}^d)$  for all  $p$  satisfying (A.11). As concerns  $U_{\rho,2}^v(x)$ , notice that since  $v \in L_\rho^1(\mathbb{R}^d) \cap L^\infty(\mathbb{R}^d)$  we have that the function  $\chi_{B_1^c(0)} \rho v$  belongs to  $L^1(\mathbb{R}^d) \cap L^\infty(\mathbb{R}^d)$ . Hence  $U_{\rho,2}^v(x)$  is continuous too. To prove that it belongs to  $L^p(\mathbb{R}^d)$  for all  $p$  satisfying (A.11), we write:

$$U_{\rho,2}^v = (\chi_{B_1(0)} I_{2s}) * (\chi_{B_1^c(0)} \rho v) + (\chi_{B_1^c(0)} I_{2s}) * (\chi_{B_1^c(0)} \rho v). \quad (\text{A.12})$$

Since  $\chi_{B_1(0)} I_{2s} \in L^1(\mathbb{R}^d)$  and  $\chi_{B_1^c(0)} \rho v \in L^1(\mathbb{R}^d) \cap L^\infty(\mathbb{R}^d)$ , the first convolution in (A.12) belongs to  $L^1(\mathbb{R}^d) \cap L^\infty(\mathbb{R}^d)$ . Using the fact that  $\chi_{B_1^c(0)} I_{2s} \in L^p(\mathbb{R}^d)$  for all  $p$  as in (A.11) and  $\chi_{B_1^c(0)} \rho v \in L^1(\mathbb{R}^d)$ , we infer that the second convolution in (A.12) belongs to  $L^p(\mathbb{R}^d)$  for all such  $p$ . The latter property is then inherited by  $U_{\rho,2}^v$ .  $\square$



## APPENDIX B

This section is devoted to give a sketch of the proofs of Theorem 3.7 and of the forthcoming Proposition B.1.

*Sketch of proof of Theorem 3.7.* We start from the validity of the fractional “integration by parts” formula

$$C_{d,s} \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \frac{(\phi(x) - \phi(y))(\psi(x) - \psi(y))}{|x - y|^{d+2s}} dx dy = \int_{\mathbb{R}^d} \phi(x) (-\Delta)^s (\psi)(x) dx, \quad (\text{B.1})$$

valid for all  $\phi, \psi \in \mathcal{D}(\mathbb{R}^d)$ , and then to extend it to all functions of  $X_{s,\rho}$ . In order to do it, the first step consists in showing that  $C^\infty(\mathbb{R}^d) \cap X_{s,\rho}$  is dense in  $X_{s,\rho}$ . This can be done by mollification arguments, which however are slightly more complicated than the standard ones, since we work with the weighted spaces  $L_\rho^2(\mathbb{R}^d)$  and  $L_{\rho^{-1}}^2(\mathbb{R}^d)$  instead of  $L^2(\mathbb{R}^d)$ . Hence, given  $v, w \in C^\infty(\mathbb{R}^d) \cap X_{s,\rho}$ , one plugs the cut-off functions  $\phi := \xi_R v$  and  $\psi := \xi_R w$  into (B.1) and lets  $R \rightarrow \infty$ . The problem is that on the r.h.s. there appear terms involving  $\|\xi_R w\|_{\dot{H}^s}$ , and a priori we do not know whether  $C^\infty(\mathbb{R}^d) \cap X_{s,\rho}$  is continuously embedded in  $\dot{H}^s(\mathbb{R}^d)$ . But this turns out to be true: the inequality

$$C_{d,s} \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \frac{(w(x) - w(y))^2}{|x - y|^{d+2s}} dx dy \leq \int_{\mathbb{R}^d} w(x) (-\Delta)^s (w)(x) dx \quad \forall w \in C^\infty(\mathbb{R}^d) \cap X_{s,\rho} \quad (\text{B.2})$$

can be proved just by repeating the above scheme with  $\phi = \psi = \xi_R w$ . In fact, on the r.h.s. of (B.1) we still have terms involving  $\|\xi_R w\|_{\dot{H}^s}$ , but the latter are small and can be absorbed into the l.h.s.; passing to the limit as  $R \rightarrow \infty$  yields (B.2). Therefore, we can now let  $R \rightarrow \infty$  safely in (B.1) (with  $\phi = \xi_R v$  and  $\psi = \xi_R w$ ) and obtain that

$$C_{d,s} \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \frac{(v(x) - v(y))(w(x) - w(y))}{|x - y|^{d+2s}} dx dy = \int_{\mathbb{R}^d} v(x) (-\Delta)^s (w)(x) dx \quad (\text{B.3})$$

for all  $v, w \in C^\infty(\mathbb{R}^d) \cap X_{s,\rho}$ , which in particular shows that (B.2) is actually an equality. Notice that in all these approximation procedures using cut-off functions, to prove that “remainder” terms go to zero we deeply exploit the results provided by Lemmas A.1, A.2 and A.3. It is in fact here that the condition  $\gamma < 2s$  plays a fundamental role: in particular, it ensures that both  $\|\rho^{-1}(-\Delta)^s(\xi_R)\|_\infty$  and  $\|\rho^{-1}l_s(\xi_R)\|_\infty$  vanish as  $R \rightarrow \infty$ . As already mentioned, we refer the reader to the note [33] for the details. However, for similar computations involving  $(-\Delta)^s(\xi_R)$  and  $l_s(\xi_R)$ , see also the proofs of Proposition 4.1, Lemma 4.3 and Lemma 5.5.

By the claimed density of  $C^\infty(\mathbb{R}^d) \cap X_{s,\rho}$ , we are allowed to extend (B.3) to the whole of  $X_{s,\rho}$ . Clearly, the r.h.s. of (B.3) can be rewritten as

$$\int_{\mathbb{R}^d} v(x) A(w)(x) \rho(x) dx,$$

and letting  $v = w$  we obtain that the operator  $A$  is positive. The fact that it is densely defined is trivial since, for instance,  $\mathcal{D}(\mathbb{R}^d) \subset X_{s,\rho}$ . Because in (B.3) one can interchange the role of  $v$  and  $w$ , we also have that  $A$  is symmetric. In order to prove that it is self-adjoint we need to show that  $D(A^*) \subset D(A)$ , namely that any function of  $D(A^*)$  also belongs to  $X_{s,\rho}$ . It is indeed straightforward to check this fact, and we leave the details to the reader.

We finally deal with the quadratic form  $Q$  associated to  $A$ . Thanks to (B.3), we have that

$$Q(v, v) = C_{d,s} \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \frac{(v(x) - v(y))^2}{|x - y|^{d+2s}} dx dy \quad \forall v \in D(A). \quad (\text{B.4})$$

As it is well-known (see e.g. [14]), the domain  $D(Q)$  of  $Q$  is just the closure of  $D(A)$  w.r.t. the norm

$$\|v\|_Q^2 := \|v\|_{2,\rho^{-1}}^2 + Q(v, v) = \|v\|_{2,\rho^{-1}}^2 + \|v\|_{\dot{H}^s}^2.$$

It is then easy to see that such a closure is nothing but  $L_\rho^2(\mathbb{R}^d) \cap \dot{H}^s(\mathbb{R}^d)$  and the quadratic form on  $D(Q) = L_\rho^2(\mathbb{R}^d) \cap \dot{H}^s(\mathbb{R}^d)$  is still represented by (B.4).

By classical results (we refer again to [14]), proving that  $A$  generates a Markov semigroup is equivalent to proving that if  $v$  belongs to  $D(Q)$  then both  $v \vee 0$  and  $v \wedge 1$  belong to  $D(Q)$  and satisfy

$$Q(v \vee 0, v \vee 0) \leq Q(v, v), \quad Q(v \wedge 1, v \wedge 1) \leq Q(v, v).$$

But the latter properties are straightforward consequences of the characterization of  $Q$  given above.

The last assertions follow from the general theory of symmetric Markov semigroups (cf. [14, Section 1.4]) and from their known analyticity properties (cf. [14, Theorem 1.4.2]). See also the discussion in the proof of Lemma 5.3.  $\square$

The next proposition extends the symmetry property of the operator  $A = \rho^{-1}(-\Delta)^s$  to functions which belong to other suitable  $L^p_\rho$  spaces. This is essential in proving our uniqueness Theorem 3.4 for certain values of  $\gamma$  in low dimensions  $d \leq 3$ , more precisely whenever  $(d - \gamma)/(d - 2s) > 2$ .

**Proposition B.1.** *Let  $d > 2s$  and assume that  $\rho$  satisfies (1.2) for some  $\gamma \in [0, 2s) \cap [0, d - 2s]$  and  $\gamma_0 \in [0, \gamma]$ . Let  $p \in [2, 2(d - \gamma)/(d - 2s))$  and  $p' = p/(p - 1)$  be its conjugate exponent. Suppose that  $v, w \in L^p_\rho(\mathbb{R}^d)$  are such that  $A(v), A(w) \in L^{p'}_\rho(\mathbb{R}^d)$ . Then  $v, w \in \dot{H}^s(\mathbb{R}^d)$  and the following formula holds true:*

$$\begin{aligned} \int_{\mathbb{R}^d} v(x)(-\Delta)^s(w)(x) \, dx &= \int_{\mathbb{R}^d} (-\Delta)^s(v)(x) w(x) \, dx \\ &= C_{d,s} \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \frac{(v(x) - v(y))(w(x) - w(y))}{|x - y|^{d+2s}} \, dx dy. \end{aligned}$$

*Sketch of proof.* The method of proof proceeds along the lines of the one of Theorem 3.7. The main difference here lies in the fact that, when using the approximation procedure by cut-off functions mentioned above, if  $p$  is *strictly larger* than 2 in order to prove that “remainder” terms go to zero one cannot exploit the fact that  $\rho^{-1}(-\Delta)^s(\xi_R)$  and  $\rho^{-1}l_s(\xi_R)$  vanish in  $L^\infty(\mathbb{R}^d)$  as  $R \rightarrow \infty$ . In fact, such remainder terms are of the form

$$\int_{\mathbb{R}^d} v^2(x)(-\Delta)^s(\xi_R)(x) \, dx \quad \text{or} \quad \int_{\mathbb{R}^d} v^2(x)l_s(\xi_R)(x) \, dx. \quad (\text{B.5})$$

Thanks to Lemmas A.1, A.2 and A.3 it is direct to see that  $\|\rho^{-1}(-\Delta)^s(\xi_R)\|_{q, -\gamma}$  and  $\|\rho^{-1}l_s(\xi_R)\|_{q, -\gamma}$  vanish as  $R \rightarrow \infty$  provided  $q > (d - \gamma)/(2s - \gamma)$ , whence the condition  $p \in [2, 2(d - \gamma)/(d - 2s))$  to ensure that also the integrals in (B.5) go to zero as  $R \rightarrow \infty$ .  $\square$

#### ACKNOWLEDGEMENTS

G. G. and M. M. have been partially supported by the MIUR-PRIN 2012 grant “Equazioni alle derivate parziali di tipo ellittico e parabolico: aspetti geometrici, disuguaglianze collegate, e applicazioni”. F. P. has been supported by MIUR-PRIN 2012 grant “Critical Point Theory and Perturbative Methods for Nonlinear Differential Equations”. All authors thank the Gruppo Nazionale per l’Analisi Matematica, la Probabilità e le loro Applicazioni (GNAMPA) of the Istituto Nazionale di Alta Matematica (INdAM).

#### REFERENCES

- [1] D. R. Adams, L. I. Hedberg, “Function Spaces and Potential Theory”, Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], 314. Springer-Verlag, Berlin, 1996.
- [2] L. Ambrosio, N. Fusco, D. Pallara, “Functions of Bounded Variation and Free Discontinuity Problems”, Oxford Mathematical Monographs. The Clarendon Press, Oxford University Press, New York, 2000.
- [3] D. G. Aronson, L. Caffarelli, *The initial trace of a solution of the porous medium equation*, Trans. Amer. Math. Soc. 280 (1983), 351–366.
- [4] B. Barrios, I. Peral, F. Soria, E. Valdinoci, *A Widder’s type theorem for the heat equation with nonlocal diffusion*, Arch. Ration. Mech. Anal. 213 (2014), 629–650.
- [5] P. Biler, C. Imbert, G. Karch, *Barenblatt profiles for a nonlocal porous medium equation*, C. R. Math. Acad. Sci. Paris 349 (2011), 641–645.
- [6] M. Bonforte, J. L. Vázquez, *Quantitative local and global a priori estimates for fractional nonlinear diffusion equations*, Adv. Math. 250 (2014), 242–284.

- [7] P. Bénilan, R. Gariepy, *Strong solutions in  $L^1$  of degenerate parabolic equations*, J. Differential Equations 119 (1995), 473–502.
- [8] H. Brézis, A. Friedman, *Nonlinear parabolic equations involving measures as initial data*, J. Math. Pures Appl. 62 (1983), 73–97.
- [9] E. Chasseigne, J. L. Vázquez, *Theory of extended solutions for fast-diffusion equations in optimal classes of data. Radiation from singularities*, Arch. Ration. Mech. Anal. 164 (2002), 133–187.
- [10] E. Chasseigne, J. L. Vázquez, *Extended solutions for general fast diffusion equations with optimal measure data*, Adv. Differential Equations 11 (2006), 627–646.
- [11] H. Chen, L. Véron, Y. Wang, *Fractional heat equations involving initial measure data and subcritical absorption*, preprint arXiv:1401.7187.
- [12] B. E. J. Dahlberg, C. E. Kenig, *Nonnegative solutions of the porous medium equation*, Comm. Partial Differential Equations 9 (1984), 409–437.
- [13] P. D’Ancona, R. Luca’, *Stein-Weiss and Caffarelli-Kohn-Nirenberg inequalities with angular integrability*, J. Math. Anal. Appl. 388 (2012), 1061–1079.
- [14] E. B. Davies, “Heat Kernels and Spectral Theory”, Cambridge Tracts in Mathematics, 92. Cambridge University Press, Cambridge, 1989.
- [15] B. Devyver, M. Fraas, Y. Pinchover, *Optimal hardy weight for second-order elliptic operator: an answer to a problem of Agmon*, J. Funct. Anal. 266 (2014), 4422–4489.
- [16] J. Dolbeault, I. Gentil, A. Guillin, F.-Y. Wang,  *$L^q$ -functional inequalities and weighted porous media equations*, Potential Anal. 28 (2008), 35–59.
- [17] J. Dolbeault, B. Nazaret, G. Savaré, *On the Bakry-Emery criterion for linear diffusions and weighted porous media equations*, Commun. Math. Sci. 6 (2008), 477–494.
- [18] A. de Pablo, F. Quirós, A. Rodríguez, J. L. Vázquez, *A fractional porous medium equation*, Adv. Math. 226 (2011), 1378–1409.
- [19] A. de Pablo, F. Quirós, A. Rodríguez, J. L. Vázquez, *A general fractional porous medium equation*, Comm. Pure Appl. Math. 65 (2012), 1242–1284.
- [20] E. Di Nezza, G. Palatucci, E. Valdinoci, *Hitchhiker’s guide to the fractional Sobolev spaces*, Bull. Sci. Math. 136 (2012), 521–573.
- [21] D. Eidus, *The Cauchy problem for the nonlinear filtration equation in an inhomogeneous medium*, J. Differential Equations 84 (1990), 309–318.
- [22] D. Eidus, S. Kamin, *The filtration equation in a class of functions decreasing at infinity*, Proc. Amer. Math. Soc. 120 (1994), 825–830.
- [23] R. L. Frank, E. H. Lieb, R. Seiringer, *Hardy-Lieb-Thirring inequalities for fractional Schrödinger operators*, J. Amer. Math. Soc. 21 (2008), 925–950.
- [24] G. Grillo, M. Muratori, *Sharp short and long time  $L^\infty$  bounds for solutions to porous media equations with Neumann boundary conditions*, J. Differential Equations 254 (2013), 2261–2288.
- [25] G. Grillo, M. Muratori, M. M. Porzio, *Porous media equations with two weights: existence, uniqueness, smoothing and decay properties of energy solutions via Poincaré inequalities*, Discrete Contin. Dyn. Syst. 33 (2013), 3599–3640.
- [26] G. Grillo, M. Muratori, F. Punzo, *On the asymptotic behaviour of solutions to the fractional porous medium equation with variable density*, preprint arXiv:1403.5293.
- [27] S. Kamin, G. Reyes, J. L. Vázquez, *Long time behavior for the inhomogeneous PME in a medium with rapidly decaying density*, Discrete Contin. Dyn. Syst. 26 (2010), 521–549.
- [28] S. Kamin, P. Rosenau, *Propagation of thermal waves in an inhomogeneous medium*, Comm. Pure Appl. Math. 34 (1981), 831–852.
- [29] S. Kamin, P. Rosenau, *Nonlinear diffusion in a finite mass medium*, Comm. Pure Appl. Math. 35 (1982), 113–127.
- [30] N. S. Landkof, “Foundations of Modern Potential Theory”, Die Grundlehren der mathematischen Wissenschaften, Band 180. Springer-Verlag, New York-Heidelberg, 1972.
- [31] M. Marcus, L. Véron, *Capacitary estimates of solutions of semilinear parabolic equations*, Calc. Var. Partial Differential Equations 48 (2013), 131–183.
- [32] G. Mingione, *Nonlinear measure data problems*, Milan J. Math. 79 (2011), 429–496.
- [33] M. Muratori, *On the self-adjointness of the fractional Laplacian in some weighted  $L^2$  spaces*, in preparation.
- [34] M. M. H. Pang,  *$L^1$  properties of two classes of singular second order elliptic operators*, J. London Math. Soc. 38 (1988), 525–543.
- [35] A. Pazy, “Semigroups of Linear Operators and Applications to Partial Differential Equations”, Applied Mathematical Sciences, 44. Springer-Verlag, New York, 1983.
- [36] M. Pierre, *Uniqueness of the solutions of  $u_t - \Delta\varphi(u) = 0$  with initial datum a measure*, Nonlinear Anal. 6 (1982), 175–187.
- [37] M. M. Porzio, F. Smarrazzo, A. Tesei, *Radon measure-valued solutions for a class of quasilinear parabolic equations*, Arch. Ration. Mech. Anal. 210 (2013), 713–772.
- [38] F. Punzo, *On the Cauchy problem for nonlinear parabolic equations with variable density*, J. Evol. Equ. 9 (2009), 429–447.

- [39] F. Punzo, G. Terrone, *Well-posedness for the Cauchy problem for a fractional porous medium equation with variable density in one space dimension*, Differential Integral Equations 27 (2014), 461–482.
- [40] F. Punzo, G. Terrone, *On the Cauchy problem for a general fractional porous medium equation with variable density*, Nonlinear Anal. 98 (2014), 27–47.
- [41] G. Reyes, J. L. Vázquez, *The Cauchy problem for the inhomogeneous porous medium equation*, Netw. Heterog. Media 1 (2006), 337–351.
- [42] G. Reyes, J. L. Vázquez, *The inhomogeneous PME in several space dimensions. Existence and uniqueness of finite energy solutions*, Commun. Pure Appl. Anal. 7 (2008), 1275–1294.
- [43] G. Reyes, J. L. Vázquez, *Long time behavior for the inhomogeneous PME in a medium with slowly decaying density*, Commun. Pure Appl. Anal. 8 (2009), 493–508.
- [44] J. L. Vázquez, *Barenblatt solutions and asymptotic behaviour for a nonlinear fractional heat equation of porous medium type*, J. Eur. Math. Soc. 16 (2014), 769–803.
- [45] J. L. Vázquez, “The Porous Medium Equation. Mathematical Theory”, Oxford Mathematical Monographs. The Clarendon Press, Oxford University Press, Oxford, 2007.

GABRIELE GRILLO, MATTEO MURATORI: DIPARTIMENTO DI MATEMATICA “F. BRIOSCHI”, POLITECNICO DI MILANO, PIAZZA LEONARDO DA VINCI 32, 20133 MILANO, ITALY  
*E-mail address:* gabriele.grillo@polimi.it  
*E-mail address:* matteo.muratori@polimi.it

FABIO PUNZO: DIPARTIMENTO DI MATEMATICA “F. ENRIQUES”, UNIVERSITÀ DEGLI STUDI DI MILANO, VIA CESARE SALDINI 50, 20133 MILANO, ITALY  
*E-mail address:* fabio.punzo@unimi.it