



# Asymptotic behavior of generalized capacities with applications to eigenvalue perturbations: The higher dimensional case



Laura Abatangelo<sup>a</sup>, Corentin Léna<sup>b,c</sup>, Paolo Musolino<sup>d,\*</sup>

<sup>a</sup> *Dipartimento di Matematica, Politecnico di Milano, P.zza Leonardo da Vinci 32, 20133 Milano, Italy*

<sup>b</sup> *Dipartimento di Tecnica e Gestione dei Sistemi Industriali (DTG), Università degli Studi di Padova, Stradella S. Nicola 3, 36100 Vicenza, Italy*

<sup>c</sup> *Dipartimento di Matematica "Tullio Levi-Civita", Università degli Studi di Padova, via Trieste 63, 35121 Padova, Italy*

<sup>d</sup> *Dipartimento di Scienze Molecolari e Nanosistemi, Università Ca' Foscari Venezia, via Torino 155, 30172 Venezia Mestre, Italy*

## ARTICLE INFO

### Article history:

Received 11 May 2023

Accepted 12 September 2023

Communicated by Enrico Valdinoci

### MSC:

35P15

31C15

31B10

35B25

35C20

### Keywords:

Dirichlet–Laplacian

Capacity

Multiple eigenvalues

Asymptotic expansion

Perforated domains

## ABSTRACT

We provide a full series expansion of a generalization of the so-called  $u$ -capacity related to the Dirichlet–Laplacian in dimension three and higher, extending the results of Abatangelo et al. (2021); Abatangelo, Léna and Musolino (2022) dealing with the planar case. We apply the result in order to study the asymptotic behavior of perturbed eigenvalues when Dirichlet conditions are imposed on a small regular subset of the domain of the eigenvalue problem.

© 2023 The Author(s). Published by Elsevier Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

## 1. Introduction

This paper is devoted to the analysis of the asymptotic behavior of a generalization of the condenser capacity of a bounded domain of  $\mathbb{R}^d$  ( $d \geq 3$ ) with a small hole of size  $\varepsilon > 0$ , as the parameter  $\varepsilon$  approaches the degenerate value 0. Such a generalization, as we shall see, has proved to be extremely useful for the analysis of the behavior of simple and multiple eigenvalues for the Dirichlet–Laplacian in a bounded domain with a small hole. A careful investigation of the behavior of such a generalized capacity has been carried out

\* Corresponding author.

*E-mail addresses:* [laura.abatangelo@polimi.it](mailto:laura.abatangelo@polimi.it) (L. Abatangelo), [corentin.lena@unipd.it](mailto:corentin.lena@unipd.it) (C. Léna), [paolo.musolino@unive.it](mailto:paolo.musolino@unive.it) (P. Musolino).

in [1,3] in the planar case, with the corresponding applications to the study of the eigenvalues in perforated domains. Here, instead, we wish to consider the case of dimension  $d$  greater than or equal to 3. As we shall see, the cases of dimension  $d = 2$  and of  $d \geq 3$  need to be treated separately because of the different aspect of the fundamental solution of the Laplace operator.

Before introducing the generalization of the capacity we are going to study, we begin by recalling the notion of *condenser capacity* and a first generalization of it, known as *u-capacity*.

We recall that for a bounded, connected open set  $\Omega$  of  $\mathbb{R}^d$  and a compact subset  $K$  of  $\Omega$ , the (*condenser capacity*) of  $K$  in  $\Omega$  is

$$\text{Cap}_\Omega(K) \equiv \inf \left\{ \int_\Omega |\nabla f|^2 dx : f \in H_0^1(\Omega) \text{ and } f - \eta_K \in H_0^1(\Omega \setminus K) \right\}, \tag{1}$$

where  $\eta_K$  is a fixed smooth function such that  $\text{supp } \eta_K \subseteq \Omega$  and  $\eta_K \equiv 1$  in a neighborhood of  $K$ . As is well known, the infimum in (1) is achieved by a function  $V_K \in H_0^1(\Omega)$  such that  $V_K - \eta_K \in H_0^1(\Omega \setminus K)$  so that

$$\text{Cap}_\Omega(K) = \int_\Omega |\nabla V_K|^2 dx,$$

where  $V_K$  (*capacitary potential*) is the unique solution of the Dirichlet problem

$$\begin{cases} \Delta V_K = 0 & \text{in } \Omega \setminus K, \\ V_K = 0 & \text{on } \partial\Omega, \\ V_K = 1 & \text{on } K. \end{cases} \tag{2}$$

By saying that  $V_K$  solves (2) we mean that  $V_K \in H_0^1(\Omega)$ ,  $V_K - \eta_K \in H_0^1(\Omega \setminus K)$ , and

$$\int_{\Omega \setminus K} \nabla V_K \cdot \nabla \phi dx = 0 \quad \forall \phi \in H_0^1(\Omega \setminus K).$$

Moreover, if  $\Omega$  and  $K$  are sufficiently regular, one can reformulate the boundary conditions of problem (2) in the trace sense. Let us also point out that  $\text{Cap}_\Omega(K)$  and  $V_K$  do not depend on the choice of  $\eta_K$ , but only on the set  $K$ .

The following set of properties for the domains will play a key role in our analysis, so we summarize it in a definition.

**Definition 1.1.** We say that  $\Omega \subset \mathbb{R}^d$  is an *admissible domain* if

1.  $\Omega$  is open, bounded and connected;
2.  $\Omega$  is in the Schauder class  $C^{1,\alpha}$  for some  $\alpha \in ]0, 1[$ ;
3.  $0 \in \Omega$  and  $\mathbb{R}^d \setminus \overline{\Omega}$  is connected.

In the case where  $d \geq 3$ , we can use the notion of *Newtonian capacity*. We restrict ourselves to the case where  $K = \overline{\omega}$ , with  $\omega$  an admissible domain. Then, there exists a unique function

$$W_K : \mathbb{R}^d \setminus \omega \rightarrow \mathbb{R}$$

that is continuous, real-analytic in the open set  $\mathbb{R}^d \setminus K$ , and satisfies

$$\begin{cases} \Delta W_K = 0 & \text{in } \mathbb{R}^d \setminus K, \\ W_K = 1 & \text{on } \partial K, \\ W_K(x) \rightarrow 0 & \text{for } x \rightarrow \infty. \end{cases} \tag{3}$$

The last condition in (3) is equivalent to the requirement that  $W_K$  be harmonic at infinity. We then define the Newtonian capacity as

$$\text{Cap}_{\mathbb{R}^d}(K) \equiv \int_{\mathbb{R}^d \setminus K} |\nabla W_K|^2 dx, \tag{4}$$

the integral being finite.

In order to study the asymptotic behavior as  $\varepsilon \rightarrow 0$  of eigenvalues of the Dirichlet–Laplacian in perforated domains of the type

$$\Omega_\varepsilon = \Omega \setminus \varepsilon\bar{\omega},$$

where  $\Omega, \omega \subset \mathbb{R}^d$  are admissible domains in the sense of Definition 1.1, a generalization of the notion of capacity –the so-called *u-capacity*–, has been introduced (see Abatangelo, Felli, Hillairet, and Léna [2]).

**Definition 1.2.** Given a function  $u \in H_0^1(\Omega)$ , the *u-capacity* of a compact set  $K \subseteq \Omega$  is

$$\text{Cap}_\Omega(K, u) \equiv \inf \left\{ \int_\Omega |\nabla f|^2 dx : f \in H_0^1(\Omega) \text{ and } f - u \in H_0^1(\Omega \setminus K) \right\}. \tag{5}$$

The infimum in (5) is achieved by a unique function  $V_{K,u} \in H_0^1(\Omega)$ , so that

$$\text{Cap}_\Omega(K, u) = \int_\Omega |\nabla V_{K,u}|^2 dx.$$

We call  $V_{K,u}$  the *potential* associated with  $u$  and  $K$ .

Furthermore,  $V_{K,u}$  is the unique weak solution of the Dirichlet problem

$$\begin{cases} \Delta V_{K,u} = 0 & \text{in } \Omega \setminus K, \\ V_{K,u} = 0 & \text{on } \partial\Omega, \\ V_{K,u} = u & \text{on } K, \end{cases}$$

where, by *weak solution*, we mean that  $V_{K,u} \in H_0^1(\Omega)$ ,  $u - V_{K,u} \in H_0^1(\Omega \setminus K)$  and  $\int_\Omega \nabla V_{K,u} \cdot \nabla \varphi dx = 0$  for all  $\varphi \in H_0^1(\Omega \setminus K)$ .

We extend Definition 1.2 to  $H^1(\Omega)$  functions, by setting, for any  $u \in H^1(\Omega)$ ,

$$\text{Cap}_\Omega(K, u) \equiv \text{Cap}_\Omega(K, \eta_K u),$$

where  $\eta_K$  is a fixed smooth function such that  $\text{supp } \eta_K \subseteq \Omega$  and  $\eta_K \equiv 1$  in a neighborhood of  $K$ . Here again,  $\text{Cap}_\Omega(K, u)$  and the associated potential  $V_{K,u}$  do not depend on the choice of  $\eta_K$ , but only on  $K$  and  $u$ .

With this tool in hand, one can obtain an asymptotic formula for the behavior of  $N$ th eigenvalue  $\lambda_N(\Omega_\varepsilon)$  of the Dirichlet–Laplacian in  $\Omega_\varepsilon$  as  $\varepsilon \rightarrow 0$ , under the assumption that the  $N$ th eigenvalue  $\lambda_N(\Omega)$  of the Dirichlet–Laplacian in the unperturbed set  $\Omega$  is simple.

In order to be more precise, we recall that if  $\tilde{\Omega}$  is a bounded open set in  $\mathbb{R}^d$  the eigenvalue problem

$$\begin{cases} -\Delta u = \lambda u & \text{in } \tilde{\Omega}, \\ u = 0 & \text{on } \partial\tilde{\Omega} \end{cases}$$

admits a sequence of real eigenvalues tending to infinity

$$0 < \lambda_1(\tilde{\Omega}) \leq \lambda_2(\tilde{\Omega}) \leq \dots \leq \lambda_N(\tilde{\Omega}) \leq \dots \rightarrow +\infty$$

where every eigenvalue is repeated as many times as its multiplicity and, in particular, the first one is simple if  $\tilde{\Omega}$  is connected. The dependence of the spectrum of the Laplace operator upon domain perturbations has long been investigated, with particular attention to the case of sets with small perforations. Here we mention, for example, Abatangelo, Felli, Hillairet, and Léna [2], Ammari, Kang, and Lee [4], Besson [5], Chavel and Feldman [7], Colbois and Courtois [8], Courtois [10], Felli, Noris, and Ognibene [17], Lamberti and Perin [23], Lanza de Cristoforis [28], Maz’ya, Movchan, and Nieves [30], Maz’ya, Nazarov, and Plamenevskiĭ [32], Ozawa [33], Rauch and Taylor [36], Samarskiĭ [37]. In particular, we recall that, by Courtois [10, Proof

of Theorem 1.2] and by Abatangelo, Felli, Hillairet, and Léna [2, Theorem 1.4], an asymptotic formula for the eigenvalues can be obtained in terms of  $\text{Cap}_\Omega(\varepsilon\bar{\omega}, u_N)$ , where  $u_N$  is a  $L^2(\Omega)$ -normalized eigenfunction corresponding to  $\lambda_N(\Omega)$ : if  $\Omega$  and  $\omega$  are admissible domains and  $\lambda_N(\Omega)$  is simple, this reads as

$$\lambda_N(\Omega \setminus (\varepsilon\bar{\omega})) = \lambda_N(\Omega) + \text{Cap}_\Omega(\varepsilon\bar{\omega}, u_N) + o(\text{Cap}_\Omega(\varepsilon\bar{\omega}, u_N)) \quad \text{as } \varepsilon \rightarrow 0^+.$$

Unfortunately, the above mentioned result a priori only holds for simple eigenvalues. In order to study multiple eigenvalues, one needs a further generalization of the capacity. The eigenspace being of dimension greater than one, one needs to take into account the interaction between different eigenfunctions corresponding to the same eigenvalue.

**Definition 1.3.** Given  $u^a, u^b \in H_0^1(\Omega)$ , the  $(u^a, u^b)$ -capacity of a compact set  $K \subseteq \Omega$  is

$$\text{Cap}_\Omega(K, u^a, u^b) \equiv \int_\Omega \nabla V_{K, u^a} \cdot \nabla V_{K, u^b} \, dx.$$

Here above the symbol  $\cdot$  denotes the scalar product in  $\mathbb{R}^d$ .

Again, as we have done for the  $u$ -capacity, we extend Definition 1.3 to  $H^1(\Omega)$  functions, by setting, for any pair of functions  $u^a, u^b \in H^1(\Omega)$ ,

$$\text{Cap}_\Omega(K, u^a, u^b) \equiv \text{Cap}_\Omega(K, \eta_K u^a, \eta_K u^b),$$

where  $\eta_K$  is a fixed smooth function such that  $\text{supp } \eta_K \subseteq \Omega$  and  $\eta_K \equiv 1$  in a neighborhood of  $K$ .

With the new object  $\text{Cap}_\Omega(K, u^a, u^b)$  we can clearly recover the classical condenser capacity as well as the  $u$ -capacity.

**Remark 1.4.** If  $u \in H^1(\Omega)$  and  $K$  is a compact subset of  $\Omega$ , then

$$\text{Cap}_\Omega(K, u, u) = \text{Cap}_\Omega(K, u).$$

Also,

$$\text{Cap}_\Omega(K, 1, 1) = \text{Cap}_\Omega(K).$$

In [3] it has been shown that by exploiting the definition of  $(u^a, u^b)$ -capacity, we can obtain the asymptotic behavior of multiple eigenvalues. More precisely, if one has an asymptotic expansion of  $\text{Cap}_\Omega(\varepsilon\bar{\omega}, u^a, u^b)$  where  $u^a$  and  $u^b$  are eigenfunctions corresponding to the same multiple eigenvalue in the unperturbed set  $\Omega$ , then one can deduce the asymptotic behavior of the corresponding eigenvalues in the perforated set  $\Omega \setminus (\varepsilon\bar{\omega})$ . Therefore, in [3], we have obtained a detailed and fully constructive representation of  $\text{Cap}_\Omega(\varepsilon\bar{\omega}, u^a, u^b)$  as a convergent series in case of dimension  $d = 2$ . Since our method is based on potential theory, the higher dimensional case differs from the two-dimensional one (mainly due to the different aspect of the fundamental solution for the Laplace equation). The goal of the present paper is to obtain a representation of  $\text{Cap}_\Omega(\varepsilon\bar{\omega}, u^a, u^b)$  for  $\varepsilon$  close to 0 in terms of a convergent series and to apply it to study the asymptotic behavior of Dirichlet eigenvalues in perforated domains, in the case  $d \geq 3$ . Such results extend the work done in [3] for the case of dimension  $d = 2$ . Moreover, taking  $u^a = u^b = 1$ , one can immediately deduce the expansion for the condenser capacity  $\text{Cap}_\Omega(\varepsilon\bar{\omega})$  as  $\varepsilon \rightarrow 0$ . We observe that several authors have considered the asymptotic behavior of condenser capacities in several geometrical situations (see Dubinin [16], Lanza de Cristoforis [25–27], Maz’ya, Nazarov, and Plamenevskij [31, §8.1], Soĭbel’man [40]). As an example, we mention that the asymptotic expansion of the capacity as the hole shrinks to a point can be deduced from

the analysis of energy integrals in perforated domains by Maz'ya, Nazarov, and Plamenevskij [31, §8.1]. For example, in dimension two, they prove that there exists  $\delta > 0$  such that

$$\text{Cap}_\Omega(\varepsilon\bar{\omega}) = -\frac{2\pi}{\log \varepsilon + 2\pi(H_{(0,0)} + N)} + o(\varepsilon^\delta), \tag{6}$$

for  $\varepsilon$  small and positive, where  $e^{2\pi N}$  is the *logarithmic capacity* (or *outer conformal radius*) of  $\omega$  and  $H_{(0,0)}$  is the value at  $x = 0$  of the unique harmonic function  $h$  in  $\Omega$  such that  $h(x) = -\log|x|/(2\pi)$  for all  $x \in \partial\Omega$ . For the definition of logarithmic capacity we refer, *e.g.*, to Landkof [24, p. 168], Pólya and Szegő [34, p. 2], and Pommerenke [35, p. 332]. In other words,  $N$  in Eq. (6) is defined as  $1/(2\pi)$  multiplied by the logarithm of the logarithmic capacity of  $\omega$ . Moreover, expansions for the capacity for the case of several small inclusions can be deduced from the corresponding expansion of the capacity potential obtained in Maz'ya, Movchan, and Nieves [29, §3.2.2].

A different method to analyze the asymptotic behavior of functionals in perforated domains is the so-called *Functional Analytic Approach* proposed by Lanza de Cristoforis in [25,26]. The goal of this method is to represent functionals in singularly perturbed domains by means of real analytic maps and possibly singular but explicitly known functions of the singular perturbation parameter (see Dalla Riva, Lanza de Cristoforis, and Musolino [11] for an introduction). As far as the condenser capacity in dimension two is concerned, by Lanza de Cristoforis [25,26], we know that there exist  $\varepsilon_1 > 0$  and a real analytic function  $\mathcal{R}$  from  $] - \varepsilon_1, \varepsilon_1[$  to  $]0, +\infty[$  such that

$$\text{Cap}_\Omega(\varepsilon\bar{\omega}) = -\frac{2\pi}{\log \varepsilon + \log \mathcal{R}[\varepsilon]} \quad \forall \varepsilon \in ]0, \varepsilon_1[.$$

Such a result implies that there exists a real analytic map  $\tilde{\mathcal{R}}$  from a neighborhood of  $(0, 0)$  in  $\mathbb{R}^2$  with values in  $\mathbb{R}$  such that

$$\text{Cap}_\Omega(\varepsilon\bar{\omega}) = \tilde{\mathcal{R}}\left[\varepsilon, \frac{1}{\log \varepsilon}\right],$$

for  $\varepsilon$  positive and close to 0. As a consequence, we have that

$$\text{Cap}_\Omega(\varepsilon\bar{\omega}) = \sum_{(k,l) \in \mathbb{N}^2} \gamma_{(k,l)} \varepsilon^k \left(\frac{1}{\log \varepsilon}\right)^l,$$

for  $\varepsilon$  positive and small enough, where the double power series  $\sum_{(k,l) \in \mathbb{N}^2} \gamma_{(k,l)} x_1^k x_2^l$  converges for  $(x_1, x_2)$  in a neighborhood of  $(0, 0)$ . Moreover, in Lanza de Cristoforis [27], the Functional Analytic Approach has been applied to the capacity  $\text{Cap}_\Omega(\varepsilon\bar{\omega})$  in  $\mathbb{R}^d$  with  $d \geq 2$  and has shown the existence of two real analytic maps  $V_1, V_2$  from a neighborhood of 0 to  $\mathbb{R}$  such that

$$\text{Cap}_\Omega(\varepsilon\bar{\omega}) = \frac{-1}{V_1[\varepsilon] + V_2[\varepsilon] \mathcal{Y}_d[\varepsilon]} \tag{7}$$

for  $\varepsilon$  positive and small enough, with  $\mathcal{Y}_d[\varepsilon]$  defined as

$$\mathcal{Y}_d[\varepsilon] \equiv \begin{cases} \frac{1}{2\pi} \log \varepsilon & \text{if } d = 2, \\ \frac{1}{(2-d)s_d} \varepsilon^{2-d} & \text{if } d \geq 3, \end{cases}$$

where  $s_d$  denotes the  $(d - 1)$ -dimensional measure of the unit sphere in  $\mathbb{R}^d$ . In dimension  $d \geq 3$  formula (7) implies the possibility of representing  $\text{Cap}_\Omega(\varepsilon\bar{\omega})$  as a convergent power series in  $\varepsilon$ , even though the explicit computation of the coefficients of the series is not available.

In [1,3], we have improved the above mentioned two-dimensional results in two directions: first we have considered a generalization of the classical capacity (the *u*-capacity), then we have explicitly computed some of the coefficients of the series representing the *u*-capacity. This result has then been applied to the

asymptotic behavior of simple and multiple eigenvalues of the Laplacian in perforated domains in dimension  $d = 2$ . The purpose of the present paper is to consider the behavior of (generalized) capacities in dimension three and higher together with its application to the asymptotic behavior of Dirichlet–Laplacian eigenvalues.

We wish to stress that to obtain an asymptotic expansion of  $\text{Cap}_\Omega(\varepsilon\bar{\omega}, u^a, u^b)$ , we follow the lines of the computations of [1,3]. Although the strategy is similar, some changes need to be taken into account due to the different dimension (here  $d \geq 3$ , whereas  $d = 2$  in [1,3]). Since they deserve some attention and care, we decided to provide these computations in the present paper.

We do believe that explicit results in dimension  $d \geq 3$  will be useful both to people interested in the behavior of capacities and to those interested in the asymptotic analysis of eigenvalues of the Laplacian in perforated domains. To the best of our knowledge, the explicit and constructive series expansion we obtain is new even for the classical capacity.

We assume that  $\Omega$  and  $\omega$  are admissible domains in the sense of Definition 1.1. These regularity assumptions on  $\Omega$  and  $\omega$  are convenient for the Functional Analytic Approach we adopt, although they can be relaxed to Lipschitz regularity as it has been done in Costabel, Dalla Riva, Dauge, and Musolino [9]

**Definition 1.5.** Let  $\Omega$  be an admissible domain. We call  $u$  an *admissible function* if

1.  $u \in H^1(\Omega)$ ;
2.  $u$  is real-analytic in a neighborhood of 0.

Our first main result, Theorem 5.2, deals with the value of  $\text{Cap}_\Omega(\varepsilon\bar{\omega}, u^a, u^b)$  for  $\varepsilon$  close to 0. We reformulate Theorem 5.2 in Theorem 1.6 below. For the proof, we refer to Section 5.

**Theorem 1.6.** Let  $\Omega, \omega$  be admissible domains and let  $u^a, u^b$  be admissible functions. Then there exist  $\varepsilon_c^\#$  positive and small enough and a sequence  $\{c_n^\#\}_{n \in \mathbb{N}}$  of real numbers such that

$$\text{Cap}_\Omega(\varepsilon\bar{\omega}, u^a, u^b) = \varepsilon^{d-2} \sum_{n=0}^\infty c_n^\# \varepsilon^n$$

for all  $\varepsilon \in ]0, \varepsilon_c^\#[$ .

In addition, we are able to give a simple description of the first few terms in the above series expansion. For an admissible function  $u$  which is not identically zero in a neighborhood of 0, we denote by  $\kappa(u) \in \mathbb{N}$  the order of vanishing of  $u$  at 0, so that  $D^\gamma u(0) = 0$  for all  $|\gamma| < \kappa(u)$  and  $D^\beta u(0) \neq 0$  for some  $\beta \in \mathbb{N}^d$  with  $|\beta| = \kappa(u)$  (we use the standard multi-index notation throughout the paper). We define the *principal part* of  $u$  by

$$u_\#(x) \equiv \sum_{\beta \in \mathbb{N}^d, |\beta| = \kappa(u)} \frac{1}{\beta!} D^\beta u(0) x^\beta,$$

so that  $u_\#$  is a homogeneous polynomial of degree  $\kappa(u)$ . Note that this includes the case when  $\kappa(u) = 0$ , i.e.,  $u(0) \neq 0$ . Then  $u_\# = u(0)$ . We finally denote by  $U$  the unique function, continuous in  $\mathbb{R}^d \setminus \omega$ , which solves the exterior boundary value problem

$$\begin{cases} \Delta U = 0 & \text{in } \mathbb{R}^d \setminus \bar{\omega}, \\ U = u_\# & \text{on } \partial\omega, \\ U(x) \rightarrow 0 & \text{for } |x| \rightarrow \infty. \end{cases} \tag{8}$$

We now assume that the admissible functions  $u^a, u^b$  have finite orders of vanishing  $\kappa(u^a) = \bar{k}_a$  and  $\kappa(u^b) = \bar{k}_b$  at 0. We write

$$\mathfrak{C}(\omega, (u^a)_\#, (u^b)_\#) \equiv \int_{\mathbb{R}^d \setminus \bar{\omega}} \nabla U^a \cdot \nabla U^b \, dx + \int_\omega \nabla u^a_\# \cdot \nabla u^b_\# \, dx,$$

where, for  $l = a, b$ , the function  $u_{\#}^l$  is the principal part of  $u^l$  and the function  $U^l$  is the solution of problem (8) with  $u_{\#}$  replaced by  $u_{\#}^l$ . Then,  $c_n^{\#} = 0$  for all  $n < \bar{k}_a + \bar{k}_b$  and

$$c_{\bar{k}_a + \bar{k}_b}^{\#} = \mathfrak{C}(\omega, (u^a)_{\#}, (u^b)_{\#}).$$

When  $u$  is a single admissible function, with finite order of vanishing  $\kappa(u) = \bar{k}$  at 0, we simplify notation further by writing

$$\mathfrak{C}(\omega, u_{\#}) \equiv \mathfrak{C}(\omega, u_{\#}, u_{\#}),$$

and we note that

$$\mathfrak{C}(\omega, u_{\#}) = \int_{\mathbb{R}^d \setminus \bar{\omega}} |\nabla U|^2 dx + \int_{\omega} |\nabla u_{\#}|^2 dx \tag{9}$$

is strictly positive. In particular,  $\text{Cap}_{\Omega}(\varepsilon\bar{\omega}, u)$  is asymptotic to  $\mathfrak{C}(\omega, u_{\#})\varepsilon^{2\bar{k}+d-2}$ .

We then apply the above results to the asymptotic behavior of eigenvalues in perforated domains. Our main results on this problem are the following [Theorems 1.7](#) and [1.8](#) (see [Theorems 7.2](#), [7.3](#), and [7.8](#) for more detailed statements). We begin with [Theorem 1.7](#) which is a reformulation of [Theorems 7.2](#) and [7.3](#) and which deals with the case when the  $N$ th eigenvalue in the unperturbed set  $\Omega$  is simple. For the proof, we refer to [Section 7.1](#).

**Theorem 1.7.** *Let  $\lambda_N(\Omega)$  be a simple eigenvalue of the Dirichlet-Laplacian in an admissible domain  $\Omega$ . Let  $u_N$  be a  $L^2(\Omega)$ -normalized eigenfunction associated to  $\lambda_N(\Omega)$ . Then*

$$\lambda_N(\Omega \setminus (\varepsilon\bar{\omega})) = \lambda_N(\Omega) + (u_N(0))^2 \text{Cap}_{\mathbb{R}^d}(\bar{\omega})\varepsilon^{d-2} + o(\varepsilon^{d-2}) \quad \text{as } \varepsilon \rightarrow 0^+.$$

Moreover, let  $\bar{k}$  be the order of vanishing of  $u_N$  at 0. Then

$$\lambda_N(\Omega \setminus (\varepsilon\bar{\omega})) = \lambda_N(\Omega) + \mathfrak{C}(\omega, (u_N)_{\#})\varepsilon^{2\bar{k}+d-2} + o(\varepsilon^{2\bar{k}+d-2}) \quad \text{as } \varepsilon \rightarrow 0^+.$$

[Theorem 1.8](#) here below is instead concerned with the case when  $\lambda_N(\Omega)$  is an eigenvalue of multiplicity  $m > 1$ . The case of multiple eigenvalues is considered in [Theorem 7.8](#) and its proof can be found in [Section 7.2](#).

**Theorem 1.8.** *Let  $\lambda_N(\Omega)$  be an eigenvalue of multiplicity  $m > 1$  and let  $E(\lambda_N(\Omega))$  be the associated eigenspace. There exists an orthonormal basis of  $E(\lambda_N(\Omega))$ ,*

$$v_1, \dots, v_i, \dots, v_m,$$

such that, for  $1 \leq i \leq m$ ,

$$\lambda_{N-1+i}(\Omega \setminus (\varepsilon\bar{\omega})) = \lambda_N(\Omega) + \text{Cap}_{\Omega}(\varepsilon\bar{\omega}, v_i) + o(\text{Cap}_{\Omega}(\varepsilon\bar{\omega}, v_i)) \quad \text{as } \varepsilon \rightarrow 0^+.$$

Moreover, we have

$$\lambda_{N-1+i}(\Omega \setminus (\varepsilon\bar{\omega})) = \lambda_N(\Omega) + \mu_i \varepsilon^{2\bar{k}_i+d-2} + o(\varepsilon^{2\bar{k}_i+d-2}) \quad \text{as } \varepsilon \rightarrow 0^+,$$

where  $\bar{k}_i$  is the order of vanishing of  $v_i$  at 0 and

$$\mu_j = \mathfrak{C}(\omega, (v_j)_{\#}) > 0.$$

**Remark 1.9.** In the previous theorem, the non-increasing finite sequence of integers  $(\bar{k}_i)$  and the non-decreasing finite sequence of positive numbers  $(\mu_i)$  are obviously independent of the choice of a basis  $(v_i)$  satisfying the properties, by uniqueness of the asymptotic expansion.

The paper is organized as follows. In Section 2 we rewrite the boundary value problem associated to the generalization of the capacity in terms of integral equations, exploiting classical potential theory. In Sections 3 and 4 we obtain series expansions for the solutions of the integral equations and for an auxiliary function. In Section 5, we deduce our main result on the series expansion of the generalized capacity  $\text{Cap}_\Omega(\varepsilon\bar{\omega}, u^a, u^b)$  for  $\varepsilon$  close to 0. In Section 6, we compute the principal term of the asymptotic expansion of  $\text{Cap}_\Omega(\varepsilon\bar{\omega}, u^a, u^b)$  under vanishing assumption for  $u^a$  and  $u^b$ . In Section 7 we outline the proofs of Theorem 1.7 and Theorem 1.8. Moreover, we supplement the present paper with a blow-up analysis for the  $u$ -capacity in Appendix. It shows more clearly that the first term of the series expansion provided in Section 5 can be seen as a suitable capacity in the whole space  $\mathbb{R}^d$ .

## 2. Integral equation formulation of capacity potentials

### 2.1. Preliminaries and classical notions of potential theory

In this paper we consider the dimension

$$d \in \mathbb{N} \setminus \{0, 1, 2\}$$

and we study the asymptotic behavior of  $\text{Cap}_\Omega(\varepsilon\bar{\omega}, u^a, u^b)$  as  $\varepsilon \rightarrow 0$ . To do so, we assume some smoothness on the sets and on the functions  $u^a$  and  $u^b$ . We work in the frame of Schauder classes and thus we assume that both  $\Omega$  and  $\omega$  are admissible domains in the sense of Definition 1.1. We can obviously find a common Schauder class  $C^{1,\alpha}$  to which they belong, up to taking a smaller  $\alpha$ . Since  $\Omega$  is open and contains 0, and since  $\omega$  bounded, it is clear that there exists  $\varepsilon_\#$  such that

$$\varepsilon_\# > 0 \text{ and } \varepsilon\bar{\omega} \subseteq \Omega \text{ for all } \varepsilon \in ]-\varepsilon_\#, \varepsilon_\#[. \tag{10}$$

To define our perforated domain, we set

$$\Omega_\varepsilon \equiv \Omega \setminus (\varepsilon\bar{\omega}) \quad \forall \varepsilon \in ]-\varepsilon_\#, \varepsilon_\#[.$$

Clearly,  $\Omega_\varepsilon$  is an open bounded connected subset of  $\mathbb{R}^d$  of class  $C^{1,\alpha}$  for all  $\varepsilon \in ]-\varepsilon_\#, \varepsilon_\#[ \setminus \{0\}$ . The boundary  $\partial\Omega_\varepsilon$  of  $\Omega_\varepsilon$  is the union of  $\partial\Omega$  and  $\partial(\varepsilon\omega) = \varepsilon\partial\omega$ , for all  $\varepsilon \in ]-\varepsilon_\#, \varepsilon_\#[ \setminus \{0\}$ . For  $\varepsilon = 0$ , we have  $\Omega_0 = \Omega \setminus \{0\}$ . We also need some regularity on the functions  $u^a, u^b$ : we assume that  $u^a, u^b$  are admissible in the sense of Definition 1.5.

Our goal is to provide accurate and explicit expansions for  $\text{Cap}_\Omega(\varepsilon\bar{\omega}, u^a, u^b)$  in terms of  $\varepsilon$ , with particular emphasis on the influence of the geometry and the data of the problem on such formulas.

Our strategy will be the same of [1,3], where we adopted the Functional Analytic Approach of Lanza de Cristoforis [25,26] for the analysis of singularly perturbed boundary value problems (see Dalla Riva, Lanza de Cristoforis, and Musolino [11] for a detailed presentation). This approach permits to deduce a representation of the solution or related functionals as real-analytic maps, and thus as convergent power series.

To analyze  $\text{Cap}_\Omega(\varepsilon\bar{\omega}, u^a, u^b)$ , we modify the techniques of [1,3], where we considered  $\text{Cap}_\Omega(\varepsilon\bar{\omega}, u)$  and  $\text{Cap}_\Omega(\varepsilon\bar{\omega}, u^a, u^b)$  but only in the planar case. Indeed, we emphasize that for considering  $d \geq 3$  some modifications need to be done, as it is customary when using Potential Theory.

By the analyticity of  $u^a$  and  $u^b$  (see Definition 1.5) and analyticity results for the composition operator (see Böhme and Tomi [6, p. 10], Henry [21, p. 29], Valent [41, Thm. 5.2, p. 44]), we deduce that, possibly shrinking  $\varepsilon_\#$ , there exists two real analytic maps  $U_\#^a, U_\#^b$  from  $]-\varepsilon_\#, \varepsilon_\#[$  to  $C^{1,\alpha}(\partial\omega)$  such that

$$u^a(\varepsilon t) = U_\#^a[\varepsilon](t), \quad u^b(\varepsilon t) = U_\#^b[\varepsilon](t), \quad \forall t \in \partial\omega, \forall \varepsilon \in ]-\varepsilon_\#, \varepsilon_\#[$$



(see Deimling [15, §15] for the definition and properties of analytic maps). Then for all  $\varepsilon \in ]-\varepsilon_{\#}, \varepsilon_{\#}[ \setminus \{0\}$ , we denote by  $u_{\varepsilon}^a$  and  $u_{\varepsilon}^b$  the unique solutions in  $C^{1,\alpha}(\overline{\Omega_{\varepsilon}})$  of the problems

$$\begin{cases} \Delta u_{\varepsilon}^a = 0 & \text{in } \Omega_{\varepsilon}, \\ u_{\varepsilon}^a(x) = 0 & \text{for all } x \in \partial\Omega, \\ u_{\varepsilon}^a(x) = U_{\#}^a[\varepsilon](x/\varepsilon) & \text{for all } x \in \varepsilon\partial\omega \end{cases} \tag{11}$$

and

$$\begin{cases} \Delta u_{\varepsilon}^b = 0 & \text{in } \Omega_{\varepsilon}, \\ u_{\varepsilon}^b(x) = 0 & \text{for all } x \in \partial\Omega, \\ u_{\varepsilon}^b(x) = U_{\#}^b[\varepsilon](x/\varepsilon) & \text{for all } x \in \varepsilon\partial\omega, \end{cases}$$

respectively. By the Divergence Theorem, we see that

$$\begin{aligned} \text{Cap}_{\Omega}(\varepsilon\overline{\omega}, u^a, u^b) &= \int_{\Omega_{\varepsilon}} \nabla u_{\varepsilon}^a \cdot \nabla u_{\varepsilon}^b \, dx + \int_{\varepsilon\omega} \nabla u^a \cdot \nabla u^b \, dx \\ &= -\varepsilon^{d-2} \int_{\partial\omega} \nu_{\omega}(t) \cdot \nabla(u_{\varepsilon}^a(\varepsilon t)) u^b(\varepsilon t) \, d\sigma_t + \varepsilon^d \int_{\omega} (\nabla u^a)(\varepsilon t) \cdot (\nabla u^b)(\varepsilon t) \, dt, \end{aligned}$$

for all  $\varepsilon \in ]-\varepsilon_{\#}, \varepsilon_{\#}[ \setminus \{0\}$ , where  $\nu_{\omega}$  denotes the outward unit normal to  $\partial\omega$ .

In order to analyze the solution to problem (11) as  $\varepsilon \rightarrow 0$ , we adapt to the present problem the method developed in Dalla Riva, Musolino, and Rogosin [14] for the solution of the Dirichlet problem in a planar perforated domain, which is based on potential theory. By exploiting some specific integral operators (the single and the double layer potentials) we convert a boundary value problem into a set of boundary integral equations.

To introduce the layer potentials, we denote by  $S_d$  the fundamental solution of  $\Delta \equiv \sum_{j=1}^d \partial_j^2$  in  $\mathbb{R}^d$ : i.e.,

$$S_d(x) \equiv \frac{1}{(2-d)s_d|x|^{d-2}} \quad \forall x \in \mathbb{R}^d \setminus \{0\}.$$

Here  $s_d$  denotes the  $(d-1)$ -dimensional measure of the unit sphere in  $\mathbb{R}^d$ . Now let  $\mathcal{O}$  be an open bounded subset of  $\mathbb{R}^d$  of class  $C^{1,\alpha}$ . If  $\phi \in C^{0,\alpha}(\partial\mathcal{O})$ , then we denote by  $v[\partial\mathcal{O}, \phi]$  the single layer potential with density  $\phi$ :

$$v[\partial\mathcal{O}, \phi](x) \equiv \int_{\partial\mathcal{O}} \phi(y) S_d(x-y) \, d\sigma_y \quad \forall x \in \mathbb{R}^d.$$

It is well known that  $v[\partial\mathcal{O}, \phi]$  is a continuous function from  $\mathbb{R}^d$  to  $\mathbb{R}$ . The restriction  $v^+[\partial\mathcal{O}, \phi] \equiv v[\partial\mathcal{O}, \phi]|_{\overline{\mathcal{O}}}$  belongs to  $C^{1,\alpha}(\overline{\mathcal{O}})$ . Moreover, if we denote by  $C_{\text{loc}}^{1,\alpha}(\mathbb{R}^d \setminus \mathcal{O})$  the space of functions on  $\mathbb{R}^d \setminus \mathcal{O}$  whose restrictions to  $\overline{\mathcal{U}}$  belong to  $C^{1,\alpha}(\overline{\mathcal{U}})$  for all open bounded subsets  $\mathcal{U}$  of  $\mathbb{R}^d \setminus \mathcal{O}$ , then  $v^-[\partial\mathcal{O}, \phi] \equiv v[\partial\mathcal{O}, \phi]|_{\mathbb{R}^d \setminus \mathcal{O}}$  belongs to  $C_{\text{loc}}^{1,\alpha}(\mathbb{R}^d \setminus \mathcal{O})$ .

If  $\psi \in C^{1,\alpha}(\partial\mathcal{O})$ , we introduce the double layer potential  $w[\partial\mathcal{O}, \psi]$  by setting

$$w[\partial\mathcal{O}, \psi](x) \equiv - \int_{\partial\mathcal{O}} \psi(y) \nu_{\mathcal{O}}(y) \cdot \nabla S_d(x-y) \, d\sigma_y \quad \forall x \in \mathbb{R}^d,$$

where  $\nu_{\mathcal{O}}$  denotes the outer unit normal to  $\partial\mathcal{O}$ . The restriction  $w[\partial\mathcal{O}, \psi]|_{\mathcal{O}}$  extends to a function  $w^+[\partial\mathcal{O}, \psi]$  of  $C^{1,\alpha}(\overline{\mathcal{O}})$  and the restriction  $w[\partial\mathcal{O}, \psi]|_{\mathbb{R}^d \setminus \overline{\mathcal{O}}}$  extends to a function  $w^-[\partial\mathcal{O}, \psi]$  of  $C_{\text{loc}}^{1,\alpha}(\mathbb{R}^d \setminus \mathcal{O})$ .

To describe the boundary behavior of the trace of the double layer potential on  $\partial\mathcal{O}$  and of the normal derivative of the single layer potential, we define the boundary integral operators  $W_{\mathcal{O}}$  and  $W_{\mathcal{O}}^*$ :

$$W_{\mathcal{O}}[\psi](x) \equiv - \int_{\partial\mathcal{O}} \psi(y) \nu_{\mathcal{O}}(y) \cdot \nabla S_d(x-y) \, d\sigma_y \quad \forall x \in \partial\mathcal{O},$$

for all  $\psi \in C^{1,\alpha}(\partial\mathcal{O})$ , and

$$W_{\mathcal{O}}^*[\phi](x) \equiv \int_{\partial\mathcal{O}} \phi(y) \nu_{\mathcal{O}}(x) \cdot \nabla S_d(x-y) \, d\sigma_y \quad \forall x \in \partial\mathcal{O},$$

for all  $\phi \in C^{0,\alpha}(\partial\mathcal{O})$ . The operators  $W_{\mathcal{O}}$  and  $W_{\mathcal{O}}^*$  are compact operators from  $C^{1,\alpha}(\partial\mathcal{O})$  to itself and from  $C^{0,\alpha}(\partial\mathcal{O})$  to itself, respectively (see Schauder [38,39]). Moreover,  $W_{\mathcal{O}}$  and  $W_{\mathcal{O}}^*$  are adjoint one to the other with respect to the duality on  $C^{1,\alpha}(\partial\mathcal{O}) \times C^{0,\alpha}(\partial\mathcal{O})$  induced by the inner product of the Lebesgue space  $L^2(\partial\mathcal{O})$  (cf., e.g., Kress [22, Chap. 4]). We refer the reader to Kress [22] and Wendland [42,43], for the theory of dual systems and the corresponding Fredholm Alternative Principle. Moreover,

$$\begin{aligned} w^\pm[\partial\mathcal{O}, \psi]_{|\partial\mathcal{O}} &= \pm \frac{1}{2}\psi + W_{\mathcal{O}}[\psi] & \forall \psi \in C^{1,\alpha}(\partial\mathcal{O}), \\ \nu_{\mathcal{O}} \cdot \nabla v^\pm[\partial\mathcal{O}, \phi]_{|\partial\mathcal{O}} &= \mp \frac{1}{2}\phi + W_{\mathcal{O}}^*[\phi] & \forall \phi \in C^{0,\alpha}(\partial\mathcal{O}) \end{aligned}$$

(see, e.g., Folland [19, Chap. 3]).

Finally, we shall need to consider the subspaces of  $C^{0,\alpha}(\partial\mathcal{O})$  and of  $C^{1,\alpha}(\partial\mathcal{O})$ , consisting of functions with zero integral on  $\partial\mathcal{O}$ :

$$C^{k,\alpha}(\partial\mathcal{O})_0 \equiv \left\{ f \in C^{k,\alpha}(\partial\mathcal{O}) : \int_{\partial\mathcal{O}} f \, d\sigma = 0 \right\} \quad \text{for } k = 0, 1. \tag{12}$$

### 2.2. An integral formulation of the boundary value problem

To convert problem (11) into integral equations, we follow the idea of Lanza de Cristoforis [27] and of Dalla Riva, Musolino, and Rogosin [14]. Therefore, we proceed as in [1,3,14] and we introduce the map  $M \equiv (M^o, M^i, M^c)$  from  $] - \varepsilon_{\#}, \varepsilon_{\#}[ \times C^{0,\alpha}(\partial\Omega) \times C^{0,\alpha}(\partial\omega)$  to  $C^{0,\alpha}(\partial\Omega) \times C^{0,\alpha}(\partial\omega)_0 \times \mathbb{R}$  by setting

$$\begin{aligned} M^o[\varepsilon, \rho^o, \rho^i](x) &\equiv \frac{1}{2}\rho^o(x) + W_{\Omega}^*[\rho^o](x) + \int_{\partial\omega} \rho^i(s) \nu_{\Omega}(x) \cdot \nabla S_d(x - \varepsilon s) \, d\sigma_s & \forall x \in \partial\Omega, \\ M^i[\varepsilon, \rho^o, \rho^i](t) &\equiv \frac{1}{2}\rho^i(t) - W_{\omega}^*[\rho^i](t) - \varepsilon^{d-1} \int_{\partial\Omega} \rho^o(y) \nu_{\omega}(t) \cdot \nabla S_d(\varepsilon t - y) \, d\sigma_y & \forall t \in \partial\omega, \\ M^c[\varepsilon, \rho^o, \rho^i] &\equiv \int_{\partial\omega} \rho^i \, d\sigma - 1, \end{aligned}$$

for all  $(\varepsilon, \rho^o, \rho^i) \in ] - \varepsilon_{\#}, \varepsilon_{\#}[ \times C^{0,\alpha}(\partial\Omega) \times C^{0,\alpha}(\partial\omega)$ . The space  $C^{0,\alpha}(\partial\omega)_0$  is defined as in Eq. (12), i.e.,

$$C^{0,\alpha}(\partial\omega)_0 \equiv \left\{ f \in C^{0,\alpha}(\partial\omega) : \int_{\partial\omega} f \, d\sigma = 0 \right\}.$$

Then we can prove the following result of Lanza de Cristoforis [27, §3] (see also Dalla Riva, Musolino, and Rogosin [14, Prop. 4.1]).

**Proposition 2.1.** *The following statements hold.*

- (i) *The map  $M$  is real analytic.*
- (ii) *If  $\varepsilon \in [0, \varepsilon_{\#}[$ , then there exists a unique pair  $(\rho_{\geq}^o[\varepsilon], \rho_{\geq}^i[\varepsilon]) \in C^{0,\alpha}(\partial\Omega) \times C^{0,\alpha}(\partial\omega)$  such that  $M[\varepsilon, \rho_{\geq}^o[\varepsilon], \rho_{\geq}^i[\varepsilon]] = 0$ .*
- (iii) *There exist  $\tilde{\varepsilon}_{\rho} \in ]0, \varepsilon_{\#}[$  and a real analytic map  $(\rho^o[\cdot], \rho^i[\cdot])$  from  $] - \tilde{\varepsilon}_{\rho}, \tilde{\varepsilon}_{\rho}[$  to  $C^{0,\alpha}(\partial\Omega) \times C^{0,\alpha}(\partial\omega)$  such that*

$$M[\varepsilon, \rho^o[\varepsilon], \rho^i[\varepsilon]] = 0 \quad \forall \varepsilon \in ] - \tilde{\varepsilon}_{\rho}, \tilde{\varepsilon}_{\rho}[.$$

*In particular,*

$$(\rho^o[\varepsilon], \rho^i[\varepsilon]) = (\rho_{\geq}^o[\varepsilon], \rho_{\geq}^i[\varepsilon]) \quad \forall \varepsilon \in [0, \tilde{\varepsilon}_{\rho}[.$$

As explained in [27], to represent the solution  $u_\varepsilon$  to problem (11), we need to consider a further operator and thus we define the map  $\Lambda \equiv (\Lambda^o, \Lambda^i)$  from  $] - \varepsilon_\#, \varepsilon_\#[ \times C^{1,\alpha}(\partial\Omega) \times C^{1,\alpha}(\partial\omega)_0$  to  $C^{1,\alpha}(\partial\Omega) \times C^{1,\alpha}(\partial\omega)$  by

$$\begin{aligned} \Lambda^o[\varepsilon, \theta^o, \theta^i](x) &\equiv \frac{1}{2}\theta^o(x) + W_\Omega[\theta^o](x) \\ &\quad + \varepsilon^{d-1} \int_{\partial\omega} \theta^i(s) \nu_\omega(s) \cdot \nabla S_d(x - \varepsilon s) d\sigma_s \quad \forall x \in \partial\Omega, \\ \Lambda^i[\varepsilon, \theta^o, \theta^i](t) &\equiv \frac{1}{2}\theta^i(t) - W_\omega[\theta^i](t) + w[\partial\Omega, \theta^o](\varepsilon t) \\ &\quad - U_\#^a[\varepsilon](t) + \int_{\partial\omega} U_\#^a[\varepsilon]\rho^i[\varepsilon] d\sigma \quad \forall t \in \partial\omega, \end{aligned}$$

for all  $(\varepsilon, \theta^o, \theta^i) \in ] - \varepsilon_\#, \varepsilon_\#[ \times C^{1,\alpha}(\partial\Omega) \times C^{1,\alpha}(\partial\omega)_0$ . Then we have the following result of Lanza de Cristoforis [27, §4] on  $\Lambda$  (cf. Dalla Riva, Musolino, and Rogosin [14, Prop. 4.3]).

**Proposition 2.2.** *The following statements hold.*

- (i) *The map  $\Lambda$  is real analytic.*
- (ii) *If  $\varepsilon \in [0, \varepsilon_\#[$ , then there exists a unique pair  $(\theta^o_\geq[\varepsilon], \theta^i_\geq[\varepsilon]) \in C^{1,\alpha}(\partial\Omega) \times C^{1,\alpha}(\partial\omega)_0$  such that  $\Lambda[\varepsilon, \theta^o_\geq[\varepsilon], \theta^i_\geq[\varepsilon]] = 0$ .*
- (iii) *There exist  $\tilde{\varepsilon}_\theta \in ]0, \varepsilon_\#[$  and a real analytic map  $(\theta^o[\cdot], \theta^i[\cdot])$  from  $] - \tilde{\varepsilon}_\theta, \tilde{\varepsilon}_\theta[$  to  $C^{1,\alpha}(\partial\Omega) \times C^{1,\alpha}(\partial\omega)_0$  such that*

$$\Lambda[\varepsilon, \theta^o[\varepsilon], \theta^i[\varepsilon]] = 0 \quad \forall \varepsilon \in ] - \tilde{\varepsilon}_\theta, \tilde{\varepsilon}_\theta[.$$

*In particular,*

$$(\theta^o[\varepsilon], \theta^i[\varepsilon]) = (\theta^o_\geq[\varepsilon], \theta^i_\geq[\varepsilon]) \quad \forall \varepsilon \in [0, \tilde{\varepsilon}_\theta[.$$

We now set

$$\varepsilon_0 \equiv \min\{\tilde{\varepsilon}_\rho, \tilde{\varepsilon}_\theta\}.$$

By summing suitable double and single layer potentials, by a modification of the argument of Dalla Riva, Musolino, and Rogosin [14, Prop. 4.5], we can represent the rescaled function  $u_\varepsilon(\varepsilon t)$  by means of the functions  $\rho^o[\varepsilon]$ ,  $\rho^i[\varepsilon]$ ,  $\theta^o[\varepsilon]$ , and  $\theta^i[\varepsilon]$  introduced in Propositions 2.1 and 2.2 (see also Lanza de Cristoforis [27, §5] and Dalla Riva, Musolino, and Rogosin [13, §2.4]).

**Proposition 2.3.** *Let  $\varepsilon \in ]0, \varepsilon_0[$ . Then*

$$\begin{aligned} u_\varepsilon^a(\varepsilon t) &\equiv w^+[\partial\Omega, \theta^o[\varepsilon]](\varepsilon t) - w^-[\partial\omega, \theta^i[\varepsilon]](t) \\ &\quad + \int_{\partial\omega} U_\#^a[\varepsilon]\rho^i[\varepsilon] d\sigma \left( v^+[\partial\Omega, \rho^o[\varepsilon]](\varepsilon t) + \varepsilon^{-(d-2)}v^-[\partial\omega, \rho^i[\varepsilon]](t) \right) \\ &\quad \times \left( \frac{1}{\int_{\partial\omega} d\sigma} \int_{\partial\omega} v[\partial\Omega, \rho^o[\varepsilon]](\varepsilon s) + \varepsilon^{-(d-2)}v[\partial\omega, \rho^i[\varepsilon]](s) d\sigma_s \right)^{-1} \\ &\equiv w^+[\partial\Omega, \theta^o[\varepsilon]](\varepsilon t) - w^-[\partial\omega, \theta^i[\varepsilon]](t) \\ &\quad + \int_{\partial\omega} U_\#^a[\varepsilon]\rho^i[\varepsilon] d\sigma \left( \varepsilon^{d-2}v^+[\partial\Omega, \rho^o[\varepsilon]](\varepsilon t) + v^-[\partial\omega, \rho^i[\varepsilon]](t) \right) \\ &\quad \times \left( \frac{1}{\int_{\partial\omega} d\sigma} \int_{\partial\omega} \varepsilon^{d-2}v[\partial\Omega, \rho^o[\varepsilon]](\varepsilon s) + v[\partial\omega, \rho^i[\varepsilon]](s) d\sigma_s \right)^{-1} \end{aligned}$$

for all  $t \in \overline{(\varepsilon^{-1}\Omega)} \setminus \omega$ .

### 3. Power series expansions of the auxiliary functions $(\rho^o[\varepsilon], \rho^i[\varepsilon])$ and $(\theta^o[\varepsilon], \theta^i[\varepsilon])$ around $\varepsilon = 0$

As in [3], now the plan is to construct an expansion for  $\nu_\omega(t) \cdot \nabla(u_\varepsilon^a(\varepsilon t))u^b(\varepsilon t)$  and then to integrate it on  $\partial\omega$ . To do so, we first obtain a representation of the auxiliary density functions  $(\rho^o[\varepsilon], \rho^i[\varepsilon])$  and  $(\theta^o[\varepsilon], \theta^i[\varepsilon])$  and then we use it to represent  $\nu_\omega(t) \cdot \nabla(u_\varepsilon^a(\varepsilon t))u^b(\varepsilon t)$ .

To compute the coefficients of the series involved in the representation, we need explicit expressions for the derivatives with respect to  $\varepsilon$  of functions of the type  $F(\varepsilon x)$ . More precisely, we will exploit the equality

$$\partial_\varepsilon^j(F(\varepsilon x)) = \sum_{\substack{\beta \in \mathbb{N}^d \\ |\beta|=j}} \frac{j!}{\beta!} x^\beta (D^\beta F)(\varepsilon x) \tag{13}$$

which holds for all  $j \in \mathbb{N}$ ,  $\varepsilon \in \mathbb{R}$ ,  $x \in \mathbb{R}^d$ , and for all functions  $F$  analytic in a neighborhood of  $\varepsilon x$ . Here, if  $\alpha \in \mathbb{N}^d$ , then  $(D^\alpha F)(y)$  denotes the partial derivative of multi-index  $\alpha$  with respect to  $x$  of the function  $F(x)$  evaluated at  $y \in \mathbb{R}^d$ . We also exploit formulas for the derivatives of layer potentials in the singularly perturbed set  $\Omega \setminus \varepsilon\bar{\omega}$  with respect to the parameter  $\varepsilon$  as it is done in Dalla Riva, Luzzini and Musolino [12].

Then we have the following variant of Dalla Riva, Musolino, and Rogosin [14, Prop. 5.1], where we represent  $(\rho^o[\varepsilon], \rho^i[\varepsilon])$  as a power series for  $\varepsilon$  close to 0.

**Proposition 3.1.** *Let  $(\rho^o[\varepsilon], \rho^i[\varepsilon])$  be as in Proposition 2.1 for all  $\varepsilon \in ]-\varepsilon_0, \varepsilon_0[$ . Then there exist  $\varepsilon_\rho \in ]0, \varepsilon_0[$  and a sequence  $\{(\rho_k^o, \rho_k^i)\}_{k \in \mathbb{N}}$  in  $C^{0,\alpha}(\partial\Omega) \times C^{0,\alpha}(\partial\omega)$  such that*

$$\rho^o[\varepsilon] = \sum_{k=0}^{+\infty} \frac{\rho_k^o}{k!} \varepsilon^k \quad \text{and} \quad \rho^i[\varepsilon] = \sum_{k=0}^{+\infty} \frac{\rho_k^i}{k!} \varepsilon^k \quad \forall \varepsilon \in ]-\varepsilon_\rho, \varepsilon_\rho[ \tag{14}$$

where the two series converge normally in  $C^{0,\alpha}(\partial\Omega)$  and in  $C^{0,\alpha}(\partial\omega)$ , respectively, for  $\varepsilon \in ]-\varepsilon_\rho, \varepsilon_\rho[$ . Moreover, the pair of functions  $(\rho_0^o, \rho_0^i)$  is the unique solution in  $C^{0,\alpha}(\partial\Omega) \times C^{0,\alpha}(\partial\omega)$  of the following system of integral equations

$$\begin{aligned} \frac{1}{2} \rho_0^o(x) + W_\Omega^*[\rho_0^o](x) &= -\nu_\Omega(x) \cdot \nabla S_d(x) & \forall x \in \partial\Omega, \\ \frac{1}{2} \rho_0^i(t) - W_\omega^*[\rho_0^i](t) &= 0 & \forall t \in \partial\omega, \\ \int_{\partial\omega} \rho_0^i \, d\sigma &= 1, \end{aligned}$$

for each  $k \in \{1, \dots, d-2\}$  the pair of functions  $(\rho_k^o, \rho_k^i)$  is the unique solution in  $C^{0,\alpha}(\partial\Omega) \times C^{0,\alpha}(\partial\omega)$  of the following system of integral equations

$$\begin{aligned} \frac{1}{2} \rho_k^o(x) + W_\Omega^*[\rho_k^o](x) &= (-1)^{k+1} \sum_{\substack{\beta \in \mathbb{N}^d \\ |\beta|=k}} \frac{k!}{\beta!} \nu_\Omega(x) \cdot (\nabla D^\beta S)(x) \int_{\partial\omega} \rho_0^i(s) s^\beta \, d\sigma_s & \forall x \in \partial\Omega, \\ \rho_k^i(t) &= 0 & \forall t \in \partial\omega, \end{aligned}$$

and for each  $k \in \mathbb{N} \setminus \{0, \dots, d - 2\}$  the pair  $(\rho_k^o, \rho_k^i)$  is the unique solution in  $C^{0,\alpha}(\partial\Omega) \times C^{0,\alpha}(\partial\omega)$  of the following system of integral equations which involves  $\{(\rho_j^o, \rho_j^i)\}_{j=0}^{k-1}$ ,

$$\begin{aligned} & \frac{1}{2} \rho_k^o(x) + W_\Omega^*[\rho_k^o](x) \\ &= \sum_{j=0}^k \binom{k}{j} (-1)^{j+1} \sum_{\substack{\beta \in \mathbb{N}^d \\ |\beta|=j}} \frac{j!}{\beta!} \nu_\Omega(x) \cdot (\nabla D^\beta S)(x) \int_{\partial\omega} \rho_{k-j}^i(s) s^\beta d\sigma_s \quad \forall x \in \partial\Omega, \\ & \frac{1}{2} \rho_k^i(t) - W_\omega^*[\rho_k^i](t) \\ &= \frac{k!}{(k - (d - 1))!} \sum_{j=0}^{k-(d-1)} \binom{k - (d - 1)}{j} \sum_{\substack{\beta \in \mathbb{N}^d \\ |\beta|=j}} \frac{j!}{\beta!} (-1)^{j+1} t^\beta \nu_\omega(t) \cdot \int_{\partial\omega} \rho_{k-(d-1)-j}^o \nabla D^\beta S_d d\sigma \\ & \quad \forall t \in \partial\omega, \\ & \int_{\partial\omega} \rho_k^i d\sigma = 0. \end{aligned}$$

**Proof.**

By the real analyticity result of Proposition 2.1 (iii) for the map

$$\varepsilon \mapsto (\rho^o[\varepsilon], \rho^i[\varepsilon]),$$

we deduce that there exist  $\varepsilon_\rho$  and  $\{(\rho_k^o, \rho_k^i)\}_{k \in \mathbb{N}}$  such that (14) holds. By the real analyticity of  $\varepsilon \mapsto (\rho^o[\varepsilon], \rho^i[\varepsilon])$ , we have  $(\rho_k^o, \rho_k^i) = (\partial_\varepsilon^k \rho^o[0], \partial_\varepsilon^k \rho^i[0])$  for all  $k \in \mathbb{N}$ . Therefore our goal is to identify the derivatives  $(\partial_\varepsilon^k \rho^o[0], \partial_\varepsilon^k \rho^i[0])$  for all  $k \in \mathbb{N}$ . Equality  $M[\varepsilon, \rho^o[\varepsilon], \rho^i[\varepsilon]] = 0$  for all  $\varepsilon \in ] - \varepsilon_0, \varepsilon_0[$  (cf. Proposition 2.1 (ii)) implies that the map

$$\varepsilon \mapsto M[\varepsilon, \rho^o[\varepsilon], \rho^i[\varepsilon]]$$

has zero derivatives, i.e.,

$$\partial_\varepsilon^k (M[\varepsilon, \rho^o[\varepsilon], \rho^i[\varepsilon]]) = 0 \quad \forall \varepsilon \in ] - \varepsilon_0, \varepsilon_0[, k \in \mathbb{N}. \tag{15}$$

Therefore we compute  $\partial_\varepsilon^k (M[\varepsilon, \rho^o[\varepsilon], \rho^i[\varepsilon]])$  and we have

$$\begin{aligned} \partial_\varepsilon^k (M^o[\varepsilon, \rho^o[\varepsilon], \rho^i[\varepsilon]])(x) &= \frac{1}{2} \partial_\varepsilon^k \rho^o[\varepsilon](x) + W_\Omega^*[\partial_\varepsilon^k \rho^o[\varepsilon]](x) \\ &+ \sum_{j=0}^k \binom{k}{j} (-1)^j \sum_{\substack{\beta \in \mathbb{N}^d \\ |\beta|=j}} \frac{j!}{\beta!} \nu_\Omega(x) \cdot \int_{\partial\omega} \partial_\varepsilon^{k-j} \rho^i[\varepsilon](s) s^\beta (\nabla D^\beta S_d)(x - \varepsilon s) d\sigma_s \\ & \quad \forall x \in \partial\Omega, \end{aligned} \tag{16}$$

$$\begin{aligned} \partial_\varepsilon^k (M^i[\varepsilon, \rho^o[\varepsilon], \rho^i[\varepsilon]])(t) &= \frac{1}{2} \partial_\varepsilon^k \rho^i[\varepsilon](t) - W_\omega^*[\partial_\varepsilon^k \rho^i[\varepsilon]](t) \\ &- \partial_\varepsilon^k \left( \varepsilon^{d-1} \int_{\partial\Omega} \rho^o[\varepsilon](y) \nu_\omega(t) \cdot \nabla S_d(\varepsilon t - y) d\sigma_y \right) \\ & \quad \forall t \in \partial\omega, \end{aligned} \tag{17}$$

$$\partial_\varepsilon^k (M^c[\varepsilon, \rho^o[\varepsilon], \rho^i[\varepsilon]]) = \int_{\partial\omega} \partial_\varepsilon^k \rho^i[\varepsilon] d\sigma - \delta_{0k}, \tag{18}$$

for all  $\varepsilon \in ] - \varepsilon_0, \varepsilon_0[$  and all  $k \in \mathbb{N}$  (see also (13)). Here above  $\delta_{ij}$  denotes the Kronecker delta function. Next we note that if  $k < d - 1$  then

$$\partial_\varepsilon^k \left( \varepsilon^{d-1} \int_{\partial\Omega} \rho^o[\varepsilon](y) \nu_\omega(t) \cdot \nabla S_d(\varepsilon t - y) d\sigma_y \right) \Big|_{\varepsilon=0} = 0 \quad \forall t \in \partial\omega.$$

Instead, if  $k \geq d - 1$  then

$$\begin{aligned} & \partial_\varepsilon^k \left( \varepsilon^{d-1} \int_{\partial\Omega} \rho^\circ[\varepsilon](y) \nu_\omega(t) \cdot \nabla S_d(\varepsilon t - y) d\sigma_y \right) \Big|_{\varepsilon=0} \\ &= \binom{k}{(d-1)} (d-1)! \partial_\varepsilon^{k-(d-1)} \left( \int_{\partial\Omega} \rho^\circ[\varepsilon](y) \nu_\omega(t) \cdot \nabla S_d(\varepsilon t - y) d\sigma_y \right) \Big|_{\varepsilon=0} \\ &= \frac{k!}{(k-(d-1))!} \sum_{j=0}^{k-(d-1)} \binom{k-(d-1)}{j} \\ &\quad \times \sum_{\substack{\beta \in \mathbb{N}^d \\ |\beta|=j}} \frac{j!}{\beta!} \int_{\partial\omega} \partial_\varepsilon^{k-(d-1)-j} (\rho^\circ[\varepsilon](y)) \Big|_{\varepsilon=0} t^\beta \nu_\omega(t) \cdot \nabla D^\beta S_d(-y) d\sigma_y \\ &= \frac{k!}{(k-(d-1))!} \sum_{j=0}^{k-(d-1)} \binom{k-(d-1)}{j} \\ &\quad \times \sum_{\substack{\beta \in \mathbb{N}^d \\ |\beta|=j}} \frac{j!}{\beta!} (-1)^{|\beta|+1} \int_{\partial\omega} \partial_\varepsilon^{k-(d-1)-j} (\rho^\circ[\varepsilon](y)) \Big|_{\varepsilon=0} t^\beta \nu_\omega(t) \cdot \nabla D^\beta S_d(y) d\sigma_y \quad \forall t \in \partial\omega. \end{aligned}$$

Then, by taking  $\varepsilon = 0$  in (16)–(18) and by equality (15), we deduce that for each  $k \in \mathbb{N}$  the pair of functions  $(\rho_k^\circ, \theta_k^i)$  is a solution of the corresponding integral equations of the statement. By the characterization of the kernels of the operators of Dalla Riva, Lanza de Cristoforis, and Musolino [11, §6.5 and §6.6], we deduce the uniqueness of the solutions of the integral equations of the statement.  $\square$

By a similar computation, in the following proposition we construct the power series expansion of  $(\theta^\circ[\varepsilon], \theta^i[\varepsilon])$ .

**Proposition 3.2.** *Let  $(\theta^\circ[\varepsilon], \theta^i[\varepsilon])$  be as in Proposition 2.2 for all  $\varepsilon \in ]-\varepsilon_0, \varepsilon_0[$ . Then there exist  $\varepsilon_\theta \in ]0, \varepsilon_0[$  and a sequence  $\{(\theta_k^\circ, \theta_k^i)\}_{k \in \mathbb{N}}$  in  $C^{1,\alpha}(\partial\Omega) \times C^{1,\alpha}(\partial\omega)_0$  such that*

$$\theta^\circ[\varepsilon] = \sum_{k=0}^\infty \frac{\theta_k^\circ}{k!} \varepsilon^k \quad \text{and} \quad \theta^i[\varepsilon] = \sum_{k=0}^\infty \frac{\theta_k^i}{k!} \varepsilon^k \quad \forall \varepsilon \in ]-\varepsilon_\theta, \varepsilon_\theta[, \tag{19}$$

where the two series converge normally in  $C^{1,\alpha}(\partial\Omega)$  and in  $C^{1,\alpha}(\partial\omega)_0$ , respectively, for  $\varepsilon \in ]-\varepsilon_\theta, \varepsilon_\theta[$ . Moreover,

$$(\theta_0^\circ, \theta_0^i) = (0, 0), \quad \theta_k^\circ = 0 \quad \forall k \in \{0, \dots, d-1\},$$

for each  $k \in \{0, \dots, d-1\}$  the function  $\theta_k^i$  is the unique solution in  $C^{1,\alpha}(\partial\omega)_0$  of

$$\frac{1}{2} \theta_k^i(t) - W_\omega[\theta_k^i](t) = \sum_{|\beta|=k} \frac{k!}{\beta!} t^\beta (D^\beta u^a)(0) - \sum_{l=0}^k \sum_{\substack{\beta \in \mathbb{N}^d \\ |\beta|=l}} \binom{k}{l} \frac{l!}{\beta!} \int_{\partial\omega} s^\beta (D^\beta u^a)(0) \rho_{k-l}^i(s) d\sigma_s \quad \forall t \in \partial\omega, \tag{20}$$

and for each  $k \in \mathbb{N} \setminus \{0, \dots, d-1\}$  the pair  $(\theta_k^\circ, \theta_k^i)$  is the unique solution in  $C^{1,\alpha}(\partial\Omega) \times C^{1,\alpha}(\partial\omega)_0$  of the following system of integral equations which involves  $\{(\theta_j^\circ, \theta_j^i)\}_{j=0}^{k-1}$ ,

$$\begin{aligned} & \frac{1}{2} \theta_k^\circ(x) + W_\Omega[\theta_k^\circ](x) \tag{21} \\ &= \frac{k!}{(k-(d-1))!} \sum_{j=0}^{k-(d-1)-1} \binom{k-(d-1)}{j} (-1)^{j+1} \sum_{|\beta|=j} \frac{j!}{\beta!} (\nabla D^\beta S_d)(x) \cdot \int_{\partial\omega} \theta_{k-(d-1)-j}^i(s) \nu_\omega(s) s^\beta d\sigma_s \\ &\quad \forall x \in \partial\Omega, \end{aligned}$$

$$\begin{aligned} \frac{1}{2}\theta_k^i(t) - W_\omega[\theta_k^i](t) &= \sum_{j=0}^{k-(d-1)} \binom{k}{j} (-1)^{j+1} \sum_{|\beta|=j} \frac{j!}{\beta!} t^\beta \int_{\partial\Omega} \theta_{k-j}^o \nu_\Omega \cdot \nabla D^\beta S_d \, d\sigma \\ &+ \sum_{|\beta|=k} \frac{k!}{\beta!} t^\beta (D^\beta u^a)(0) - \sum_{l=0}^k \sum_{\substack{\beta \in \mathbb{N}^d \\ |\beta|=l}} \binom{k}{l} \frac{l!}{\beta!} \int_{\partial\omega} s^\beta (D^\beta u^a)(0) \rho_{k-l}^i(s) \, d\sigma_s \quad \forall t \in \partial\omega. \end{aligned} \tag{22}$$

**Proof.** We proceed as in Dalla Riva, Musolino, and Rogosin [14, Prop. 5.2] and in [1, Prop 3.2]. Proposition 2.2 (iii) implies that the map

$$\varepsilon \mapsto (\theta^o[\varepsilon], \theta^i[\varepsilon])$$

is real analytic. As a consequence, there exist  $\varepsilon_\theta$  and  $\{(\theta_k^o, \theta_k^i)\}_{k \in \mathbb{N}}$  such that (19) holds. Our goal is to identify the terms  $(\theta_k^o, \theta_k^i)$  for each  $k \in \mathbb{N}$  and we do that by computing the derivatives of  $(\theta^o[\varepsilon], \theta^i[\varepsilon])$  with respect to  $\varepsilon$ . By Proposition 2.2 (ii), we have

$$A[\varepsilon, \theta^o[\varepsilon], \theta^i[\varepsilon]] = 0 \quad \forall \varepsilon \in ]-\varepsilon_0, \varepsilon_0[.$$

We can compute the derivative with respect to  $\varepsilon$  in the equality above and we deduce that

$$\partial_\varepsilon^k (A[\varepsilon, \theta^o[\varepsilon], \theta^i[\varepsilon]]) = 0 \quad \forall \varepsilon \in ]-\varepsilon_0, \varepsilon_0[, \forall k \in \mathbb{N}. \tag{23}$$

Hence we compute  $\partial_\varepsilon^k (A[\varepsilon, \theta^o[\varepsilon], \theta^i[\varepsilon]])$  and we have

$$\begin{aligned} \partial_\varepsilon^k (A^o[\varepsilon, \theta^o[\varepsilon], \theta^i[\varepsilon]])(x) &= \frac{1}{2} \partial_\varepsilon^k \theta^o[\varepsilon](x) + W_\Omega[\partial_\varepsilon^k \theta^o[\varepsilon]](x) \\ &+ \partial_\varepsilon^k (\varepsilon^{d-1} \int_{\partial\omega} \theta^i[\varepsilon](s) \nu_\omega(s) \cdot \nabla S_d(x - \varepsilon s) \, d\sigma_s) \quad \forall x \in \partial\Omega, \end{aligned} \tag{24}$$

$$\begin{aligned} \partial_\varepsilon^k (A^i[\varepsilon, \theta^o[\varepsilon], \theta^i[\varepsilon]])(t) &= \frac{1}{2} \partial_\varepsilon^k \theta^i[\varepsilon](t) - W_\omega[\partial_\varepsilon^k \theta^i[\varepsilon]](t) \\ &- \sum_{j=0}^k \binom{k}{j} \sum_{|\beta|=j} \frac{j!}{\beta!} t^\beta \int_{\partial\Omega} \partial_\varepsilon^{k-j} \theta^o[\varepsilon](y) \nu_\Omega(y) \cdot (\nabla D^\beta S_d)(\varepsilon t - y) \, d\sigma_y \\ &- \sum_{|\beta|=k} \frac{k!}{\beta!} t^\beta (D^\beta u^a)(\varepsilon t) \\ &+ \sum_{l=0}^k \sum_{\substack{\beta \in \mathbb{N}^d \\ |\beta|=l}} \binom{k}{l} \frac{l!}{\beta!} \int_{\partial\omega} s^\beta (D^\beta u^a)(\varepsilon s) \partial_\varepsilon^{k-l} \rho^i[\varepsilon](s) \, d\sigma_s \quad \forall t \in \partial\omega, \end{aligned} \tag{25}$$

for all  $\varepsilon \in ]-\varepsilon_0, \varepsilon_0[$  and all  $k \in \mathbb{N}$ . Then we note that if  $k < d - 1$  then

$$\partial_\varepsilon^k \left( \varepsilon^{d-1} \int_{\partial\omega} \theta^i[\varepsilon](s) \nu_\omega(s) \cdot \nabla S_d(x - \varepsilon s) \, d\sigma_s \right) \Big|_{\varepsilon=0} = 0 \quad \forall x \in \partial\Omega.$$

Instead, if  $k \geq d - 1$  then

$$\begin{aligned} &\partial_\varepsilon^k \left( \varepsilon^{d-1} \int_{\partial\omega} \theta^i[\varepsilon](s) \nu_\omega(s) \cdot \nabla S_d(x - \varepsilon s) \, d\sigma_s \right) \Big|_{\varepsilon=0} \\ &= \binom{k}{(d-1)} (d-1)! \partial_\varepsilon^{k-(d-1)} \left( \int_{\partial\omega} \theta^i[\varepsilon](s) \nu_\omega(s) \cdot \nabla S_d(x - \varepsilon s) \, d\sigma_s \right) \Big|_{\varepsilon=0} \\ &= \frac{k!}{(k-(d-1))!} \sum_{j=0}^{k-(d-1)} \binom{k-(d-1)}{j} (-1)^j \\ &\quad \times \sum_{|\beta|=j} \frac{j!}{\beta!} \int_{\partial\omega} \partial_\varepsilon^{k-(d-1)-j} (\theta^i[\varepsilon](s)) \Big|_{\varepsilon=0} s^\beta \nu_\omega(s) \cdot \nabla D^\beta S_d(x) \, d\sigma_s \quad \forall x \in \partial\Omega. \end{aligned}$$

The real analyticity of  $\varepsilon \mapsto (\theta^\circ[\varepsilon], \theta^i[\varepsilon])$  implies that  $(\theta_k^\circ, \theta_k^i) = (\partial_\varepsilon^k \theta^\circ[0], \partial_\varepsilon^k \theta^i[0])$  for all  $k \in \mathbb{N}$ . Therefore, by taking  $\varepsilon = 0$  in (24) and (25) and by equality (23), we deduce that  $(\theta_0^\circ, \theta_0^i) = (0, 0)$ , that  $\theta_k^\circ = 0$  for all  $k \in \{0, \dots, d-1\}$ , that for each  $k \in \{0, \dots, d-1\}$  the function  $\theta_k^i$  solves Eq. (20), and that  $(\theta_k^\circ, \theta_k^i)$  is a solution of (21) and (22) for all  $k \in \mathbb{N} \setminus \{0, \dots, d-1\}$ . To conclude, we note that, by classical potential theory (cf., e.g., Dalla Riva, Lanza de Cristoforis, and Musolino [11, §6.5 and §6.6]), Eq. (20) has a unique solution in  $C^{1,\alpha}(\partial\omega)_0$  and Eqs. (21)–(22) have a unique solution in  $C^{1,\alpha}(\partial\Omega) \times C^{1,\alpha}(\partial\omega)_0$ .  $\square$

**4. Series expansion of  $\nu_\omega(\cdot) \cdot \nabla(u_\varepsilon^a(\varepsilon \cdot))u^b(\varepsilon \cdot)$  around  $\varepsilon = 0$**

The next step is to obtain a series expansion of  $\nu_\omega(\cdot) \cdot \nabla(u_\varepsilon^a(\varepsilon \cdot))u^b(\varepsilon \cdot)$  for  $\varepsilon$  close to 0. The coefficients of such series will be defined by means of the sequences  $\{(\rho_k^\circ, \rho_k^i)\}_{k \in \mathbb{N}}$  and  $\{(\theta_k^\circ, \theta_k^i)\}_{k \in \mathbb{N}}$  introduced in Section 3. We begin with the intermediate result of Proposition 4.1 below, whose proof can be obtained by arguing as in the proof of [3, Prop. 2.9].

**Proposition 4.1.** *Let  $\{(\rho_k^\circ, \rho_k^i)\}_{k \in \mathbb{N}}$  and  $\{(\theta_k^\circ, \theta_k^i)\}_{k \in \mathbb{N}}$  be as in Propositions 3.1 and 3.2, respectively. Let*

$$\begin{aligned} u_{m,0}^a(t) &\equiv 0 && \forall t \in \mathbb{R}^d \setminus \omega, \\ u_{m,k}^a(t) &\equiv -w^-[\partial\omega, \theta_k^i](t) && \forall t \in \mathbb{R}^d \setminus \omega, k \in \{0, \dots, d-1\}, \\ u_{m,k}^a(t) &\equiv \frac{1}{k!} \sum_{j=0}^{k-d} \binom{k}{j} (-1)^j \sum_{\substack{\beta \in \mathbb{N}^d \\ |\beta|=j}} \frac{j!}{\beta!} t^\beta \int_{\partial\Omega} \theta_{k-j}^\circ \nu_\Omega \cdot (\nabla D^\beta S_d) d\sigma \\ &\quad - \frac{1}{k!} w^-[\partial\omega, \theta_k^i](t) && \forall t \in \mathbb{R}^d \setminus \omega, \quad \forall k \geq d \end{aligned}$$

and

$$\begin{aligned} v_{m,k}(t) &\equiv \frac{1}{(k-(d-2))!} \sum_{j=0}^{k-(d-2)} \binom{k-(d-2)}{j} (-1)^j \sum_{\substack{\beta \in \mathbb{N}^d \\ |\beta|=j}} \frac{j!}{\beta!} t^\beta \int_{\partial\Omega} \rho_{k-(d-2)-j}^\circ D^\beta S_d d\sigma + \frac{1}{k!} v^-[\partial\omega, \rho_k^i](t) \\ &&& \forall t \in \mathbb{R}^d \setminus \omega, \\ g_k^a &\equiv \frac{1}{k!} \sum_{l=0}^k \sum_{\substack{\beta \in \mathbb{N}^d \\ |\beta|=l}} \binom{k}{l} \frac{l!}{\beta!} \int_{\partial\omega} s^\beta (D^\beta u^a)(0) \rho_{k-l}^i(s) d\sigma_s, \\ r_k &\equiv \frac{1}{(k-(d-2))! \int_{\partial\omega} d\sigma} \sum_{j=0}^{k-(d-2)} \binom{k-(d-2)}{j} (-1)^j \sum_{\substack{\beta \in \mathbb{N}^d \\ |\beta|=j}} \frac{j!}{\beta!} \int_{\partial\omega} s^\beta d\sigma_s \int_{\partial\Omega} \rho_{k-(d-2)-j}^\circ D^\beta S_d d\sigma \\ &\quad + \frac{1}{k! \int_{\partial\omega} d\sigma} \int_{\partial\omega} v[\partial\omega, \rho_k^i] d\sigma, \end{aligned}$$

for all  $k \in \mathbb{N}$ . Here above the sum  $\sum_{j=0}^{k-(d-2)}$  is omitted if  $k - (d - 2) < 0$ . Let

$$\begin{aligned} u_{\#,k}^l(t) &\equiv \sum_{\substack{\beta \in \mathbb{N}^d \\ |\beta|=k}} \frac{D^\beta u^l(0)}{\beta!} t^\beta && \forall t \in \mathbb{R}^d, \quad l = a, b, \\ \tilde{u}_k(t) &\equiv \sum_{l=0}^k \nu_\omega(t) \cdot \nabla u_{m,l}^a|_{\partial\omega}(t) u_{\#,k-l}^b(t) && \forall t \in \partial\omega, \\ \tilde{v}_k(t) &\equiv \nu_\omega(t) \cdot \nabla v_{m,k}|_{\partial\omega}(t) && \forall t \in \partial\omega, \\ \tilde{g}_k(t) &\equiv \sum_{l=0}^k g_l^a u_{\#,k-l}^b(t) && \forall t \in \partial\omega, \end{aligned}$$



for all  $k \in \mathbb{N}$ . Then there exists  $\tilde{\varepsilon} \in ]0, \varepsilon_0]$  such that

$$\nu_\omega(\cdot) \cdot \nabla(u_\varepsilon^a(\varepsilon \cdot))|_{\partial\omega} u^b(\varepsilon \cdot)|_{\partial\omega} = \sum_{k=1}^\infty \tilde{u}_k(\cdot) \varepsilon^k + \left( \sum_{k=0}^\infty \tilde{g}_k(\cdot) \varepsilon^k \right) \frac{\sum_{k=0}^\infty \tilde{v}_k(\cdot) \varepsilon^k}{\sum_{k=0}^\infty r_k \varepsilon^k} \tag{26}$$

for all  $\varepsilon \in ]-\tilde{\varepsilon}, \tilde{\varepsilon}[\setminus\{0\}$ . Moreover, the series  $\sum_{k=0}^\infty \tilde{g}_k \varepsilon^k$ ,  $\sum_{k=1}^\infty \tilde{u}_k \varepsilon^k$ , and  $\sum_{k=0}^\infty \tilde{v}_k \varepsilon^k$  converge normally in  $C^{0,\alpha}(\partial\omega)$  for  $\varepsilon \in ]-\tilde{\varepsilon}, \tilde{\varepsilon}[$  and  $\sum_{k=0}^\infty r_k \varepsilon^k$  converge absolutely in  $]-\tilde{\varepsilon}, \tilde{\varepsilon}[$ .

We would like to have a representation formula for  $\nu_\omega(\cdot) \cdot \nabla(u_\varepsilon^a(\varepsilon \cdot))|_{\partial\omega} u^b(\varepsilon \cdot)|_{\partial\omega}$  in the form of a convergent power series of the type  $\sum_{n=0}^\infty \varphi_\varepsilon(\cdot) \varepsilon^n$ . By exploiting an argument similar to that of Dalla Riva, Musolino, and Rogosin [14, Thm. 6.3], we can prove Theorem 4.2 below where we obtain from formula (26) a series expansion for the map which takes  $\varepsilon$  to  $\nu_\omega(\cdot) \cdot \nabla(u_\varepsilon^a(\varepsilon \cdot))|_{\partial\omega} u^b(\varepsilon \cdot)|_{\partial\omega}$  (see also [1, Thm. 2.10] and [3, Thm. 4.3]).

**Theorem 4.2.** *With the notation introduced in Proposition 4.1, let  $\{\tilde{a}_n\}_{n \in \mathbb{N}}$  be the sequence of functions from  $\partial\omega$  to  $\mathbb{R}$  defined by*

$$\tilde{a}_n \equiv \sum_{k=0}^n \tilde{g}_{n-k} \tilde{v}_k \quad \forall n \in \mathbb{N}.$$

Let  $\{\tilde{\lambda}_n\}_{n \in \mathbb{N}}$  be the sequence of functions from  $\partial\omega$  to  $\mathbb{R}$  defined by

$$\tilde{\lambda}_0 \equiv \tilde{a}_0/r_0, \quad \tilde{\lambda}_n \equiv \tilde{u}_n + \tilde{a}_n/r_0 + \sum_{k=1}^n \tilde{a}_{n-k} \sum_{j=1}^k \frac{(-1)^j}{r_0^{j+1}} \sum_{\substack{\beta \in (\mathbb{N} \setminus \{0\})^j \\ |\beta|=k}} \prod_{h=1}^j r_{\beta_h} \quad \forall n \geq 1.$$

Then there exists  $\tilde{\varepsilon}' \in ]0, \varepsilon_0]$  such that

$$\nu_\omega(\cdot) \cdot \nabla(u_\varepsilon^a(\varepsilon \cdot))|_{\partial\omega} u^b(\varepsilon \cdot)|_{\partial\omega} = \sum_{n=0}^\infty \tilde{\lambda}_n(\cdot) \varepsilon^n \tag{27}$$

for all  $\varepsilon \in ]-\tilde{\varepsilon}', \tilde{\varepsilon}'[\setminus\{0\}$ . Moreover, the series

$$\sum_{n=0}^\infty \tilde{\lambda}_n(\cdot) \varepsilon^n$$

converges normally in  $C^{0,\alpha}(\partial\omega)$  for  $\varepsilon \in ]-\tilde{\varepsilon}', \tilde{\varepsilon}'[$  and

$$\tilde{\lambda}_0 = \frac{u^a(0)u^b(0)}{r_0} \frac{\partial}{\partial\nu_\omega} v^-[\partial\omega, \rho_0^i].$$

### 5. Series expansion of $\text{Cap}_\Omega(\varepsilon\bar{\omega}, u^a, u^b)$

We recall that the  $(u^a, u^b)$ -capacity  $\text{Cap}_\Omega(\varepsilon\bar{\omega}, u^a, u^b)$  can be represented as the sum of  $\int_{\Omega_\varepsilon} \nabla u_\varepsilon^a \cdot \nabla u_\varepsilon^b dx$  and of  $\int_{\varepsilon\omega} \nabla u^a \cdot \nabla u^b dx$ . Therefore, in order to compute a series expansion of  $\text{Cap}_\Omega(\varepsilon\bar{\omega}, u^a, u^b)$ , we begin by providing an expansion for  $\int_{\varepsilon\omega} \nabla u^a \cdot \nabla u^b dx$  around  $\varepsilon = 0$ .

**Lemma 5.1.** *Let  $\{\xi_n\}_{n \in \mathbb{N}}$  be the sequence of real numbers defined by*

$$\xi_0 \equiv \xi_1 \equiv \dots \equiv \xi_{d-1} \equiv 0, \quad \xi_n \equiv \sum_{j=1}^d \sum_{l=0}^{n-d} \int_\omega \partial_j u_{\#,l+1}^a(t) \partial_j u_{\#,n-l-(d-1)}^b(t) dt \quad \forall n \geq d.$$

Then there exists  $\varepsilon_\xi \in ]0, \varepsilon_0]$  such that

$$\int_{\varepsilon\omega} \nabla u^a \cdot \nabla u^b \, dx = \sum_{n=d}^\infty \xi_n \varepsilon^n$$

for all  $\varepsilon \in ]0, \varepsilon_\xi[$ . Moreover,

$$\xi_d = \nabla u^a(0) \cdot \nabla u^b(0) m_d(\omega),$$

and the series

$$\sum_{n=d}^\infty \xi_n \varepsilon^n$$

converges absolutely for  $\varepsilon \in ]-\varepsilon_\xi, \varepsilon_\xi[$ . (The symbol  $m_d(\dots)$  denotes the  $d$ -dimensional Lebesgue measure of a set).

**Proof.** We argue as in the proof of [3, Lem. 2.12]. We first note that the Theorem of change of variable in integrals implies that

$$\int_{\varepsilon\omega} \nabla u^a \cdot \nabla u^b \, dx = \varepsilon^d \int_\omega \nabla u^a(\varepsilon t) \cdot \nabla u^b(\varepsilon t) \, dt \quad \forall \varepsilon \in ]0, \varepsilon_0[.$$

Then analyticity of  $u^a$  and  $u^b$  (see Definition 1.5) and analyticity results for the composition operator (cf. Böhme and Tomi [6, p. 10], Henry [21, p. 29], Valent [41, Thm. 5.2, p. 44]), imply that there exists  $\varepsilon_\xi \in ]0, \varepsilon_0]$  such that the map from  $]-\varepsilon_\xi, \varepsilon_\xi[$  to  $C^{0,\alpha}(\bar{\omega})$  which takes  $\varepsilon$  to  $(\partial_j u^l)(\varepsilon \cdot)|_{\bar{\omega}}$  is real analytic for  $l = a, b$  and that

$$(\partial_j u^l)(\varepsilon t) = \sum_{h=0}^\infty \partial_j u_{\#,h+1}^l(t) \varepsilon^h \quad \forall t \in \bar{\omega}, \quad l = a, b,$$

where for  $l = a, b$  the series  $\sum_{h=0}^\infty \partial_j u_{\#,h+1}^l|_{\bar{\omega}} \varepsilon^h$  converges normally in  $C^{0,\alpha}(\bar{\omega})$  for  $\varepsilon \in ]-\varepsilon_\xi, \varepsilon_\xi[$ . Accordingly,

$$(\partial_j u^a)(\varepsilon t)(\partial_j u^b)(\varepsilon t) = \sum_{n=0}^\infty \left( \sum_{l=0}^n \partial_j u_{\#,l+1}^a(t) \partial_j u_{\#,n-l+1}^b(t) \right) \varepsilon^n \quad \forall t \in \bar{\omega}, \forall \varepsilon \in ]-\varepsilon_\xi, \varepsilon_\xi[ \setminus \{0\}.$$

Possibly taking a smaller  $\varepsilon_\xi$ , we also have

$$\int_\omega \nabla u^a(\varepsilon t) \cdot \nabla u^b(\varepsilon t) \, dt = \sum_{n=0}^\infty \left( \sum_{j=1}^d \sum_{l=0}^n \int_\omega \partial_j u_{\#,l+1}^a(t) \partial_j u_{\#,n-l+1}^b(t) \, dt \right) \varepsilon^n, \tag{28}$$

for all  $\varepsilon \in ]0, \varepsilon_\xi[$ . Moreover,

$$\sum_{j=1}^d \int_\omega \partial_j u_{\#,1}^a(t) \partial_j u_{\#,1}^b(t) \, dt = \nabla u^a(0) \cdot \nabla u^b(0) m_d(\omega).$$

Finally, to deduce the validity of the lemma, it is enough to multiply Eq. (28) by  $\varepsilon^d$ .  $\square$

To deduce our main result on the asymptotic behavior of  $\text{Cap}_\Omega(\varepsilon\bar{\omega}, u^a, u^b)$ , it suffices to integrate formula (27) over  $\partial\omega$  adding the coefficients of Lemma 5.1 and to apply Theorem 4.2.

**Theorem 5.2.** *With the notation introduced in Proposition 4.1, Theorem 4.2 and Lemma 5.1, let  $\{c_n\}_{n \in \mathbb{N}}$  be the sequence of real numbers defined by*

$$c_n \equiv 0 \quad \forall n \in \{0, \dots, d-3\}, \quad c_n \equiv - \int_{\partial\omega} \tilde{\lambda}_{n-(d-2)} \, d\sigma + \xi_n \quad \forall n \geq d-2.$$

Then there exists  $\varepsilon_c \in ]0, \varepsilon_0]$  such that

$$\text{Cap}_\Omega(\varepsilon\bar{\omega}, u^a, u^b) = \sum_{n=0}^\infty c_n \varepsilon^n$$

for all  $\varepsilon \in ]0, \varepsilon_c[$ . Moreover, the series

$$\sum_{n=0}^\infty c_n \varepsilon^n$$

converges absolutely for  $\varepsilon \in ]-\varepsilon_c, \varepsilon_c[$  and

$$c_{d-2} = - \int_{\partial\omega} \frac{u^a(0)u^b(0)}{r_0} \frac{\partial}{\partial\nu_\omega} v^-[\partial\omega, \rho_0^i] d\sigma = - \frac{u^a(0)u^b(0)}{r_0} \int_{\partial\omega} \rho_0^i d\sigma = - \frac{u^a(0)u^b(0)}{r_0}. \tag{29}$$

Our next aim is to better understand the value  $r_0$  which appears in formula (29) and, possibly, to link it to some boundary value problem related to the geometric setting. We begin with the lemma below where  $v^+[\partial\omega, \rho_0^i]$  is related to the solution of some exterior Dirichlet problem in  $\mathbb{R}^d \setminus \omega$ .

**Lemma 5.3.** *Let  $H^i \in C_{\text{loc}}^{1,\alpha}(\mathbb{R}^d \setminus \omega)$  be the solution of*

$$\begin{cases} \Delta H^i = 0 & \text{in } \mathbb{R}^d \setminus \bar{\omega}, \\ H^i(t) = 1 & \text{for all } t \in \partial\omega, \\ \lim_{t \rightarrow \infty} H^i(t) = 0. \end{cases} \tag{30}$$

Then the restriction  $v^+[\partial\omega, \rho_0^i]$  is constant and equal to  $((2-d)s_d \lim_{t \rightarrow \infty} |t|^{d-2} H^i(t))^{-1}$ .

**Proof.** Let  $u \in C_{\text{loc}}^{1,\alpha}(\mathbb{R}^d \setminus \omega)$ ,  $\Delta u = 0$  in  $\mathbb{R}^d \setminus \bar{\omega}$ , and  $\lim_{t \rightarrow \infty} u(t) = 0$ . Then by classical potential theory there exists  $\mu \in C^{1,\alpha}(\partial\omega)$  such that

$$u = w_\omega^-[\mu] + \frac{\int_{\partial\omega} u \rho_0^i d\sigma}{\int_{\partial\omega} \frac{1}{d\sigma} \int_{\partial\omega} v[\partial\omega, \rho_0^i] d\sigma} v^-[\partial\omega, \rho_0^i]$$

(cf., e.g., Folland [19, Chap. 3]). Then

$$\begin{aligned} \lim_{t \rightarrow \infty} |t|^{d-2} u(t) &= \lim_{t \rightarrow \infty} |t|^{d-2} w_\omega^-[\mu](t) + \lim_{t \rightarrow \infty} |t|^{d-2} \frac{\int_{\partial\omega} u \rho_0^i d\sigma}{\int_{\partial\omega} \frac{1}{d\sigma} \int_{\partial\omega} v[\partial\omega, \rho_0^i] d\sigma} v^-[\partial\omega, \rho_0^i](t) \\ &= \lim_{t \rightarrow \infty} \frac{\int_{\partial\omega} u \rho_0^i d\sigma}{\int_{\partial\omega} \frac{1}{d\sigma} \int_{\partial\omega} v[\partial\omega, \rho_0^i] d\sigma} |t|^{d-2} v^-[\partial\omega, \rho_0^i](t) \\ &= \frac{\int_{\partial\omega} u \rho_0^i d\sigma}{\int_{\partial\omega} \frac{1}{d\sigma} \int_{\partial\omega} v[\partial\omega, \rho_0^i] d\sigma} \frac{1}{(2-d)s_d} \int_{\partial\omega} \rho_0^i d\sigma \\ &= \frac{1}{(2-d)s_d} \frac{\int_{\partial\omega} u \rho_0^i d\sigma}{\int_{\partial\omega} \frac{1}{d\sigma} \int_{\partial\omega} v[\partial\omega, \rho_0^i] d\sigma}. \end{aligned}$$

As a consequence,

$$\begin{aligned} \lim_{t \rightarrow \infty} |t|^{d-2} H^i(t) &= \frac{1}{(2-d)s_d} \frac{\int_{\partial\omega} 1 \rho_0^i d\sigma}{\int_{\partial\omega} \frac{1}{d\sigma} \int_{\partial\omega} v[\partial\omega, \rho_0^i] d\sigma} \\ &= \frac{1}{(2-d)s_d} \frac{1}{\int_{\partial\omega} \frac{1}{d\sigma} \int_{\partial\omega} v[\partial\omega, \rho_0^i] d\sigma}, \end{aligned}$$

and thus

$$\frac{1}{\int_{\partial\omega} \frac{1}{d\sigma} \int_{\partial\omega} v[\partial\omega, \rho_0^i] d\sigma} = (2-d)s_d \lim_{t \rightarrow \infty} |t|^{d-2} H^i(t).$$

Moreover, by the jump properties of the single layer potential we have  $\nu_\omega \cdot \nabla v^+[\partial\omega, \rho_0^i]|_{\partial\omega} = 0$ . Thus  $v^+[\partial\omega, \rho_0^i]$  is constant in  $\bar{\omega}$  and the validity of the statement follows.  $\square$

**Corollary 5.4.** *Let  $g \in C^{1,\alpha}(\partial\omega)$ . Let  $u \in C_{\text{loc}}^{1,\alpha}(\mathbb{R}^d \setminus \omega)$  be the solution of*

$$\begin{cases} \Delta u = 0 & \text{in } \mathbb{R}^d \setminus \bar{\omega}, \\ u(t) = g(t) & \text{for all } t \in \partial\omega, \\ \lim_{t \rightarrow \infty} u(t) = 0. \end{cases}$$

Then

$$\lim_{t \rightarrow \infty} |t|^{d-2} u(t) = \frac{1}{(2-d)s_d} \frac{\int_{\partial\omega} u \rho_0^i d\sigma}{\int_{\partial\omega} \frac{1}{d\sigma} \int_{\partial\omega} v[\partial\omega, \rho_0^i] d\sigma} = \int_{\partial\omega} g \rho_0^i d\sigma \lim_{t \rightarrow \infty} |t|^{d-2} H^i(t).$$

**Remark 5.5.** Let  $H^i$  be the unique solution in  $C_{\text{loc}}^{1,\alpha}(\mathbb{R}^d \setminus \omega)$  of problem (30). Then by Lemma 5.3 we have

$$-\frac{1}{r_0} = (d-2)s_d \lim_{t \rightarrow \infty} |t|^{d-2} H^i(t).$$

Accordingly,

$$\text{Cap}_\Omega(\varepsilon\bar{\omega}, u^a, u^b) = u^a(0)u^b(0)(d-2)s_d \lim_{t \rightarrow \infty} |t|^{d-2} H^i(t)\varepsilon^{d-2} + \varepsilon^{d-1} \left( \sum_{n=d-1}^\infty c_n \varepsilon^{n-(d-1)} \right) \tag{31}$$

for all  $\varepsilon \in ]0, \varepsilon_c[$ .

We now wish to provide an alternative characterization of the quantity

$$(d-2)s_d \lim_{t \rightarrow \infty} |t|^{d-2} H^i(t).$$

By the Divergence theorem in exterior domains for functions which are harmonic at infinity (see Dalla Riva, Lanza de Cristoforis, and Musolino [11, §4.2]), we have

$$\begin{aligned} \int_{\mathbb{R}^d \setminus \bar{\omega}} |\nabla H^i(t)|^2 dt &= - \int_{\partial\omega} H^i(t) \frac{\partial}{\partial \nu_\omega} H^i(t) d\sigma_t \\ &= - \int_{\partial\omega} \frac{\partial}{\partial \nu_\omega} H^i(t) d\sigma_t. \end{aligned}$$

On the other hand, by Lemma 5.3 one verifies that

$$H^i = ((2-d)s_d \lim_{t \rightarrow \infty} |t|^{d-2} H^i(t)) v^-[\partial\omega, \rho_0^i] \quad \text{in } \mathbb{R}^d \setminus \omega,$$

and that

$$\begin{aligned} \frac{\partial}{\partial \nu_\omega} H^i &= ((2-d)s_d \lim_{t \rightarrow \infty} |t|^{d-2} H^i(t)) \left( \frac{1}{2} \rho_0^i + W_\omega^*[\rho_0^i] \right) \\ &= ((2-d)s_d \lim_{t \rightarrow \infty} |t|^{d-2} H^i(t)) \rho_0^i \quad \text{on } \partial\omega. \end{aligned} \tag{32}$$

As a consequence,

$$\begin{aligned} - \int_{\partial\omega} \frac{\partial}{\partial \nu_\omega} H^i(t) d\sigma_t &= ((d-2)s_d \lim_{t \rightarrow \infty} |t|^{d-2} H^i(t)) \int_{\partial\omega} \rho_0^i d\sigma \\ &= ((d-2)s_d \lim_{t \rightarrow \infty} |t|^{d-2} H^i(t)). \end{aligned}$$

Accordingly,

$$\int_{\mathbb{R}^d \setminus \bar{\omega}} |\nabla H^i(t)|^2 dt = (d - 2)s_d \lim_{t \rightarrow \infty} |t|^{d-2} H^i(t).$$

In other words, the quantity

$$(d - 2)s_d \lim_{t \rightarrow \infty} |t|^{d-2} H^i(t)$$

equals the energy integral of  $H^i$ , *i.e.*, of the unique function in  $C_{loc}^{1,\alpha}(\mathbb{R}^d \setminus \omega)$  which is harmonic in  $\mathbb{R}^d \setminus \bar{\omega}$  and at infinity and that is equal to 1 on  $\partial\omega$ . Such quantity is thus equal to the Newtonian capacity  $\text{Cap}_{\mathbb{R}^d}(\bar{\omega})$  (see (4)).

**Remark 5.6.** In case  $\omega$  is equal to the open unit ball  $B_1$  in  $\mathbb{R}^d$  of center 0 and radius 1, one verifies that

$$H^i(t) = |t|^{2-d} \quad \forall t \in \mathbb{R}^d \setminus B_1.$$

As a consequence,

$$(d - 2)s_d \lim_{t \rightarrow \infty} |t|^{d-2} H^i(t) = (d - 2)s_d,$$

and thus

$$\text{Cap}_{\mathbb{R}^d}(\bar{B}_1) = (d - 2)s_d.$$

In particular, if  $d = 3$ , then

$$\text{Cap}_{\mathbb{R}^3}(\bar{B}_1) = 4\pi.$$

### 6. Asymptotic behavior of $\text{Cap}_{\Omega}(\varepsilon\bar{\omega}, u^a, u^b)$ under vanishing assumption for $u^a$ and $u^b$

In this section, we investigate the behavior of  $\text{Cap}_{\Omega}(\varepsilon\bar{\omega}, u^a, u^b)$  assuming that the functions  $u^a$  and  $u^b$  and their derivatives up to a certain order could vanish at 0 and we modify the computation of [1, §5.1] and [3, §2.7]. So we consider the following assumption

$$\begin{aligned} u^a \text{ and } u^b \text{ are admissible functions as in Definition 1.5,} \\ \text{not identically zero in a neighborhood of 0.} \end{aligned} \tag{33}$$

Assumption (33) implies that  $u^a, u^b$  have finite order of vanishing  $\bar{k}^a, \bar{k}^b \in \mathbb{N}$  at 0, that is

$$D^\gamma u^l(0) = 0 \quad \forall |\gamma| < \bar{k}^l, \quad D^{\beta^l} u^l(0) \neq 0 \quad \text{for some } \beta^l \in \mathbb{N}^d \text{ with } |\beta^l| = \bar{k}^l, \quad l = a, b.$$

By Remark 5.5, we already know the principal term of the asymptotic expansion of  $\text{Cap}_{\Omega}(\varepsilon\bar{\omega}, u^a, u^b)$  as  $\varepsilon \rightarrow 0$  in case  $\bar{k}^a = \bar{k}^b = 0$  (*i.e.*,  $u^a(0)u^b(0) \neq 0$ ). We now wish to investigate the case when at least one between  $u^a$  and  $u^b$  vanishes at 0. The computation below notably simplifies if  $\bar{k}^a$  or  $\bar{k}^b$  is equal to 0, although it is not necessary to assume that this is the case. By condition (33) and Proposition 3.2 we note that

$$(\theta_k^a, \theta_k^b) = (0, 0) \quad \forall k < \bar{k}^a, \quad \theta_{\bar{k}^a}^a = 0, \tag{34}$$

and that  $\theta_{\bar{k}^a}^a$  is the unique solution in  $C^{1,\alpha}(\partial\omega)_0$  of

$$\frac{1}{2}\theta_{\bar{k}^a}^a(t) - W_{\omega}[\theta_{\bar{k}^a}^a](t) = \sum_{|\beta|=\bar{k}^a} \frac{\bar{k}^a!}{\beta!} t^\beta (D^\beta u^a)(0) - \sum_{\substack{\beta \in \mathbb{N}^d \\ |\beta|=\bar{k}^a}} \frac{\bar{k}^a!}{\beta!} \int_{\partial\omega} s^\beta (D^\beta u^a)(0) \rho_0^i(s) d\sigma_s \quad \forall t \in \partial\omega,$$

*i.e.*,

$$\frac{1}{2}\theta_{\bar{k}^a}^a(t) - W_{\omega}[\theta_{\bar{k}^a}^a](t) = \bar{k}^a! \left( u_{\#, \bar{k}^a}(t) - \int_{\partial\omega} u_{\#, \bar{k}^a} \rho_0^i d\sigma \right) \quad \forall t \in \partial\omega. \tag{35}$$

Then Eqs. (34), (35), and Proposition 4.1 imply that

$$u_{m,k}^a = 0 \quad \forall k < \bar{k}^a, \quad u_{m,\bar{k}^a}^a = -\frac{1}{\bar{k}^a} w^- [\partial\omega, \theta_{\bar{k}^a}^i]. \tag{36}$$

Hence, by the properties of the double layer potential, one verifies that  $u_{m,\bar{k}^a}^a$  is the unique solution in  $C_{loc}^{1,\alpha}(\mathbb{R}^d \setminus \omega)$  of

$$\begin{cases} \Delta u_{m,\bar{k}^a}^a = 0 & \text{in } \mathbb{R}^d \setminus \bar{\omega}, \\ u_{m,\bar{k}^a}^a(t) = u_{\#, \bar{k}^a}^a(t) - \int_{\partial\omega} u_{\#, \bar{k}^a}^a \rho_0^i d\sigma & \text{for all } t \in \partial\omega, \\ \lim_{t \rightarrow \infty} u_{m,\bar{k}^a}^a(t) = 0. \end{cases}$$

In addition, by assumption (33) and Proposition 4.1 we deduce that

$$g_k^a = 0 \quad \forall k < \bar{k}^a, \quad g_{\bar{k}^a}^a = \int_{\partial\omega} u_{\#, \bar{k}^a}^a \rho_0^i d\sigma. \tag{37}$$

Then by (33) and by Proposition 4.1 we show that

$$u_{\#,k}^l = 0 \quad \forall k < \bar{k}^l, \quad l = a, b. \tag{38}$$

As a consequence, Proposition 4.1 and Eqs. (36), (38) imply

$$\tilde{u}_k = 0 \quad \forall k < \bar{k}^a + \bar{k}^b, \quad \tilde{u}_{\bar{k}^a + \bar{k}^b} = \left( \frac{\partial u_{m,\bar{k}^a}^a}{\partial \nu_\omega} \right) u_{\#, \bar{k}^b}^b. \tag{39}$$

By (37) and (38) we have

$$\tilde{g}_k = 0 \quad \forall k < \bar{k}^a + \bar{k}^b, \quad \tilde{g}_{\bar{k}^a + \bar{k}^b} = g_{\bar{k}^a}^a u_{\#, \bar{k}^b}^b = \left( \int_{\partial\omega} u_{\#, \bar{k}^a}^a \rho_0^i d\sigma \right) u_{\#, \bar{k}^b}^b. \tag{40}$$

We now consider the quantities  $\tilde{a}_n, \tilde{\lambda}_n$  introduced in Theorem 4.2 for representing the behavior of  $\nu_\omega(\cdot) \cdot \nabla(u_\varepsilon^a(\varepsilon \cdot))u^b(\varepsilon \cdot)$ . By a direct computation based on (39), (40) we have

$$\tilde{a}_n = 0 \quad \forall n < \bar{k}^a + \bar{k}^b, \quad \tilde{a}_{\bar{k}^a + \bar{k}^b} = \tilde{g}_{\bar{k}^a + \bar{k}^b} \tilde{v}_0 = \tilde{v}_0 \left( \int_{\partial\omega} u_{\#, \bar{k}^a}^a \rho_0^i d\sigma \right) u_{\#, \bar{k}^b}^b,$$

and thus

$$\begin{aligned} \tilde{\lambda}_n &= 0 \quad \forall n < \bar{k}^a + \bar{k}^b, \\ \tilde{\lambda}_{\bar{k}^a + \bar{k}^b} &= \tilde{u}_{\bar{k}^a + \bar{k}^b} + \frac{\tilde{a}_{\bar{k}^a + \bar{k}^b}}{r_0} = \left( \frac{\partial u_{m,\bar{k}^a}^a}{\partial \nu_\omega} \right) u_{\#, \bar{k}^b}^b + \frac{1}{r_0} \tilde{v}_0 \left( \int_{\partial\omega} u_{\#, \bar{k}^a}^a \rho_0^i d\sigma \right) u_{\#, \bar{k}^b}^b. \end{aligned} \tag{41}$$

Moreover, a simple computation shows that

$$\xi_n = 0 \quad \forall n < \bar{k}^a + \bar{k}^b + d - 2, \quad \xi_{\bar{k}^a + \bar{k}^b + d - 2} = \int_{\omega} \nabla u_{\#, \bar{k}^a}^a \cdot \nabla u_{\#, \bar{k}^b}^b dt.$$

Finally, by Theorem 5.2 and by integrating equalities (41), we obtain

$$\begin{aligned} c_n &= 0 \quad \forall n < \bar{k}^a + \bar{k}^b + d - 2, \\ c_{\bar{k}^a + \bar{k}^b + d - 2} &= - \int_{\partial\omega} \left( \tilde{u}_{\bar{k}^a + \bar{k}^b} + \frac{\tilde{a}_{\bar{k}^a + \bar{k}^b}}{r_0} \right) d\sigma + \int_{\omega} \nabla u_{\#, \bar{k}^a}^a \cdot \nabla u_{\#, \bar{k}^b}^b dt \\ &= - \int_{\partial\omega} \left( \frac{\partial u_{m,\bar{k}^a}^a}{\partial \nu_\omega} \right) u_{\#, \bar{k}^b}^b d\sigma - \frac{1}{r_0} \int_{\partial\omega} \tilde{v}_0 u_{\#, \bar{k}^b}^b d\sigma \int_{\partial\omega} u_{\#, \bar{k}^a}^a \rho_0^i d\sigma + \int_{\omega} \nabla u_{\#, \bar{k}^a}^a \cdot \nabla u_{\#, \bar{k}^b}^b dt \\ &= - \int_{\partial\omega} \left( \frac{\partial u_{m,\bar{k}^a}^a}{\partial \nu_\omega} \right) u_{\#, \bar{k}^b}^b d\sigma - \frac{1}{r_0} \int_{\partial\omega} u_{\#, \bar{k}^a}^a \rho_0^i d\sigma \int_{\partial\omega} u_{\#, \bar{k}^b}^b \rho_0^i d\sigma + \int_{\omega} \nabla u_{\#, \bar{k}^a}^a \cdot \nabla u_{\#, \bar{k}^b}^b dt. \end{aligned}$$

Now let  $H_{u^a, \bar{k}^a} \in C_{loc}^{1,\alpha}(\mathbb{R}^d \setminus \omega)$  be the solution of

$$\begin{cases} \Delta H_{u^a, \bar{k}^a} = 0 & \text{in } \mathbb{R}^d \setminus \bar{\omega}, \\ H_{u^a, \bar{k}^a}(t) = \int_{\partial\omega} u_{\#, \bar{k}^a}^a \rho_0^i d\sigma & \text{for all } t \in \partial\omega, \\ \lim_{t \rightarrow \infty} H_{u^a, \bar{k}^a} = 0. \end{cases}$$

Then by Remark 5.5 we have

$$H_{u^a, \bar{k}^a} = \left( \int_{\partial\omega} u_{\#, \bar{k}^a}^a \rho_0^i d\sigma \right) H^i \quad \text{in } \mathbb{R}^d \setminus \omega,$$

and accordingly by (32)

$$\frac{\partial}{\partial\nu_\omega} H_{u^a, \bar{k}^a} = \frac{1}{r_0} \left( \int_{\partial\omega} u_{\#, \bar{k}^a}^a \rho_0^i d\sigma \right) \rho_0^i \quad \text{on } \partial\omega.$$

As a consequence,

$$\frac{1}{r_0} \int_{\partial\omega} u_{\#, \bar{k}^a}^a \rho_0^i d\sigma \int_{\partial\omega} u_{\#, \bar{k}^b}^b \rho_0^i d\sigma = \int_{\partial\omega} \left( \frac{\partial H_{u^a, \bar{k}^a}}{\partial\nu_\omega} \right) u_{\#, \bar{k}^b}^b|_{\partial\omega} d\sigma.$$

Then for  $l = a, b$  we denote by  $u_{\bar{k}^l}^l$  the unique solution in  $C_{loc}^{1,\alpha}(\mathbb{R}^d \setminus \omega)$  of

$$\begin{cases} \Delta u_{\bar{k}^l}^l = 0 & \text{in } \mathbb{R}^d \setminus \bar{\omega}, \\ u_{\bar{k}^l}^l(t) = u_{\#, \bar{k}^l}^l(t) & \text{for all } t \in \partial\omega, \\ \lim_{t \rightarrow \infty} u_{\bar{k}^l}^l(t) = 0. \end{cases} \tag{42}$$

Thus

$$u_{\bar{k}^a}^a = u_{m, \bar{k}^a}^a + H_{u^a, \bar{k}^a} \quad \text{in } \mathbb{R}^d \setminus \omega,$$

and therefore

$$- \int_{\partial\omega} \left( \frac{\partial u_{m, \bar{k}^a}^a}{\partial\nu_\omega} \right) u_{\#, \bar{k}^b}^b|_{\partial\omega} d\sigma - \frac{1}{r_0} \int_{\partial\omega} u_{\#, \bar{k}^a}^a \rho_0^i d\sigma \int_{\partial\omega} u_{\#, \bar{k}^b}^b \rho_0^i d\sigma = - \int_{\partial\omega} \left( \frac{\partial u_{\bar{k}^a}^a}{\partial\nu_\omega} \right) u_{\bar{k}^b}^b d\sigma.$$

On the other hand, the harmonicity at infinity of  $u_{\bar{k}^a}^a$  and of  $u_{\bar{k}^b}^b$  and the Divergence Theorem imply that

$$\int_{\mathbb{R}^d \setminus \bar{\omega}} \nabla u_{\bar{k}^a}^a \cdot \nabla u_{\bar{k}^b}^b dt = - \int_{\partial\omega} \left( \frac{\partial u_{\bar{k}^a}^a}{\partial\nu_\omega} \right) u_{\bar{k}^b}^b d\sigma$$

(cf. Folland [19, p. 118], Dalla Riva, Lanza de Cristoforis, and Musolino [11, Cor. 4.7]). Accordingly,

$$c_{\bar{k}^a + \bar{k}^b + d - 2} = \int_{\mathbb{R}^d \setminus \bar{\omega}} \nabla u_{\bar{k}^a}^a \cdot \nabla u_{\bar{k}^b}^b dt + \int_{\omega} \nabla u_{\#, \bar{k}^a}^a \cdot \nabla u_{\#, \bar{k}^b}^b dt.$$

Incidentally, we note that if for example  $u^b(0) \neq 0$  (i.e., if  $\bar{k}^b = 0$ ), then  $u_{\#, \bar{k}^b}^b = u^b(0)$  (and so  $\nabla u_{\#, \bar{k}^b}^b = 0$ ) and  $u_{\bar{k}^b}^b = u^b(0)H^i$ . Therefore, if  $\bar{k}^b = 0$  the term  $c_{\bar{k}^a + \bar{k}^b + d - 2}$  reduces to  $u^b(0) \int_{\mathbb{R}^d \setminus \bar{\omega}} \nabla u_{\bar{k}^a}^a \cdot \nabla H^i dt$ . Similarly, if  $\bar{k}^a = 0$  the term  $c_{\bar{k}^a + \bar{k}^b + d - 2}$  reduces to  $u^a(0) \int_{\mathbb{R}^d \setminus \bar{\omega}} \nabla u_{\bar{k}^b}^b \cdot \nabla H^i dt$ . We also note that if both  $\bar{k}^a = 0$  and  $\bar{k}^b = 0$ , then for  $l = a, b$  we have  $u_{\#, \bar{k}^l}^l = u^l(0)$  (and so  $\nabla u_{\#, \bar{k}^l}^l = 0$ ) and  $u_{\bar{k}^l}^l = u^l(0)H^i$ , and accordingly

$$\begin{aligned} \int_{\mathbb{R}^d \setminus \bar{\omega}} \nabla u_{\bar{k}^a}^a \cdot \nabla u_{\bar{k}^b}^b dt + \int_{\omega} \nabla u_{\#, \bar{k}^a}^a \cdot \nabla u_{\#, \bar{k}^b}^b dt &= u^a(0)u^b(0) \int_{\mathbb{R}^d \setminus \bar{\omega}} |\nabla H^i|^2 dt \\ &= u^a(0)u^b(0)(d-2)s_d \lim_{t \rightarrow \infty} |t|^{d-2} H^i(t). \end{aligned}$$

In other words, the quantity  $u^a(0)u^b(0)(d - 2)s_d \lim_{t \rightarrow \infty} |t|^{d-2} H^i(t)$  can be seen as the specific value of  $\int_{\mathbb{R}^d \setminus \bar{\omega}} \nabla u_{\bar{k}^a}^a \cdot \nabla u_{\bar{k}^b}^b dt + \int_{\omega} \nabla u_{\#, \bar{k}^a}^a \cdot \nabla u_{\#, \bar{k}^b}^b dt$  when both  $\bar{k}^a$  and  $\bar{k}^b$  are equal to 0.

As a consequence, under assumption (33), by Theorem 5.2 and formula (31), we can deduce the validity of the following.

**Theorem 6.1.** *Let assumption (33) hold. For  $l = a, b$ , let  $u_{\bar{k}^l}^l$  be the unique solution in  $C_{\text{loc}}^{1,\alpha}(\mathbb{R}^d \setminus \omega)$  of (42). Then*

$$\begin{aligned} & \text{Cap}_{\Omega}(\varepsilon \bar{\omega}, u^a, u^b) \\ &= \varepsilon^{\bar{k}^a + \bar{k}^b + d - 2} \left( \int_{\mathbb{R}^d \setminus \bar{\omega}} \nabla u_{\bar{k}^a}^a \cdot \nabla u_{\bar{k}^b}^b dt + \int_{\omega} \nabla u_{\#, \bar{k}^a}^a \cdot \nabla u_{\#, \bar{k}^b}^b dt \right) + \sum_{n=\bar{k}^a + \bar{k}^b + d - 1}^{\infty} \varepsilon^n c_n, \end{aligned} \tag{43}$$

for all  $\varepsilon \in ]0, \varepsilon_c[$ .

**Remark 6.2.** Under assumption (33), by (43) we have

$$\begin{aligned} & \text{Cap}_{\Omega}(\varepsilon \bar{\omega}, u^a, u^b) \\ &= \varepsilon^{\bar{k}^a + \bar{k}^b + d - 2} \left( \int_{\mathbb{R}^d \setminus \bar{\omega}} \nabla u_{\bar{k}^a}^a \cdot \nabla u_{\bar{k}^b}^b dt + \int_{\omega} \nabla u_{\#, \bar{k}^a}^a \cdot \nabla u_{\#, \bar{k}^b}^b dt \right) + o(\varepsilon^{\bar{k}^a + \bar{k}^b + d - 2}) \quad \text{as } \varepsilon \rightarrow 0. \end{aligned}$$

Moreover, we note that the coefficient of  $\varepsilon^{\bar{k}^a + \bar{k}^b + d - 2}$  depends both on the geometrical properties of the set  $\omega$  and on the behavior at 0 of the functions  $u^a$  and  $u^b$ , but does not depend on  $\Omega$ .

### 7. Asymptotic behavior of the eigenvalues of the Dirichlet-Laplacian in perforated domains

It is well known that if  $\Omega$  is a bounded open set in  $\mathbb{R}^d$ ,  $K$  a compact subset of  $\Omega$ , and if we denote by

$$0 < \lambda_1(\Omega) < \lambda_2(\Omega) \leq \dots \leq \lambda_N(\Omega) \leq \dots$$

and

$$0 < \lambda_1(\Omega \setminus K) < \lambda_2(\Omega \setminus K) \leq \dots \leq \lambda_N(\Omega \setminus K) \leq \dots$$

the sequences of the eigenvalues of the Dirichlet-Laplacian in  $\Omega$  and in  $\Omega \setminus K$ , respectively, then  $\lambda_N(\Omega \setminus K)$  is close to  $\lambda_N(\Omega)$  if and only if the capacity  $\text{Cap}_{\Omega}(K)$  of  $K$  in  $\Omega$  is small (see Rauch and Taylor [36]). A typical example is when we fix  $\Omega$  and  $\omega$  admissible domains, and we set

$$K = \varepsilon \bar{\omega} \quad \forall \varepsilon \in ] - \varepsilon_{\#}, \varepsilon_{\#}[ ,$$

with  $\varepsilon_{\#}$  as in (10). We define

$$\Omega_{\varepsilon} \equiv \Omega \setminus (\varepsilon \bar{\omega}) \quad \forall \varepsilon \in ] - \varepsilon_{\#}, \varepsilon_{\#}[ ,$$

as before, and we wish to study the convergence of the  $N$ th Dirichlet eigenvalues  $\lambda_N(\Omega_{\varepsilon}) = \lambda_N(\Omega \setminus (\varepsilon \bar{\omega}))$  to  $\lambda_N(\Omega)$  as  $\varepsilon \rightarrow 0$ .

In this section we show how the results on the asymptotic behavior of the generalizations of the capacity can be employed to obtain accurate asymptotic estimates for  $\lambda_N(\Omega_{\varepsilon})$  when  $\varepsilon \rightarrow 0$ , both in the case when  $\lambda_N(\Omega)$  is a simple eigenvalue of the Dirichlet-Laplacian in  $\Omega$  and in the case when it is a multiple eigenvalue, as we have done in [1,3] for the planar case. We treat these two situations in two different subsections.



### 7.1. Simple eigenvalues

We begin our analysis by considering the case when  $\lambda_N(\Omega)$  is a simple eigenvalue of the Dirichlet–Laplacian in the open set  $\Omega$

The following result by Courtois [10, Proof of Theorem 1.2] and Abatangelo, Felli, Hillairet, and Léna [2, Theorem 1.4] shows that the  $u$ -capacity can be successfully used to study the asymptotic behavior of the eigenvalues  $\lambda_N(\Omega \setminus (\varepsilon\bar{\omega}))$  as  $\varepsilon \rightarrow 0$ .

**Theorem 7.1.** *Let  $\lambda_N(\Omega)$  be a simple eigenvalue of the Dirichlet–Laplacian in a bounded, connected, and open set  $\Omega$ . Let  $u_N$  be a  $L^2(\Omega)$ -normalized eigenfunction associated to  $\lambda_N(\Omega)$  and let  $(K_\varepsilon)_{\varepsilon>0}$  be a family of compact sets contained in  $\Omega$  concentrating to a compact set  $K$  with  $\text{Cap}_\Omega(K) = 0$ . Then*

$$\lambda_N(\Omega \setminus K_\varepsilon) = \lambda_N(\Omega) + \text{Cap}_\Omega(K_\varepsilon, u_N) + o(\text{Cap}_\Omega(K_\varepsilon, u_N)), \quad \text{as } \varepsilon \rightarrow 0. \tag{44}$$

In view of Theorem 7.1, we can produce an asymptotic expansion of  $\lambda_N(\Omega \setminus (\varepsilon\bar{\omega}))$  by combining the expansion of  $\text{Cap}_\Omega(\varepsilon\bar{\omega}, u)$  and the asymptotic formula (44) for the eigenvalues.

Therefore, we fix  $\Omega$  and  $\omega$  admissible domains and we assume that

$$\begin{aligned} &\text{the } N\text{th eigenvalue } \lambda_N(\Omega) \text{ for the Dirichlet–Laplacian is simple} \\ &\text{and } u_N \text{ is a } L^2(\Omega)\text{-normalized eigenfunction related to } \lambda_N(\Omega). \end{aligned} \tag{45}$$

As a first step, we formulate our result on the asymptotic behavior of  $\text{Cap}_\Omega(\varepsilon\bar{\omega}, u_N)$ . Standard elliptic regularity theory (see for instance Friedman [20, Thm. 1.2, p. 205]) implies that  $u_N$  is analytic in a neighborhood of 0. Accordingly, by (31) we have

$$\begin{aligned} \text{Cap}_\Omega(\varepsilon\bar{\omega}, u_N) &= \text{Cap}_\Omega(\varepsilon\bar{\omega}, u_N, u_N) \\ &= (u_N(0))^2 \text{Cap}_{\mathbb{R}^d}(\bar{\omega}) \varepsilon^{d-2} + \varepsilon^{d-1} \left( \sum_{n=d-1}^\infty c_n \varepsilon^{n-(d-1)} \right) \end{aligned}$$

for  $\varepsilon$  positive and small, where the sequence of coefficients  $\{c_n\}_{n \in \mathbb{N}}$  is as in Theorem 5.2 and  $\text{Cap}_{\mathbb{R}^d}(\bar{\omega})$  is as in Eq. (4).

Then by formula (44) we immediately deduce the validity of the following result.

**Theorem 7.2.** *Let assumption (45) hold. Then*

$$\begin{aligned} &\lambda_N(\Omega \setminus (\varepsilon\bar{\omega})) \\ &= \lambda_N(\Omega) + (u_N(0))^2 \text{Cap}_{\mathbb{R}^d}(\bar{\omega}) \varepsilon^{d-2} + o(\varepsilon^{d-2}) \quad \text{as } \varepsilon \rightarrow 0^+. \end{aligned} \tag{46}$$

The above result agrees with the one of Maz’ya, Nazarov, and Plamenevskii [32] for  $N = 1$  and  $d = 3$ , with that of Ozawa [33] for  $d = 2, 3$ , and with that of Flucher [18] if  $d \geq 2$ .

If we assume that

$$u_N(0) = 0, \tag{47}$$

then formula (46) of Theorem 7.2 reduces to

$$\lambda_N(\Omega \setminus (\varepsilon\bar{\omega})) = \lambda_N(\Omega) + o(\varepsilon^{d-2}) \quad \text{as } \varepsilon \rightarrow 0^+.$$

Therefore, if (47) holds,

$$\begin{aligned} &\text{there exists } \bar{k} \in \mathbb{N} \setminus \{0\} \text{ such that } D^\gamma u_N(0) = 0 \quad \forall |\gamma| < \bar{k} \\ &\text{and that } D^\beta u_N(0) \neq 0 \text{ for some } \beta \in \mathbb{N}^d \text{ with } |\beta| = \bar{k}. \end{aligned} \tag{48}$$

Then we set

$$u_{N,\#\bar{k}}(t) \equiv \sum_{\substack{\beta \in \mathbb{N}^d \\ |\beta|=\bar{k}}} \frac{D^\beta u_N(0)}{\beta!} t^\beta \quad \forall t \in \mathbb{R}^d, \tag{49}$$

and we denote by  $u_{N,\bar{k}}$  the unique solution in  $C_{loc}^{1,\alpha}(\mathbb{R}^d \setminus \omega)$  of

$$\begin{cases} \Delta u_{N,\bar{k}} = 0 & \text{in } \mathbb{R}^d \setminus \bar{\omega}, \\ u_{N,\bar{k}}(t) = u_{N,\#\bar{k}}(t) & \text{for all } t \in \partial\omega, \\ \lim_{t \rightarrow \infty} u_{N,\bar{k}}(t) = 0. \end{cases} \tag{50}$$

Hence, Remark 6.2 implies that

$$\begin{aligned} \text{Cap}_\Omega(\varepsilon\bar{\omega}, u_N) &= \text{Cap}_\Omega(\varepsilon\bar{\omega}, u_N, u_N) \\ &= \varepsilon^{2\bar{k}+d-2} \left( \int_{\mathbb{R}^d \setminus \bar{\omega}} |\nabla u_{N,\bar{k}}|^2 dt + \int_\omega |\nabla u_{N,\#\bar{k}}|^2 dt \right) + o(\varepsilon^{2\bar{k}+d-2}) \\ &= \varepsilon^{2\bar{k}+d-2} \mathfrak{C}(\omega, u_{N,\#\bar{k}}) + o(\varepsilon^{2\bar{k}+d-2}) \quad \text{as } \varepsilon \rightarrow 0 \end{aligned}$$

(see (9)). Thus, by formula (44) of Theorem 7.1 we deduce the following result.

**Theorem 7.3.** *Let assumption (45) hold. Let  $\bar{k}$  be as in (48) and  $u_{N,\#\bar{k}}$  be as in (49). Let  $u_{N,\bar{k}}$  be the unique solution in  $C_{loc}^{1,\alpha}(\mathbb{R}^d \setminus \omega)$  of (50). Then*

$$\begin{aligned} \lambda_N(\Omega \setminus (\varepsilon\bar{\omega})) \\ = \lambda_N(\Omega) + \varepsilon^{2\bar{k}+d-2} \left( \int_{\mathbb{R}^d \setminus \bar{\omega}} |\nabla u_{N,\bar{k}}|^2 dt + \int_\omega |\nabla u_{N,\#\bar{k}}|^2 dt \right) + o(\varepsilon^{2\bar{k}+d-2}) \quad \text{as } \varepsilon \rightarrow 0. \end{aligned} \tag{51}$$

Clearly, Theorems 7.2 and 7.3 can be restated as Theorem 1.7 of the Introduction.

**Remark 7.4.** By the computation of Section 6, we note that

$$\begin{aligned} \int_{\mathbb{R}^d \setminus \bar{\omega}} |\nabla u_{N,0}|^2 dt + \int_\omega |\nabla u_{N,\#0}|^2 dt &= (u_N(0))^2 (d-2) s_d \lim_{t \rightarrow \infty} |t|^{d-2} H^i(t) \\ &= (u_N(0))^2 \text{Cap}_{\mathbb{R}^d}(\bar{\omega}), \end{aligned}$$

and thus Eq. (51) holds also when the order of vanishing  $\bar{k}$  of  $u_N$  is equal to 0.

### 7.2. Multiple eigenvalues

In the previous subsection, we have shown the asymptotic formula (51) for simple eigenvalues. In this subsection, instead, we follow the lines of the arguments of [3] and we consider the case where the  $N$ th eigenvalue  $\lambda_N(\Omega)$  for the Dirichlet–Laplacian is multiple.

So let  $\lambda_N(\Omega)$  be an eigenvalue of multiplicity  $m > 1$  of the Dirichlet–Laplacian in  $\Omega$  and let  $E(\lambda_N(\Omega))$  be the associated eigenspace. In particular, we have

$$\lambda_{N-1}(\Omega) < \lambda_N(\Omega) = \lambda_{N+j}(\Omega) < \lambda_{N+m}(\Omega) \quad \forall j = 0, \dots, m-1.$$

By [3, Appendix A], we have the following result on the decomposition of  $E(\lambda_N(\Omega))$ .

**Proposition 7.5.** *There exists a decomposition of  $E(\lambda_N(\Omega))$  into a sum of orthogonal subspaces*

$$E(\lambda_N(\Omega)) = E_1 \oplus \dots \oplus E_p$$

and an associated finite decreasing sequence of integers

$$k_1 > \dots > k_p \geq 0$$

such that, for all  $1 \leq j \leq p$ , a function in  $E_j \setminus \{0\}$  has the order of vanishing  $k_j$  at 0. In addition, such a decomposition is unique. We call it the order decomposition of  $E(\lambda_N(\Omega))$ .

We proceed as in [3] and we introduce some notation. So let  $u, v \in E(\lambda_N(\Omega)) \setminus \{0\}$ . Then  $u$  and  $v$  are admissible functions in the sense of Definition 1.5 and are not identically zero in a neighborhood of 0. Then  $u$  has a finite order of vanishing  $\kappa(u)$  at 0, with a corresponding principal part  $u_{\#}$ , meaning that

$$D^\gamma u(0) = 0 \quad \forall |\gamma| < \kappa(u)$$

and that  $D^\beta u(0) \neq 0$  for some  $\beta \in \mathbb{N}^d$  with  $|\beta| = \kappa(u)$ ,

and

$$u_{\#}(t) \equiv \sum_{\substack{\beta \in \mathbb{N}^d \\ |\beta| = \kappa(u)}} \frac{D^\beta u(0)}{\beta!} t^\beta \quad \forall t \in \mathbb{R}^d. \tag{52}$$

Consistent with Eq. (50), we denote by  $\mathbf{u}$  the unique solution in  $C_{\text{loc}}^{1,\alpha}(\mathbb{R}^d \setminus \omega)$  of

$$\begin{cases} \Delta \mathbf{u} = 0 & \text{in } \mathbb{R}^d \setminus \bar{\omega}, \\ \mathbf{u}(t) = u_{\#}(t) & \text{for all } t \in \partial\omega, \\ \lim_{t \rightarrow \infty} \mathbf{u}(t) = 0. \end{cases}$$

We write the order  $\kappa(v)$  and the functions  $v_{\#}, \mathbf{v}$  also for the function  $v$ . As in [3], we define

$$\mathcal{Q}(u, v) \equiv \int_{\mathbb{R}^d \setminus \bar{\omega}} \nabla \mathbf{u} \cdot \nabla \mathbf{v} \, dt + \int_{\omega} \nabla u_{\#} \cdot \nabla v_{\#} \, dt.$$

If instead  $u$  or  $v$  is identically zero, we set  $\mathcal{Q}(u, v) = 0$ . We observe that  $\mathcal{Q}$  is not a bilinear form, but the restriction of  $\mathcal{Q}$  to suitable subspaces of  $E(\lambda_N(\Omega))$  defines bilinear forms (cf. Definition 7.7 below).

By Remark 5.5 and Theorem 6.1, we can deduce the following result where we link the asymptotic behavior of  $\text{Cap}_{\Omega}(\varepsilon\bar{\omega}, u, v)$  with  $\mathcal{Q}(u, v)$ .

**Corollary 7.6.** *Let us fix  $u, v \in E(\lambda_N(\Omega)) \setminus \{0\}$ . Then,*

$$\text{Cap}_{\Omega}(\varepsilon\bar{\omega}, u, v) = \varepsilon^{\kappa(u)+\kappa(v)+d-2} \mathcal{Q}(u, v) + o\left(\varepsilon^{\kappa(u)+\kappa(v)+d-2}\right) \text{ as } \varepsilon \rightarrow 0^+.$$

We now consider again an eigenvalue  $\lambda_N(\Omega)$  of multiplicity  $m > 1$  and the associated eigenspace  $E(\lambda_N(\Omega))$  and we give the following definition (cf. Proposition 7.5).

**Definition 7.7.** For all  $1 \leq j \leq p$ , we define  $\mathcal{Q}_j$  on  $E_j$  by  $\mathcal{Q}_j(u, v) \equiv \mathcal{Q}(u, v)$ . It is a strictly positive (in particular non-degenerate) symmetric bilinear form on  $E_j$ .

We can now describe the behavior of the eigenvalues  $(\lambda_i(\Omega \setminus (\varepsilon\bar{\omega}))_{N \leq i \leq N+m-1})$ , and more specifically give the principal part of the spectral shift  $\lambda_i(\Omega \setminus (\varepsilon\bar{\omega})) - \lambda_N(\Omega)$  for each eigenvalue branch departing from  $\lambda_N(\Omega)$ .

**Theorem 7.8.** For  $1 \leq j \leq p$ , we write

$$m_j \equiv \dim(E_j),$$

so that

$$m = m_1 + \dots + m_j + \dots + m_p,$$

and we denote by

$$0 < \mu_{j,1} \leq \dots \leq \mu_{j,\ell} \leq \dots \leq \mu_{j,m_j}$$

the eigenvalues of the quadratic form  $\mathcal{Q}_j$ . Then, for all  $1 \leq j \leq p$  and  $1 \leq \ell \leq m_j$ ,

$$\lambda_{N-1+m_1+\dots+m_{j-1}+\ell}(\Omega \setminus (\varepsilon\bar{\omega})) = \lambda_N(\Omega) + \mu_{j,\ell} \varepsilon^{2k_j+(d-2)} + o(\varepsilon^{2k_j+(d-2)}) \text{ as } \varepsilon \rightarrow 0^+.$$

The proof of [Theorem 7.8](#) is identical with that of [[3](#), Th. 1.17]. To see how the above results imply [Theorem 1.8](#), we fix, for each  $1 \leq j \leq p$ , an orthonormal basis of  $E_j$ ,

$$v_{j,1}, \dots, v_{j,\ell}, \dots, v_{j,m_j}$$

such that, for all  $1 \leq \ell, \ell' \leq m_j$ ,

$$\mathcal{Q}_j(v_{j,\ell}, v_{j,\ell'}) = \delta_{\ell\ell'} \mu_{j,\ell}.$$

From [Theorem 7.8](#) and [Corollary 7.6](#), it then follows that, for all  $1 \leq j \leq p$  and  $1 \leq \ell \leq m_j$ ,

$$\lambda_{N-1+m_1+\dots+m_{j-1}+\ell}(\Omega \setminus (\varepsilon\bar{\omega})) = \lambda_N(\Omega) + \text{Cap}_\Omega(\varepsilon\bar{\omega}, v_{j,\ell}) + o(\text{Cap}_\Omega(\varepsilon\bar{\omega}, v_{j,\ell})) \text{ as } \varepsilon \rightarrow 0^+.$$

By relabeling the  $v_{j,\ell}$  with  $1 \leq i \leq m$ , in increasing order, first of  $j$ , then of  $\ell$ , we obtain an orthonormal basis of  $E(\lambda_N(\Omega))$ ,

$$v_1, \dots, v_i, \dots, v_m,$$

which has the properties of [Theorem 1.8](#).

Then, to illustrate our result, as in [[3](#), Cor. 1.18], we deduce the following corollary in a specific situation

**Corollary 7.9.** Let us assume that  $\lambda_N(\Omega)$  has multiplicity 2 (i.e.,  $m = 2$ ). Then one of the following alternatives holds.

- (1) There exist two normalized eigenfunctions  $u_1, u_2 \in E(\lambda_N(\Omega)) \setminus \{0\}$ , with respective order of vanishing  $k_1, k_2$  such that  $k_1 > k_2$ . In that case,

$$\begin{aligned} \lambda_N(\Omega \setminus (\varepsilon\bar{\omega})) &= \lambda_N(\Omega) + \mathcal{Q}(u_1, u_1) \varepsilon^{2k_1+(d-2)} + o(\varepsilon^{2k_1+(d-2)}); \\ \lambda_{N+1}(\Omega \setminus (\varepsilon\bar{\omega})) &= \lambda_N(\Omega) + \mathcal{Q}(u_2, u_2) \varepsilon^{2k_2+(d-2)} + o(\varepsilon^{2k_2+(d-2)}). \end{aligned}$$

- (2) All eigenfunctions in  $E(\lambda_N(\Omega))$  have the same order of vanishing, which we denote by  $k$ . Let us note that necessarily  $k \geq 1$ . In that case, let us choose eigenfunctions  $u_1, u_2$  forming an orthonormal basis of  $E(\lambda_N(\Omega))$  and let us denote by  $0 < \mu_1 \leq \mu_2$  the eigenvalues of the symmetric and positive definite matrix

$$\begin{pmatrix} \mathcal{Q}(u_1, u_1) & \mathcal{Q}(u_1, u_2) \\ \mathcal{Q}(u_1, u_2) & \mathcal{Q}(u_2, u_2) \end{pmatrix}.$$

Then

$$\begin{aligned} \lambda_N(\Omega \setminus (\varepsilon\bar{\omega})) &= \lambda_N(\Omega) + \mu_1 \varepsilon^{2k+(d-2)} + o(\varepsilon^{2k+(d-2)}); \\ \lambda_{N+1}(\Omega \setminus (\varepsilon\bar{\omega})) &= \lambda_N(\Omega) + \mu_2 \varepsilon^{2k+(d-2)} + o(\varepsilon^{2k+(d-2)}). \end{aligned}$$

### Acknowledgments

L.A. has been supported by MUR grant Dipartimento di Eccellenza 2023–2027. C.L. and P.M. are members of the “Gruppo Nazionale per l’Analisi Matematica, la Probabilità e le loro Applicazioni” (GNAMPA) of the “Istituto Nazionale di Alta Matematica” (INdAM) and acknowledge the support of the “INdAM GNAMPA Project” codice CUP\_E53C22001930001 “Operatori differenziali e integrali in geometria spettrale”. P.M. also acknowledges the support from EU through the H2020-MSCA-RISE-2020 project EffectFact, Grant agreement ID: 101008140, and the support of the SPIN Project “DOMain perturbation problems and Interactions Of scales - DOMINO” of the Ca’ Foscari University of Venice.

### Appendix. Blow-up analysis for the $u$ -capacity

Following the outline proposed in [17], we perform a blow up analysis for  $\text{Cap}_\Omega(\varepsilon\bar{\omega}, u)$ , with  $\Omega, \omega$  admissible domains and  $u$  an admissible function. Without loss of generality, we assume that  $u$  is not identically zero in a neighborhood of 0. In the final remark of this appendix we connect this with the results established in Section 6.

Firstly we introduce the following notation. Let  $K \subseteq \mathbb{R}^d$  be a compact set. For  $d \geq 3$ , the Beppo-Levi spaces  $\mathcal{D}^{1,2}(\mathbb{R}^d)$  and  $\mathcal{D}^{1,2}(\mathbb{R}^d \setminus K)$  are defined as the completion, with respect to the  $L^2$ -norm of the gradient, of  $C_c^\infty(\mathbb{R}^d)$  and  $C_c^\infty(\mathbb{R}^d \setminus K)$ , respectively. We recall as well that thanks to the well-known Hardy Inequality,

$$\left(\frac{d-2}{2}\right)^2 \int_{\mathbb{R}^d} \frac{v^2}{|x|^2} \leq \int_{\mathbb{R}^d} |\nabla v|^2 \quad (\text{Hardy Inequality}), \tag{53}$$

for all  $v \in C_c^\infty(\mathbb{R}^d)$ , the space  $\mathcal{D}^{1,2}(\mathbb{R}^d)$  can be characterized in the following way

$$\mathcal{D}^{1,2}(\mathbb{R}^d) = \left\{ v \in L^1_{\text{loc}}(\mathbb{R}^d) : \nabla v \in L^2(\mathbb{R}^d) \text{ and } \frac{v}{|x|} \in L^2(\mathbb{R}^d) \right\}. \tag{54}$$

For any  $f \in H^1_{\text{loc}}(\mathbb{R}^d)$  we introduce the quantity

$$\text{Cap}_{\mathbb{R}^d}(K, f) := \inf \left\{ \int_{\mathbb{R}^d} |\nabla v|^2 : v \in \mathcal{D}^{1,2}(\mathbb{R}^d), v - \eta_K f \in \mathcal{D}^{1,2}(\mathbb{R}^d \setminus K) \right\},$$

where  $\eta_K \in C_c^\infty(\mathbb{R}^d)$  such that  $\eta_K = 1$  in a neighborhood of  $K$ . In particular, when  $f = 1$  we have

$$\text{Cap}_{\mathbb{R}^d}(K, 1) = \text{Cap}_{\mathbb{R}^d}(K)$$

(for the definition of  $\text{Cap}_{\mathbb{R}^d}(K)$  see (4)). Let now  $u$  be a fixed admissible function, in the sense of Definition 1.5, not identically zero in a neighborhood of 0. We denote by  $\bar{k}$  its order of vanishing  $\kappa(u)$  at 0 and we define

$$\tilde{u}_\varepsilon(t) \equiv \begin{cases} \frac{u(\varepsilon t)}{\varepsilon^{\bar{k}}} & \forall t \in \frac{1}{\varepsilon}\Omega, \\ 0 & \forall t \in \mathbb{R}^d \setminus \frac{1}{\varepsilon}\Omega. \end{cases}$$

Since  $u$  is analytic in a neighborhood of 0,  $\tilde{u}_\varepsilon$  converges to  $u_\#$ , the principal part of  $u$  defined by Eq. (52), uniformly in every compact subset of  $\mathbb{R}^d$ , and similarly for the derivatives of  $\tilde{u}_\varepsilon$  to any order. This implies in turn that for any  $R > 0$ , as  $\varepsilon \rightarrow 0$ ,

$$\frac{1}{\varepsilon^{d+2\bar{k}}} \int_{\Omega \cap B_{R\varepsilon}} u^2 \rightarrow \int_{B_R} u_\#^2, \tag{55}$$

$$\frac{1}{\varepsilon^{d-2+2\bar{k}}} \int_{\Omega \cap B_{R\varepsilon}} |\nabla u|^2 \rightarrow \int_{B_R} |\nabla u_\#|^2. \tag{56}$$

Here, if  $\tilde{R}$  is a positive real number, the symbol  $B_{\tilde{R}}$  denotes the open ball in  $\mathbb{R}^d$  of radius  $\tilde{R}$  and center 0. Moreover, if  $K$  is a compact set with  $\text{Cap}_{\mathbb{R}^d}(K) > 0$ , we have that, for any  $R > 0$  such that  $K \subset B_R$ , there exists  $C_P > 0$  such that, for any  $v$  in the closure of  $C_c^\infty(\overline{B_R} \setminus K)$  in  $H^1(B_R)$ ,

$$\int_{B_R} v^2 \leq C_P \int_{B_R} |\nabla v|^2 \quad (\text{Poincaré Inequality}). \tag{57}$$

For a proof of the basic inequalities (53) and (57) in a similar context we refer the reader to [17, Lemma 6.5] and [17, Lemma 6.7], respectively.

In the following lemma, we deduce the vanishing rate of  $\text{Cap}_\Omega(\varepsilon\bar{\omega}, u)$  as  $\varepsilon \rightarrow 0$ .

**Lemma A.1.** For  $\varepsilon \rightarrow 0$ ,

$$\text{Cap}_\Omega(\varepsilon\bar{\omega}, u) = O(\varepsilon^{d-2+2\bar{k}}).$$

**Proof.** If  $\varepsilon$  is sufficiently small there exists  $R > 0$  such that  $\varepsilon\bar{\omega} \subset B_{R\varepsilon} \subset B_{2R\varepsilon} \subset \Omega$ . Let  $\eta_\varepsilon \in C_c^\infty(\Omega)$  such that  $0 \leq \eta_\varepsilon \leq 1$ ,  $\eta_\varepsilon \equiv 1$  in  $B_{R\varepsilon}$ ,  $\eta_\varepsilon \equiv 0$  in  $\Omega \setminus B_{2R\varepsilon}$  and  $|\nabla\eta_\varepsilon| \leq \frac{4}{\varepsilon R}$  in  $\Omega$ . Then  $\eta_\varepsilon u \in H_0^1(\Omega)$  and  $\eta_\varepsilon u - u \in H_0^1(\Omega \setminus \varepsilon\bar{\omega})$ . Therefore

$$\text{Cap}_\Omega(\varepsilon\bar{\omega}, u) \leq \int_\Omega |\nabla(\eta_\varepsilon u)|^2 = \int_\Omega |\eta_\varepsilon \nabla u + u \nabla \eta_\varepsilon|^2 \leq 2 \int_\Omega \eta_\varepsilon^2 |\nabla u|^2 + 2 \int_\Omega u^2 |\nabla \eta_\varepsilon|^2 = O(\varepsilon^{d-2+2\bar{k}}),$$

thanks to (55) and (56) together with the properties of  $\eta_\varepsilon$ .  $\square$

We now define a suitable rescaling of the  $u$ -capacitary potential, that is

$$\tilde{V}_\varepsilon(t) \equiv \begin{cases} \frac{V_{\varepsilon\bar{\omega},u}(\varepsilon t)}{\varepsilon^{\bar{k}}} & \forall t \in \frac{1}{\varepsilon}\Omega, \\ 0 & \forall t \in \mathbb{R}^d \setminus \frac{1}{\varepsilon}\Omega, \end{cases}$$

for any positive  $\varepsilon$ . Let us note that  $\tilde{V}_\varepsilon \in H_{\text{loc}}^1(\mathbb{R}^d)$ .

**Lemma A.2.** Let  $R > 0$ . There exists  $C > 0$  such that

$$\|\tilde{V}_\varepsilon\|_{H^1(B_R)} \leq C$$

if  $\varepsilon$  is small enough.

**Proof.** Let  $R_0 > 0$  such that  $B_{R_0} \subset \Omega$ . We have

$$\int_\Omega |\nabla V_{\varepsilon\bar{\omega},u}|^2 \geq \int_{B_{R_0}} |\nabla V_{\varepsilon\bar{\omega},u}|^2 = \varepsilon^{d-2+2\bar{k}} \int_{B_{\frac{R_0}{\varepsilon}}} \frac{|\nabla(V_{\varepsilon\bar{\omega},u}(\varepsilon t))|^2}{\varepsilon^{2\bar{k}}} = \varepsilon^{d-2+2\bar{k}} \int_{B_{\frac{R_0}{\varepsilon}}} |\nabla \tilde{V}_\varepsilon|^2.$$

From Lemma A.1 and the previous inequality, we deduce that

$$\int_{B_{\frac{R_0}{\varepsilon}}} |\nabla \tilde{V}_\varepsilon|^2 = O(1)$$

as  $\varepsilon \rightarrow 0$  and that there exists  $C_1 > 0$  such that for any  $R > 0$

$$\int_{B_R} |\nabla \tilde{V}_\varepsilon|^2 \leq C_1 \quad \text{for any } \varepsilon \in \left]0, \frac{R_0}{R}\right[. \tag{58}$$

On the other hand,

$$\begin{aligned} \int_{B_R} |\tilde{V}_\varepsilon|^2 &= \int_{B_R} |\tilde{V}_\varepsilon - \tilde{u}_\varepsilon + \tilde{u}_\varepsilon|^2 \leq 2 \int_{B_R} |\tilde{V}_\varepsilon - \tilde{u}_\varepsilon|^2 + 2 \int_{B_R} |\tilde{u}_\varepsilon|^2 \\ &\leq 2C_P \int_{B_R} |\nabla(\tilde{V}_\varepsilon - \tilde{u}_\varepsilon)|^2 + 2 \int_{B_R} |\tilde{u}_\varepsilon|^2 \leq 4C_P \left( \int_{B_R} |\nabla \tilde{V}_\varepsilon|^2 + \int_{B_R} |\nabla \tilde{u}_\varepsilon|^2 \right) + 2 \int_{B_R} |\tilde{u}_\varepsilon|^2 \end{aligned}$$

thanks to (57) applied on  $\tilde{V}_\varepsilon - \tilde{u}_\varepsilon$ . Every term in the last sum can be proved to be bounded following the preceding argument and taking into account (55) and (56), so that there exists  $\bar{\varepsilon} \in ]0, 1[$  and  $C_2 > 0$  such that

$$\int_{B_R} |\tilde{V}_\varepsilon|^2 \leq C_2 \quad \text{for any } \varepsilon \in ]0, \min\left\{\bar{\varepsilon}, \frac{R_0}{R}\right\}[. \tag{59}$$

Eqs. (58) and (59) conclude the proof.  $\square$

From this last lemma and a diagonal process over a sequence of  $R_n \rightarrow +\infty$  we deduce that there exist  $W \in H^1_{loc}(\mathbb{R}^d)$  and a subsequence (still denoted with  $\varepsilon$ ) such that

$$\tilde{V}_\varepsilon \rightharpoonup W \quad \text{as } \varepsilon \rightarrow 0$$

weakly in  $H^1(B_R)$ , strongly in  $L^2(B_R)$  for any  $R > 0$  and almost everywhere in  $\mathbb{R}^d$ .

By a change of variables we have

$$\int_{\mathbb{R}^d} |\nabla \tilde{V}_\varepsilon|^2 = \frac{1}{\varepsilon^{d-2+2k}} \int_{\Omega} |\nabla V_{\varepsilon\bar{\omega},u}|^2 = \frac{1}{\varepsilon^{d-2+2k}} \text{Cap}_\Omega(\varepsilon\bar{\omega}, u) = O(1) \tag{60}$$

as  $\varepsilon \rightarrow 0$  thanks to Lemma A.1. Via Hardy Inequality (53) we obtain also

$$\int_{\mathbb{R}^d} \frac{|\tilde{V}_\varepsilon|^2}{|x|^2} \leq \left(\frac{2}{d-2}\right)^2 \int_{\mathbb{R}^d} |\nabla \tilde{V}_\varepsilon|^2 \leq C$$

uniformly with respect to  $\varepsilon$ . Via Fatou’s lemma, we have

$$\int_{\mathbb{R}^d} \frac{|W|^2}{|x|^2} = \int_{\mathbb{R}^d} \liminf_{\varepsilon \rightarrow 0} \frac{|\tilde{V}_\varepsilon|^2}{|x|^2} \leq \liminf_{\varepsilon \rightarrow 0} \int_{\mathbb{R}^d} \frac{|\tilde{V}_\varepsilon|^2}{|x|^2} \leq \bar{C} \tag{61}$$

uniformly with respect to  $\varepsilon$  small enough. Moreover, for any  $R > 0$  we have

$$\int_{B_R} |\nabla W|^2 \leq \liminf_{\varepsilon \rightarrow 0} \int_{B_R} |\nabla \tilde{V}_\varepsilon|^2 \leq \bar{C}$$

where the last inequality follows from (60) (the constant  $\bar{C}$  does not depend on  $R$ ). We then deduce that

$$\int_{\mathbb{R}^d} |\nabla W|^2 \leq \bar{C}. \tag{62}$$

From (61) and (62) we deduce that  $W \in \mathcal{D}^{1,2}(\mathbb{R}^d)$  using the characterization (54).

On the other hand,  $V_{\varepsilon\bar{\omega},u}$  weakly solves the problem

$$\begin{cases} -\Delta V_{\varepsilon\bar{\omega},u} = 0 & \text{in } \Omega \setminus \varepsilon\bar{\omega}, \\ V_{\varepsilon\bar{\omega},u} = u & \text{on } \varepsilon\bar{\omega}, \\ V_{\varepsilon\bar{\omega},u} = 0 & \text{on } \partial\Omega, \end{cases} \tag{63}$$

and in particular

$$\int_{\frac{1}{\varepsilon}\Omega \setminus \bar{\omega}} \nabla \tilde{V}_\varepsilon \cdot \nabla \varphi = 0 \quad \text{for all } \varphi \in H^1_0\left(\frac{1}{\varepsilon}\Omega \setminus \bar{\omega}\right).$$

Taking the limit for  $\varepsilon \rightarrow 0$  in the previous equation we obtain that for any  $R > 0$

$$\int_{B_R \setminus \bar{\omega}} \nabla W \cdot \nabla \varphi = 0 \quad \text{for all } \varphi \in H_0^1(B_R \setminus \bar{\omega})$$

and therefore

$$\int_{\mathbb{R}^d \setminus \bar{\omega}} \nabla W \cdot \nabla \varphi = 0 \quad \text{for all } \varphi \in C_c^\infty(\mathbb{R}^d \setminus \bar{\omega}).$$

As a consequence, thanks also to the uniform convergence  $\tilde{u}_\varepsilon \rightarrow u_\#$  on compact sets,  $W$  is a weak solution of the problem

$$\begin{cases} -\Delta W = 0 & \text{in } \mathbb{R}^d \setminus \bar{\omega} \\ W = u_\# & \text{on } \bar{\omega} \\ W \in \mathcal{D}^{1,2}(\mathbb{R}^d) \end{cases}$$

By standard variational methods, the problem above for  $W$  has a unique weak solution which we denote by  $V_{\mathbb{R}^d, u_\#}$ , i.e.,  $W = V_{\mathbb{R}^d, u_\#}$ .

Let  $\eta$  be a cut-off function with compact support such that  $\eta \equiv 1$  on  $\bar{\omega}$ . Testing the previous equation with  $W - \eta u_\# \in \mathcal{D}^{1,2}(\mathbb{R}^d \setminus \bar{\omega})$  we obtain

$$\int_{\mathbb{R}^d \setminus \bar{\omega}} |\nabla W|^2 = \int_{\mathbb{R}^d \setminus \bar{\omega}} \nabla W \cdot \nabla(\eta u_\#). \tag{64}$$

Letting  $\eta_\varepsilon(x) \equiv \eta(\frac{x}{\varepsilon})$ , we can test Eq. (63) with  $\varphi = V_{\varepsilon\bar{\omega}, u} - \eta_\varepsilon u$ , obtaining

$$\int_{\Omega \setminus \varepsilon\bar{\omega}} |\nabla V_{\varepsilon\bar{\omega}, u}|^2 = \int_{\Omega \setminus \varepsilon\bar{\omega}} \nabla V_{\varepsilon\bar{\omega}, u} \cdot \nabla(\eta_\varepsilon u).$$

By a change of variables we obtain

$$\frac{1}{\varepsilon^{d-2+2\bar{k}}} \int_{\Omega \setminus \varepsilon\bar{\omega}} |\nabla V_{\varepsilon\bar{\omega}, u}|^2 = \int_{\frac{1}{\varepsilon}\Omega \setminus \bar{\omega}} \nabla \tilde{V}_\varepsilon \cdot \nabla(\eta \tilde{u}_\varepsilon) \rightarrow \int_{\mathbb{R}^d \setminus \bar{\omega}} \nabla W \cdot \nabla(\eta u_\#) \tag{65}$$

as  $\varepsilon \rightarrow 0$  thanks to the weak convergence  $\tilde{V}_\varepsilon \rightharpoonup W$  in  $H^1(B_R)$  for any  $R > 0$  and the uniform convergence  $\tilde{u}_\varepsilon \rightarrow u_\#$  on compact sets. Moreover, the very same convergence implies

$$\frac{1}{\varepsilon^{d-2+2\bar{k}}} \int_{\varepsilon\bar{\omega}} |\nabla u|^2 = \int_{\bar{\omega}} |\nabla \tilde{u}_\varepsilon|^2 \rightarrow \int_{\bar{\omega}} |\nabla u_\#|^2 \quad \text{as } \varepsilon \rightarrow 0. \tag{66}$$

In view of (66), (64) and (65) we deduce that

$$\varepsilon^{-(d-2+2\bar{k})} \text{Cap}_\Omega(\varepsilon\bar{\omega}, u) \rightarrow \text{Cap}_{\mathbb{R}^d}(\bar{\omega}, u_\#) \quad \text{as } \varepsilon \rightarrow 0, \tag{67}$$

independently from the subsequence.

Let us note that, when  $u$  is the eigenfunction  $u_N$ ,  $V_{\mathbb{R}^d, u_{N, \#, \bar{k}}|_{\bar{\omega}}} = u_{N, \#, \bar{k}}|_{\bar{\omega}}$  and  $V_{\mathbb{R}^d, u_{N, \#, \bar{k}}|_{\mathbb{R}^d \setminus \omega}} = u_{N, \bar{k}}$ . Eq. (67) can therefore be written

$$\text{Cap}_\Omega(\varepsilon\bar{\omega}, u_N) = \text{Cap}_{\mathbb{R}^d}(\bar{\omega}, u_{N, \#, \bar{k}}) \varepsilon^{2\bar{k}+d-2} + o\left(\varepsilon^{2\bar{k}+d-2}\right) \quad \text{as } \varepsilon \rightarrow 0.$$

**Remark A.3.** We recall that in our context  $\omega$  is an open regular bounded set in  $\mathbb{R}^d$  and  $u_\#$  is a harmonic homogeneous polynomial of degree  $\bar{k}$ . As a consequence, the quantity  $\text{Cap}_{\mathbb{R}^d}(\bar{\omega}, u_\#)$  is strictly positive and (67) is in fact a sharp asymptotics.

Moreover, from Remark 6.2 we deduce that

$$\left( \int_{\mathbb{R}^d \setminus \bar{\omega}} |\nabla u_{N, \bar{k}}|^2 dt + \int_{\omega} |\nabla u_{N, \#, \bar{k}}|^2 dt \right) = \text{Cap}_{\mathbb{R}^d}(\bar{\omega}, u_{N, \#, \bar{k}}) > 0.$$



**Remark A.4.** The blow-up analysis presented above can be also performed for the  $(u, v)$ -capacity provided suitable modifications.

## References

- [1] L. Abatangelo, V. Bonnaillie-Noël, C. Léna, P. Musolino, Asymptotic behavior of  $u$ -capacities and singular perturbations for the Dirichlet-Laplacian, *ESAIM Control Optim. Calc. Var.* 27 (2021) 43, suppl., Paper No. S25.
- [2] L. Abatangelo, V. Felli, L. Hillairet, C. Léna, Spectral stability under removal of small capacity sets and applications to Aharonov-Bohm operators, *J. Spectr. Theory* 9 (2) (2019) 379–427.
- [3] L. Abatangelo, C. Léna, P. Musolino, Ramification of multiple eigenvalues for the Dirichlet-Laplacian in perforated domains, *J. Funct. Anal.* 283 (12) (2022) 50, Paper (109718).
- [4] H. Ammari, H. Kang, H. Lee, Layer Potential Techniques in Spectral Analysis, in: *Mathematical Surveys and Monographs*, vol. 153, American Mathematical Society, Providence, RI, 2009.
- [5] G. Besson, Comportement asymptotique des valeurs propres du laplacien dans un domaine avec un trou. (French) [Asymptotic behavior of the eigenvalues of the Laplacian in a domain with a hole], *Bull. Soc. Math. France* 113 (2) (1985) 211–230.
- [6] R. Böhme, F. Tomi, Zur struktur der Lösungsmenge des plateauproblems, *Math. Z.* 133 (1973) 1–29.
- [7] I. Chavel, E.A. Feldman, Spectra of manifolds less a small domain, *Duke Math. J.* 56 (2) (1988) 399–414.
- [8] B. Colbois, G. Courtois, Convergence de variétés and convergence du spectre du laplacien. (French) [Convergence of manifolds and convergence of the spectrum of the Laplacian], *Ann. Sci. Éc. Norm. Supér.* 24 (4) (1991) 507–518.
- [9] M. Costabel, M. Dalla Riva, M. Dauge, P. Musolino, Converging expansions for Lipschitz self-similar perforations of a plane sector, *Integral Equations Operator Theory* 88 (3) (2017) 401–449.
- [10] G. Courtois, Spectrum of manifolds with holes, *J. Funct. Anal.* 134 (1995) 194–221.
- [11] M. Dalla Riva, M. Lanza de Cristoforis, P. Musolino, *Singularly Perturbed Boundary Value Problems. A Functional Analytic Approach*, Springer, Cham, 2021.
- [12] M. Dalla Riva, P. Luzzini, P. Musolino, Shape analyticity and singular perturbations for layer potential operators, *ESAIM Math. Model. Numer. Anal.* 56 (6) (2022) 1889–1910.
- [13] M. Dalla Riva, P. Musolino, Real analytic families of harmonic functions in a planar domain with a small hole, *J. Math. Anal. Appl.* 422 (2015) 37–55.
- [14] M. Dalla Riva, P. Musolino, S.V. Rogosin, Series expansions for the solution of the Dirichlet problem in a planar domain with a small hole, *Asymptot. Anal.* 92 (2015) 339–361.
- [15] K. Deimling, *Nonlinear Functional Analysis*, Springer-Verlag, Berlin, 1985.
- [16] V.N. Dubinin, Asymptotic behavior of the capacity of a condenser as some of its plates contract to points, *Transl. Mat. Zametki* 96 (2) (2014) 194–206, *Math. Notes* 96 (2014) no. 1-2, 187–198.
- [17] V. Felli, B. Noris, R. Ognibene, Eigenvalues of the Laplacian with moving mixed boundary conditions: the case of disappearing Dirichlet region, *Calc. Var. Partial Differential Equations* 60 (1) (2021) 12.
- [18] M. Flucher, Approximation of Dirichlet eigenvalues on domains with small holes, *J. Math. Anal. Appl.* 193 (1) (1995) 169–199.
- [19] G.B. Folland, *Introduction to Partial Differential Equations*, Princeton University Press, Princeton NJ, 1995.
- [20] A. Friedman, *Partial Differential Equations*, Holt, Rinehart and Winston, Inc., New York- Montreal, Que.-London, 1969.
- [21] D. Henry, *Topics in Nonlinear Analysis*, in: *Trabalho de Matemática*, vol. 192, Brasília, 1982.
- [22] R. Kress, *Linear Integral Equations*, Third edition, in: *Applied Mathematical Sciences*, vol. 82, Springer-Verlag, New York, 2014.
- [23] P.D. Lamberti, M. Perin, On the sharpness of a certain spectral stability estimate for the Dirichlet-Laplacian, *Eurasian Math. J.* 1 (1) (2010) 111–122.
- [24] N.S. Landkof, *Foundations of Modern Potential Theory*, in: *Die Grundlehren der mathematischen Wissenschaften*, Band, vol. 180, Springer-Verlag, New York-Heidelberg, 1972.
- [25] M. Lanza de Cristoforis, Asymptotic behaviour of the conformal representation of a Jordan domain with a small hole in schauder spaces, *Comput. Methods Funct. Theory* 2 (2002) 1–27.
- [26] M. Lanza de Cristoforis, Asymptotic behaviour of the conformal representation of a Jordan domain with a small hole, and relative capacity, in: *Complex Analysis and Dynamical Systems*, in: *Contemp. Math.*, vol. 364, Amer. Math. Soc., Providence, RI, 2004, pp. 155–167.
- [27] M. Lanza de Cristoforis, Asymptotic behaviour of the solutions of the Dirichlet problem for the Laplace operator in a domain with a small hole. a functional analytic approach, *Analysis (Munich)* 28 (2008) 63–93.
- [28] M. Lanza de Cristoforis, Simple Neumann eigenvalues for the Laplace operator in a domain with a small hole. A functional analytic approach, *Rev. Mat. Complut.* 25 (2) (2012) 369–412.
- [29] V.G. Maz'ya, A.B. Movchan, M.J. Nieves, Green's Kernels and Meso-Scale Approximations in Perforated Domains, in: *Lecture Notes in Mathematics*, vol. 2077, Springer, Berlin, 2013.
- [30] V.G. Maz'ya, A.B. Movchan, M.J. Nieves, Eigenvalue problem in a solid with many inclusions: asymptotic analysis, *Multiscale Model. Simul.* 15 (2) (2017) 1003–1047.
- [31] V.G. Maz'ya, S.A. Nazarov, B.A. Plamenevskij, *Asymptotic Theory of Elliptic Boundary Value Problems in Singularly Perturbed Domains. Vol. I*, in: *Operator Theory: Advances and Applications*, vol. 111, Birkhäuser Verlag, Basel, 2000.
- [32] V.G. Maz'ya, S.A. Nazarov, B.A. Plamenevskii, Asymptotic expansions of eigenvalues of boundary value problems for the Laplace operator in domains with small openings, (Russian), *Izv. Akad. Nauk SSSR Ser. Mat.* 48 (2) (1984) 347–371, English translation: *Math. USSR-Izv.* 24(2) (1985) 321–346.

- [33] S. Ozawa, Singular variation of domains and eigenvalues of the Laplacian, *Duke Math. J.* 48 (4) (1981) 767–778.
- [34] G. Pólya, G. Szegő, *Isoperimetric Inequalities in Mathematical Physics*, in: *Annals of Mathematics Studies*, No. 27, Princeton University Press, Princeton, N. J., 1951.
- [35] C. Pommerenke, *Univalent functions*, in: *With a chapter on quadratic differentials by Gerd Jensen* *Studia Mathematica/Mathematische Lehrbücher*, Band XXV. Vandenhoeck & Ruprecht, Göttingen, 1975.
- [36] J. Rauch, M. Taylor, Potential and scattering theory on wildly perturbed domains, *J. Funct. Anal.* 18 (1975) 27–59.
- [37] A. Samarskiĭ, On the influence of constraints on the characteristic frequencies of closed volumes (Russian), *Doklady Akad. Nauk SSSR (N.S.)* 63 (1948) 631–634.
- [38] J. Schauder, Potentialtheoretische untersuchungen, *Math. Z.* 33 (1931) 602–640.
- [39] J. Schauder, Bemerkung zu meiner arbeit potentialtheoretische untersuchungen I (Anhang), *Math. Z.* 35 (1932) 536–538.
- [40] Ya.S. Soĭbel'man, An asymptotic formula for condenser capacity (Russian), *Dokl. Akad. Nauk SSSR* 258 (3) (1981) 590–593.
- [41] T. Valent, *Boundary Value Problems of Finite Elasticity. Local Theorems on Existence, Uniqueness and Analytic Dependence on Data*, Springer-Verlag, New York, 1988.
- [42] W. Wendland, Die fredholmsche alternative für operatoren, die bezüglich eines bilinearen funktionalen adjungiert sind, *Math. Z* 101 (1967) 61–64.
- [43] W. Wendland, Bemerkungen über die fredholmschen sätze, *Methoden Verfahren Math. Phys.* 3, B.I.-Hochschulskripten 722/722a (1970) 141–176.