

# Measuring the gravitational free-fall of antihydrogen

**J. Storey · S. Aghion · O. Ahlén · C. Amsler · A. Ariga · T. Ariga · A. S. Belov · G. Bonomi · P. Bräunig · J. Bremer · R. S. Brusa · L. Cabaret · C. Canali · R. Caravita · F. Castelli · G. Cerchiari · S. Cialdi · D. Comparat · G. Consolati · J. H. Derking · S. Di Domizio · L. Di Noto · M. Doser · A. Dudarev · A. Ereditato · R. Ferragut · A. Fontana · P. Genova · M. Giammarchi · A. Gligorova · S. N. Gninenko · S. Haider · S. D. Hogan · T. Huse · E. Jordan · L. V. Jørgensen · T. Kaltenbacher · J. Kawada · A. Kellerbauer · M. Kimura · A. Knecht · D. Krasnický · V. Lagomarsino · S. Lehner · C. Malbrunot · S. Mariazzi · V. A. Matveev · F. Merkt · F. Moia · G. Nebbia · P. Nédélec · M. K. Oberthaler · N. Pacifico · V. Petráček · C. Pistillo · F. Prelz · M. Prevedelli · C. Regenfus · C. Riccardi · O. Røhne · A. Rotondi · H. Sandaker · P. Scampoli · M. A. Subieta Vasquez · M. Špaček · G. Testera · D. Trezzi · R. Vaccarone · E. Widmann · S. Zavatarelli · J. Zmeskal**

Published online: 7 May 2014

Proceedings of the 11th International Conference on Low Energy Antiproton Physics (LEAP 2013) held in Uppsala, Sweden, 10–15 June, 2013

J. Storey (✉) · C. Amsler · A. Ariga · T. Ariga · A. Ereditato · J. Kawada · M. Kimura · C. Pistillo · P. Scampoli

Albert Einstein Center for Fundamental Physics, Laboratory for High Energy Physics,  
University of Bern, 3012 Bern, Switzerland  
e-mail: james.storey@cern.ch

S. Aghion · G. Consolati · R. Ferragut · F. Moia  
Politecnico di Milano, Piazza Leonardo da Vinci 32, 20133 Milano, Italy

S. Aghion · G. Consolati · R. Ferragut · M. Giammarchi · F. Moia · F. Prelz · D. Trezzi  
Istituto Nazionale di Fisica Nucleare, Sez. di Milano, Via Celoria 16, 20133 Milano, Italy

O. Ahlén · J. Bremer · J. H. Derking · M. Doser · A. Dudarev · S. Haider · L. V. Jørgensen ·  
T. Kaltenbacher · A. Knecht  
Physics Department, European Organisation for Nuclear Research, 1211 Geneva 23, Switzerland

A. S. Belov · S. N. Gninenko · V. A. Matveev  
Institute for Nuclear Research of the Russian Academy of Sciences, Moscow 117312, Russia

G. Bonomi · M. A. Subieta Vasquez  
Department of Mechanical and Industrial Engineering, University of Brescia, Via Branze 38, 25133  
Brescia, Italy

G. Bonomi · A. Fontana · P. Genova · C. Riccardi · A. Rotondi · M. A. Subieta Vasquez  
Istituto Nazionale di Fisica Nucleare, Sez. di Pavia, Via Agostino Bassi 6, 27100 Pavia, Italy

P. Bräunig · M. K. Oberthaler  
Kirchhoff Institute for Physics, University of Heidelberg, Im Neuenheimer Feld 227, 69120  
Heidelberg, Germany

R. S. Brusa · L. Di Noto · S. Mariazzi  
Dipartimento di Fisica, Università di Trento and INFN, Gruppo Collegato di Trento, Via Sommarive  
14, 38050 Povo, Trento, Italy

L. Cabaret · D. Comparat  
Laboratoire Aimé Cotton, CNRS, Université Paris Sud, ENS Cachan, Bâtiment 505, Campus dOrsay,  
91405 Orsay Cedex, France

C. Canali · C. Regenfus  
Physics Institute, University of Zurich, Winterthurerstrasse 190, 8057 Zurich, Switzerland

R. Caravita · F. Castelli · G. Cerchiari · S. Cialdi  
Department of Physics, University of Milano, Via Celoria 16, 20133 Milano, Italy

S. Di Domizio · G. Testera · R. Vaccarone · S. Zavatarelli  
Istituto Nazionale di Fisica Nucleare, Sez. di Genova, Via Dodecaneso 33, 16146 Genova, Italy

A. Gligorova · N. Pacifico · H. Sandaker  
Institute of Physics and Technology, University of Bergen, Alleegaten 55, 5007 Bergen, Norway

S. D. Hogan  
Department of Physics and Astronomy, University College London, Gower Street, London WC1E  
6BT, UK

T. Huse · O. Røhne  
Department of Physics, University of Oslo, Sem Sælands vei 24, 0371 Oslo, Norway

E. Jordan · A. Kellerbauer  
Max Planck Institute for Nuclear Physics, Saupfercheckweg 1, 69117 Heidelberg, Germany

D. Krasnický · V. Lagomarsino  
Department of Physics, University of Genoa, Via Dodecaneso 33, 16146 Genova, Italy

## 1 The AEGIS experiment

The principle of equivalence between gravitational mass and inertial mass is a cornerstone of general relativity. The universality of free-fall (UFF), the experimental evidence on which the weak equivalence principle is based, has been tested to be valid to a very high precision (1 part in 10 trillion) by many experiments using a variety of techniques [1]. However, the UFF has never been tested directly with antimatter. The scientific goal of the AEGIS experiment is to measure, in the first instance, the acceleration of antihydrogen in the Earth's gravitational field with a 1 % accuracy [2].

The conceptual design of the AEGIS experiment is shown in Fig. 1. Details of the design and current status of the AEGIS experiment are described elsewhere in these proceedings [3]. The production of the antihydrogen beam will require many technical problems to be overcome. It is therefore highly desirable that the flux of antihydrogen required for the gravity measurement is minimized. Emulsion based detectors offer the opportunity to reduce the antihydrogen flux required for a measurement of  $\bar{g}$  with a 1 % relative error by enabling the reconstruction of the antihydrogen annihilation vertex with micron-level precision [4].

## 2 Antihydrogen flux requirement for gravity measurement with 1 % accuracy

The experimental challenge is to produce and accelerate antihydrogen atoms through the moiré deflectometer and onto the free-fall detector. For an antihydrogen atom to reach the free-fall detector it must have a radial temperature less than 100 mK. To produce such

---

V. A. Matveev

Joint Institute for Nuclear Research, 141980 Dubna, Russia

S. Lehner · C. Malbrunot · E. Widmann · J. Zmeskal

Stefan Meyer Institute for Subatomic Physics, Austrian Academy of Sciences, Boltzmannngasse 3,  
1090 Vienna, Austria

F. Merkt

ETH Zurich, Laboratory for Physical Chemistry, 8093 Zurich, Switzerland

G. Nebbia

Istituto Nazionale di Fisica Nucleare, Sez. di Padova, Via Marzolo 8, 35131 Padova, Italy

P. Nédélec

Institut de Physique Nucléaire de Lyon, Claude Bernard University Lyon 1, 4 Rue Enrico Fermi,  
69622 Villeurbanne, France

V. Petráček · M. Špaček

FNSPE, Břehová, Czech Technical University in Prague, 11519 Praha 1, Czech Republic

M. Prevedelli


Department of Physics, University of Bologna, Via Irnerio 46, 40126 Bologna, Italy

C. Riccardi · A. Rotondi

Department of Nuclear and Theoretical Physics, University of Pavia, Via Bassi 6, 27100 Pavia, Italy

P. Scamporrì


Department of Physics, University of Napoli Federico II, Via Cinthia, 80126 Napoli, Italy



**Fig. 1** Conceptual design of the AEgIS experiment. The divergent antihydrogen beam traverses a pair of identical gratings with horizontal slits (moiré deflectometer) and annihilates on the position and time sensitive free-fall detector. The free-fall detector is based on a hybrid of three detector technologies: nuclear emulsions, a scintillating fiber tracker and a silicon strip detector. Emulsion based detectors provide unrivaled positional information but no time information. Therefore, a scintillating fiber tracker and silicon strip detector are used to measure the arrival time of each antihydrogen anti-atom on the free-fall detector with a  $1 \approx m$  time resolution. The time of flight (TOF) of each anti-atom is inferred from the arrival time on the free-fall detector and the (known) time at which the pulsed antihydrogen beam is produced. A *light pattern* is produced by illuminating the gratings homogeneously with an incoherent light source (LED with spatial diffuser) and is recorded directly on the emulsion film. The *shadow-like spatial* distribution of the annihilation vertices (*solid line*) is displaced relative to that of light (*dashed lines*) by an amount which depends on the time of flight between the gratings and the gravitational strength  $\bar{g}$  between the antihydrogen atom and the Earth

atoms two experimental approaches will be pursued: the production of antihydrogen which is as cold as possible and the synthesis of as many anti-atoms as possible. Since there are considerable technical difficulties to be overcome, it is illuminating to investigate the absolute antihydrogen flux required for a measurement of gravity with a 1 % accuracy and the dependency of this flux on the spatial resolution of the free-fall detector.

The influence of the spatial resolution of the free-fall detector on the flux of antihydrogen required for a measurement of  $\bar{g}$  with a 1 % precision has been studied with a Monte Carlo Simulation. Antihydrogen atoms were uniformly generated in a cylinder of 3 mm transverse radius and 2 cm axial length, with initial velocities distributed according to a Maxwell-Boltzmann distribution with a mean temperature of 100 mK. The atoms were then accelerated towards the moiré deflectometer (pitch  $d=40 \approx m$  between horizontal slits of  $12 \approx m$  width) and the free-fall detector. A Gaussian smearing was then applied to the annihilation vertex to simulate the detector resolution. Figure 2 left, shows the expected intensity distribution as a function of the phase  $\phi = (2\pi/d)\Delta x$ , where  $\Delta x = \bar{g}T^2$  and T is the time of flight between the 2 gratings. The contrast of the shadow-like image is greatly improved by a higher spatial resolution. The benefit of the improved contrast is shown in Fig. 2 right, which shows the dependency of the relative  $\bar{g}$  precision on the antihydrogen flux for various spatial resolutions. A free-fall detector with a  $\sim 2 \approx m$  spatial resolution requires  $\sim 600$  fully reconstructed and time tagged antihydrogen atoms for a 1 % relative error on  $\bar{g}$ , which compared to a  $10 \approx m$  spatial resolution is an order of magnitude reduction in the required antihydrogen flux.



**Fig. 2** *Left*: Simulated distribution of reconstructed annihilation vertices on the free-fall detector over one grating period for vertex resolutions between 1 and 15  $\mu\text{m}$  for an antihydrogen beam with a velocity of 700 m/s. The contrast of the shadow-like image increases with improved vertex resolution. *Right*: Simulation of the relative precision on  $\bar{g}$  vs. the number of fully reconstructed and time tagged antihydrogen atoms for vertex resolutions between 1 and 15  $\mu\text{m}$ . The horizontal black line indicates a 1% relative error on  $\bar{g}$  [5]


### 3 Reconstruction of $\bar{\text{H}}$ annihilation vertices with micron level spatial resolution

In light of the considerable benefit to AEGIS of a free-fall detector with a 1–2  $\mu\text{m}$  spatial resolution, a detector based on nuclear emulsions is currently under development. Nuclear emulsions are photographic films which are optimized for use as particle tracking detectors and are still the most precise particle tracking detector technology currently available [6]. A typical detector consists of a gel with a suspension of silver bromide crystals, in which a track is formed after the passage of an ionizing particle. After chemical development silver grains along the path of the track, with dimensions of 1  $\mu\text{m}$  or less, are visible with an optical microscope. With just a single 50  $\mu\text{m}$  layer it is possible for the full three-dimensional reconstruction of the particle track and a measurement of the energy loss. The intrinsic resolution for tracks in a nuclear emulsion is a mere 50 nm [5].


In order to evaluate the suitability of an emulsion based free-fall detector for AEGIS, a series of measurements were performed during the 2012 antiproton decelerator (AD) run at CERN. The 5 MeV antiprotons from the AD were directed onto a nuclear emulsion the surface of which was partially covered with a 20  $\mu\text{m}$  stainless steel foil. The foil simulates the separation window which will be used in the actual experiment to isolate the emulsion detector from the ultra high vacuum region. After passing through a series of degrading foils and vacuum separation windows, the average energy of the antiprotons reaching the emulsion was  $\approx 150$  keV.

Images of the antiproton annihilations in the bare emulsion are shown in Fig. 3. A full 3D image of the annihilation star is constructed by focusing the microscope at different depths in the emulsion. The nuclear fragments and charged pions from the antiproton annihilation are clearly discernible in the emulsion as black (heavily ionising) and thin (minimum ionising) tracks respectively.

In order to determine the vertex resolution that is achievable for the  $\bar{g}$  experiment, the vertex position was determined by finding the coordinates of the point which minimised the quadratic sum of distances to all tracks. Figure 4 left, shows the distribution of the vertex position in the direction of the antiproton beam axis. For the bare sample the distribution has a sharp peak at the surface of the emulsion, while the distribution for the stainless steel sample is much broader due to the fact that the foil was not in perfect contact with the



**Fig. 3** *Left*: Antiproton annihilations in a bare emulsion. *Blue circles* indicate the location of the annihilations. *Right*: 3D image of antiproton annihilation in a new emulsion prepared at LHEP from a gel produced by the University of Nagoya. The size of the 3D image is  $150 \times 120 \times 50 \text{ }\mu\text{m}$




**Fig. 4** *Left*: Annihilation vertex resolution in the direction of the antiproton beam (perpendicular to the emulsion surface). *Right*: Impact parameter distribution for annihilations [5]

emulsion surface. Figure 4 right, shows the distribution of the distance between the reconstructed vertex and the tracks (impact parameter). This demonstrates that an r.m.s resolution of  $\sigma \simeq 1 \text{ }\mu\text{m}$  on the vertical position of the annihilation can, in principal, be achieved with an emulsion based detector.

#### 4 Conceptual design for a free-fall detector

Emulsion based detectors provide spatial but no temporal information. To measure the time of flight of the antihydrogen atom requires that the time of the annihilation on the emulsion is also measured. A conceptual design for a hybrid detector that combines the emulsion detector with timing information from a scintillating/silicon tracker and silicon strip detector is shown in Fig. 5. The time of the annihilation is determined from information from both the silicon strip detector and the downstream tracker. The tracker will also be used to seed the search for annihilation vertices in the emulsion by reconstructing the annihilation pions.



**Fig. 5** Conceptual design for a free-fall detector for AEGIS. The emulsion detector is used to identify the precise location of the annihilation vertex and the downstream tracker and silicon strip detector are used to measure the time of the annihilation

## 5 Summary

Emulsion based detectors can be used to reconstruct the annihilation vertex position with an r.m.s resolution of  $\sigma \simeq 1 \text{ }\mu\text{m}$  on the vertical position of the annihilation. If such a precision can be realised in the final free-fall detector, a measurement of  $\bar{g}$  with 1 % relative error will require the reconstruction of  $\sim 600$  antihydrogen atoms. A hybrid detector which combines the spatial precision of emulsion detectors with temporal information from silicon/scintillating fiber technologies is currently under investigation.

**Acknowledgments** We warmly acknowledge our technical collaborators Luca Dassa, Roger Hänni and Jacky Rochet without whom the experiment could not have been built. E.W. wishes to acknowledge support by the European Research Council grant no. 291242-HBAR-HFS and the Austrian Ministry for Science and Research. J.S. wishes to acknowledge support from the Swiss National Science Foundation.

## References

1. Heckel, B. et al.: Adv. Space Res. **25**(6), 1225–1230 (2000)
2. AEGIS-proto collaboration: Proposed antimatter gravity measurement with an antihydrogen beam. Nucl. Instrum. Meth. B **266**, 351 (2008)
3. Knecht, A., et al.: in these proceedings
4. Amsler, C. et al.: A new application of emulsions to measure the gravitational force on antihydrogen. JINST **8**, P02015 (2013). arXiv:1211.1370
5. Aghion, S. et al.: JINST **8**, P08013 (2013)
6. de Lellis, G., Ereditato, A., Niwa, K.: Nuclear emulsions. In: Fabjan, C.W., Schopper, H. (eds) Springer Materials, Landolt-Börnstein Database. Springer, Heidelberg (2011). <http://www.springermaterials.com>