

A diffuse interface model for two-phase incompressible flows with nonlocal interactions and nonconstant mobility

Sergio Frigeri ^{*} Maurizio Grasselli[†] Elisabetta Rocca[‡]

Abstract

We consider a diffuse interface model for incompressible isothermal mixtures of two immiscible fluids with matched constant densities. This model consists of the Navier-Stokes system coupled with a convective nonlocal Cahn-Hilliard equation with non-constant mobility. We first prove the existence of a global weak solution in the case of non-degenerate mobilities and regular potentials of polynomial growth. Then we extend the result to degenerate mobilities and singular (e.g. logarithmic) potentials. In the latter case we also establish the existence of the global attractor in dimension two. Using a similar technique, we show that there is a global attractor for the convective nonlocal Cahn-Hilliard equation with degenerate mobility and singular potential in dimension three.

Keywords: Navier-Stokes equations, nonlocal Cahn-Hilliard equations, degenerate mobility, incompressible binary fluids, weak solutions, global attractors.

MSC 2010: 35Q30, 37L30, 45K05, 76D03, 76T99.

^{*}Weierstrass Institute for Applied Analysis and Stochastics, Mohrenstr. 39, D-10117 Berlin, Germany. E-mail: *SergioPietro.Frigeri@wias-berlin.de* The author is supported by the FP7-IDEAS-ERC-StG Grant #256872 (EntroPhase)

[†]Dipartimento di Matematica, Politecnico di Milano, Milano I-20133, Italy. E-mail: *maurizio.grasselli@polimi.it*

[‡]Weierstrass Institute for Applied Analysis and Stochastics, Mohrenstr. 39, D-10117 Berlin, Germany. E-mail: *Elisabetta.Rocca@wias-berlin.de* The author is supported by the FP7-IDEAS-ERC-StG Grant #256872 (EntroPhase)

1 Introduction

Model H is a diffuse interface model for incompressible isothermal two-phase flows which consists of the Navier-Stokes equations for the (averaged) velocity u nonlinearly coupled with a convective Cahn-Hilliard equation for the (relative) concentration difference φ (cf., for instance, [7, 31, 32, 33, 34, 35, 39]). The resulting evolution system has been studied by several authors (see, e.g., [1, 2, 12, 13, 22, 23, 24, 41, 49, 51, 52] and references therein, cf. also [5, 11, 36, 30] for models with shear dependent viscosity). In the case of matched densities, setting the constant density equal to one, the system can be written as follows

$$u_t - \nu \Delta u + (u \cdot \nabla)u + \nabla \pi = \mu \nabla \varphi + h \quad \text{in } \Omega \times (0, T) \quad (1.1)$$

$$\operatorname{div}(u) = 0 \quad \text{in } \Omega \times (0, T) \quad (1.2)$$

$$\varphi_t + u \cdot \nabla \varphi = \operatorname{div}(m(\varphi) \nabla \mu) \quad \text{in } \Omega \times (0, T) \quad (1.3)$$

$$\mu = -\sigma \Delta \varphi + \frac{1}{\sigma} F'(\varphi) \quad \text{in } \Omega \times (0, T), \quad (1.4)$$

where $\Omega \subset \mathbb{R}^d$, $d = 2, 3$, is a bounded domain. Here $\nu > 0$ is the viscosity (supposed to be constant for simplicity), π is the pressure, h is a given (non-gradient) external force, m is the so-called mobility and $\sigma > 0$ is related to the (diffuse) interface thickness.

A more realistic version of the Cahn-Hilliard equation is characterized by a (spatially) nonlocal free energy. The physical relevance of nonlocal interactions was already pointed out in the pioneering paper [46] (see also [17, 4.2] and references therein). Though isothermal and nonisothermal models containing nonlocal terms have only recently been studied from the analytical viewpoint (cf., e.g., [10, 15, 26, 27, 28, 29, 38] and their references). The difference between local and nonlocal models consists in the choice of the interaction potential. The nonlocal contribution to the free energy has typically the form $\int_{\Omega} J(x, y) |\varphi(x) - \varphi(y)|^2 dy$ with a given symmetric kernel J defined on $\Omega \times \Omega$; its local Ginzburg-Landau counterpart has the form $(\sigma/2) |\nabla \varphi(x)|^2$ with a positive parameter σ . The latter can be obtained as a formal limit as $\zeta \rightarrow \infty$ from the nonlocal one with the choice $J(x, y) = \zeta^{d+2} J(|\zeta(x - y)|^2)$, where J is a nonnegative function with support in $[0, 1]$. This follows from the formula formally deduced in [37]

$$\begin{aligned} \int_{\Omega} \zeta^{d+2} J(|\zeta(x - y)|^2) |\varphi(x) - \varphi(y)|^2 dy &= \int_{\Omega_{\zeta}(x)} J(|z|^2) \left| \frac{\varphi\left(x + \frac{z}{\zeta}\right) - \varphi(x)}{\frac{1}{\zeta}} \right|^2 dz \\ &\xrightarrow{\zeta \rightarrow \infty} \int_{\mathbb{R}^d} J(|z|^2) \langle \nabla \varphi(x), z \rangle^2 dz = \frac{\sigma}{2} |\nabla \varphi(x)|^2 \end{aligned}$$

for a sufficiently regular φ , where $\sigma = 2/d \int_{\mathbb{R}^d} J(|z|^2) |z|^2 dz$ and $\Omega_{\zeta}(x) = \zeta(\Omega - x)$. Here we have used the identity $\int_{\mathbb{R}^d} J(|z|^2) \langle e, z \rangle^2 dz = 1/d \int_{\mathbb{R}^d} J(|z|^2) |z|^2 dz$ for every unit vector

$e \in \mathbb{R}^d$. As a consequence, the Cahn-Hilliard equation (1.3)-(1.4) can be viewed as an approximation of the nonlocal one.

Nonlocal interactions have been taken into account in a series of recent papers (see [14, 18, 19, 20, 21]) where a modification of the model H with matched densities has been considered and analyzed. More precisely, a system of the following form has been considered (cf. [14])

$$\varphi_t + u \cdot \nabla \varphi = \operatorname{div}(m(\varphi)\nabla\mu) \quad \text{in } \Omega \times (0, T) \quad (1.5)$$

$$\mu = a\varphi - J * \varphi + F'(\varphi) \quad \text{in } \Omega \times (0, T) \quad (1.6)$$

$$u_t - \nu\Delta u + (u \cdot \nabla)u + \nabla\pi = \mu\nabla\varphi + h \quad \text{in } \Omega \times (0, T) \quad (1.7)$$

$$\operatorname{div}(u) = 0 \quad \text{in } \Omega \times (0, T) \quad (1.8)$$

$$\frac{\partial\mu}{\partial n} = 0, \quad u = 0 \quad \text{on } \partial\Omega \times (0, T) \quad (1.9)$$

$$u(0) = u_0, \quad \varphi(0) = \varphi_0 \quad \text{in } \Omega, \quad (1.10)$$

where n stands for the outward normal to $\partial\Omega$, while u_0 and φ_0 are given initial conditions. Here the interaction kernel $J : \mathbb{R}^d \rightarrow \mathbb{R}$ is an even function and $a(x) = \int_{\Omega} J(x-y)dy$.

Nonlocal system (1.5)-(1.8) is more challenging with respect to (1.1)-(1.4), even in dimension two. One of the reasons is that φ has a poorer regularity and this influences the treatment of the Navier-Stokes system through the so-called Korteweg force $\mu\nabla\varphi$. For instance, uniqueness in dimension two is an open issue under sufficiently general conditions which ensures the existence of a weak solution (see [14, Remark 8], cf. also [18]). Due to this difficulty, only the constant mobility case has been considered so far (though viscosity depending on φ has been handled). Let us briefly recall the main existing results for system (1.5)–(1.8) with m constant.

In [14] the authors proved the existence of global dissipative weak solutions in 2D and 3D, and study some regularity properties, for the case of regular potentials of arbitrary polynomial growth. For such potentials the longterm behavior of weak solutions was analyzed in [19]. More precisely, the existence of the global attractor in 2D and of the trajectory attractor in 3D were established. In [20] the previous results of [14, 19] were extended to the case of singular potentials. The existence of (unique) strong solutions in 2D for regular potentials with arbitrary polynomial growth were obtained in [21]. There, in addition, the regularity of the global attractor and the convergence of weak solutions to single equilibria were shown.

Uniqueness of weak solutions in 2D has been demonstrated only recently for both regular and singular potentials (see [18]). In the same paper, further results have been proven. For instance, the existence of strong solutions and the weak-strong uniqueness for the case of nonconstant (i.e., φ -dependent) viscosity and regular potentials, the ex-

istence of exponential attractors (regular potentials), and the connectedness of the global attractor in the case of constant viscosity.

On the other hand, despite the variety of results with m constant, in the rigorous derivation of the nonlocal Cahn-Hilliard equation done in [27] the mobility depends on φ and degenerates at the pure phases, while the potential is of logarithmic type. This motivates the main goal of this contribution, namely, the analysis of the so-called non-local Cahn-Hilliard-Navier-Stokes system in the case of degenerate mobility and singular potential.

The local Cahn-Hilliard equation with degenerate mobility (i.e., system (1.5–1.10) with $u = 0$) was considered in the seminal paper [16], where the authors established the existence of a weak solution (cf. also [40, 45] and references therein, for non-degenerate mobility see [9, 47]). This result was then extended to the standard (local) Cahn-Hilliard-Navier-Stokes system in [12]. The nonlocal Cahn-Hilliard equation with degenerate mobility and logarithmic potential was rigorously justified and analyzed in [27] (see also [29] and references therein). In particular, in the case of periodic boundary conditions, an existence and uniqueness result was proven in [28]. Then a more general case was considered in [26]. The convergence to single equilibria was recently studied in [42, 43] (cf. also [25] for further results).

Inspired by the strategy devised in [16], we first analyze the nonlocal system by taking a non-degenerate mobility m and a regular potential F with polynomial growth. We prove the existence of a global weak solution which satisfies an energy inequality (equality if $d = 2$). This result extends [19] and allows us to construct a rigorous approximation of the case where m is degenerate and F is singular (e.g. logarithmic). Therefore we can pass to the limit and obtain a similar result for the latter case. In addition, since the energy identity holds in two dimensions, we can construct a generalized semiflow which possesses a global attractor by using Ball’s method (see [8]). In the above mentioned recent contribution [18], uniqueness of weak solutions to system (1.5)-(1.8) with degenerate mobility and singular potential in 2D has also been proven (constant viscosity). Hence, we can say that our semiflow is indeed a semigroup and the global attractor is connected. Regarding the 3D case, the validity of a suitable energy inequality (cf. (3.45)) allows to generalize the results on the trajectory attractors (cf. [19, 20]) to system (1.5)-(1.8) with degenerate mobility and singular potential. By means of Ball’s approach, we can also show that the convective nonlocal Cahn-Hilliard equation with degenerate mobility and singular potential possesses a (connected) global attractor. In this case uniqueness can be proven even in dimension three. Note that this result entails, in particular, that the nonlocal Cahn-Hilliard equation which has been obtained as hydrodynamic limit in [27] has indeed a global attractor.

We point out that uniqueness of solutions is still an open issue for the local Cahn-Hilliard equation analyzed in [16]. This is one of the main advantages of the nonlocal versus the local. We remind that uniqueness of weak solutions and continuous dependence estimates are fundamental starting points, for example, in view of the study of related optimal control problems. This issue will be the subject of forthcoming contributions.

Let us notice here that the main difficulty encountered while dealing with the degenerate mobility case is that the gradient of the chemical potential μ in (1.6) can no longer be controlled in any L^p space. Hence, in order to get an existence result a suitable notion of weak solution needs to be introduced (cf. [16] for the local Cahn-Hilliard equation). More precisely, in this new formulation the gradient of μ does not appear anymore (cf. Definition 2 in Section 4). It worth observing that, in the present case, our main Thm. 2 does not require the (conserved) mean value of the order parameter φ to be strictly in between -1 and 1 , but $|\int_{\Omega} \varphi_0| \leq |\Omega|$ suffices. Thus the model allows pure phase solutions for all $t \geq 0$. This is not possible in the case of constant or *strongly degenerate* mobility (cf. Remarks 7 and 8 for further comments on this topic).

We conclude by observing that it would be particularly interesting (albeit nontrivial) to extend the present and previous results on nonlocal Cahn-Hilliard-Navier-Stokes systems to a model with unmatched densities (for the local case see [3, 4] and references therein) or to the compressible case (cf. [6] for the local Cahn-Hilliard equation).

The plan of the paper goes as follows. In Section 2 we introduce the notation and the functional setting. Section 3 is devoted to prove the existence of a global weak solution satisfying a suitable energy inequality (equality in the 2D case) when the mobility is non-degenerate and F is regular. In Section 4, using a convenient approximation scheme, we extend the proven result to the case of degenerate mobility and a singular F (e.g. of logarithmic type). Some regularity issues for φ and μ are also discussed. Section 5 is devoted to the existence of global attractor in the two dimensional case. Finally, in Section 6, we consider the convective nonlocal Cahn-Hilliard equation with degenerate mobility. We deduce the existence of a global weak solution from the previous result for the coupled system. Then, even in dimension three, we establish the uniqueness of weak solutions as well as the existence of a (connected) global attractor under rather general assumptions on F , J and m .

2 Notation and functional setting

We set $H := L^2(\Omega)$ and $V := H^1(\Omega)$, where Ω is supposed to have a sufficiently smooth boundary (say, e.g., of class $C^{1,1}$). If X is a (real) Banach space, X' will denote its dual. For every $f \in V'$ we denote by \bar{f} the average of f over Ω , i.e., $\bar{f} := |\Omega|^{-1} \langle f, 1 \rangle$. Here $|\Omega|$

is the Lebesgue measure of Ω . Let us introduce also the spaces $V_0 := \{v \in V : \bar{v} = 0\}$, $V'_0 := \{f \in V' : \bar{f} = 0\}$ and the operator $A : V \rightarrow V'$, $A \in \mathcal{L}(V, V')$ defined by

$$\langle Au, v \rangle := \int_{\Omega} \nabla u \cdot \nabla v \quad \forall u, v \in V.$$

We recall that A maps V onto V'_0 and the restriction of A to V_0 maps V_0 onto V'_0 isomorphically. Let us denote by $\mathcal{N} : V'_0 \rightarrow V_0$ the inverse map defined by

$$A\mathcal{N}f = f, \quad \forall f \in V'_0 \quad \text{and} \quad \mathcal{N}Au = u, \quad \forall u \in V_0.$$

As is well known, for every $f \in V'_0$, $\mathcal{N}f$ is the unique solution with zero mean value of the Neumann problem

$$\begin{cases} -\Delta u = f, & \text{in } \Omega \\ \frac{\partial u}{\partial n} = 0, & \text{on } \partial\Omega. \end{cases}$$

Furthermore, the following relations hold

$$\begin{aligned} \langle Au, \mathcal{N}f \rangle &= \langle f, u \rangle, \quad \forall u \in V, \quad \forall f \in V'_0, \\ \langle f, \mathcal{N}g \rangle &= \langle g, \mathcal{N}f \rangle = \int_{\Omega} \nabla(\mathcal{N}f) \cdot \nabla(\mathcal{N}g), \quad \forall f, g \in V'_0. \end{aligned}$$

Recall that A can be also viewed as an unbounded operator $A : D(A) \subset H \rightarrow H$ from the domain $D(A) = \{\phi \in H^2(\Omega) : \partial\phi/\partial n = 0 \text{ on } \partial\Omega\}$ onto H and that \mathcal{N} can also be viewed as a self-adjoint compact operator $\mathcal{N} = A^{-1} : H \rightarrow H$ in H . Hence, the fractional powers A^r and \mathcal{N}^s , for $r, s \geq 0$, can be defined through classical spectral theory.

We shall repeatedly need the standard Hilbert spaces for the Navier-Stokes equations with no-slip boundary condition (see, e.g., [50])

$$G_{div} := \overline{\{u \in C_0^\infty(\Omega)^d : \operatorname{div}(u) = 0\}}^{L^2(\Omega)^d}, \quad V_{div} := \{u \in H_0^1(\Omega)^d : \operatorname{div}(u) = 0\}.$$

We denote by $\|\cdot\|$ and (\cdot, \cdot) the norm and the scalar product on both H and G_{div} , respectively. Instead, V_{div} is endowed with the scalar product

$$(u, v)_{V_{div}} = (\nabla u, \nabla v), \quad \forall u, v \in V_{div}.$$

In the proof of Theorem 1 we shall introduce the family of the eigenfunctions of the Stokes operator S with no-slip boundary condition. We recall that $S : D(S) \subset G_{div} \rightarrow G_{div}$ is defined as $S := -P\Delta$ with domain $D(S) = H^2(\Omega)^d \cap V_{div}$, where $P : L^2(\Omega)^d \rightarrow G_{div}$ is the Leray projector. Notice that we have

$$(Su, v) = (u, v)_{V_{div}} = (\nabla u, \nabla v), \quad \forall u \in D(S), \quad \forall v \in V_{div},$$

and $S^{-1} : G_{div} \rightarrow G_{div}$ is a self-adjoint compact operator in G_{div} . Thus, according with classical results, S possesses a sequence of eigenvalues $\{\lambda_j\}$ with $0 < \lambda_1 \leq \lambda_2 \leq \dots$ and

$\lambda_j \rightarrow \infty$, and a family $\{w_j\} \subset D(S)$ of eigenfunctions which is orthonormal in G_{div} . Let us also recall the Poincaré inequality

$$\lambda_1 \|u\|^2 \leq \|\nabla u\|^2, \quad \forall u \in V_{div}.$$

The trilinear form b which appears in the weak formulation of the Navier-Stokes equations is defined as follows

$$b(u, v, w) = \int_{\Omega} (u \cdot \nabla) v \cdot w, \quad \forall u, v, w \in V_{div}.$$

We recall that we have

$$b(u, w, v) = -b(u, v, w), \quad \forall u, v, w \in V_{div}.$$

For the basic estimates satisfied by the trilinear form b the reader is referred to, e.g., [50].

Finally, if X is a (real) Banach space and $\tau \in \mathbb{R}$, we shall denote by $L_{tb}^p(\tau, \infty; X)$, $1 \leq p < \infty$, the space of functions $f \in L_{loc}^p([\tau, \infty); X)$ that are translation bounded in $L_{loc}^p([\tau, \infty); X)$, i.e. such that

$$\|f\|_{L_{tb}^p(\tau, \infty; X)}^p := \sup_{t \geq \tau} \int_t^{t+1} \|f(s)\|_X^p ds < \infty. \quad (2.1)$$

3 Non-degenerate mobility

Let us first consider the case where the mobility m does not degenerate, i.e. m satisfies the following assumption

(H1) $m \in C_{loc}^{0,1}(\mathbb{R})$ and there exist $m_1, m_2 > 0$ such that

$$m_1 \leq m(s) \leq m_2, \quad \forall s \in \mathbb{R}.$$

The other assumptions we need are the ones on the kernel J , on the potential F and the forcing term h which are the same as in [14]

(H2) $J(\cdot - x) \in W^{1,1}(\Omega)$ for almost any $x \in \Omega$ and satisfies

$$J(x) = J(-x), \quad a(x) := \int_{\Omega} J(x - y) dy \geq 0, \quad \text{a.e. } x \in \Omega,$$

$$a^* := \sup_{x \in \Omega} \int_{\Omega} |J(x - y)| dy < \infty, \quad b := \sup_{x \in \Omega} \int_{\Omega} |\nabla J(x - y)| dy < \infty.$$

(H3) $F \in C_{loc}^{2,1}(\mathbb{R})$ and there exists $c_0 > 0$ such that

$$F''(s) + a(x) \geq c_0, \quad \forall s \in \mathbb{R}, \quad \text{a.e. } x \in \Omega.$$

(H4) There exist $c_1 > (a^* - a_*)/2$, where

$$a_* := \inf_{x \in \Omega} \int_{\Omega} J(x-y) dy,$$

and $c_2 \in \mathbb{R}$ such that

$$F(s) \geq c_1 s^2 - c_2, \quad \forall s \in \mathbb{R}.$$

(H5) There exist $c_3 > 0$, $c_4 \geq 0$ and $r \in (1, 2]$ such that

$$|F'(s)|^r \leq c_3 |F(s)| + c_4, \quad \forall s \in \mathbb{R}.$$

(H6) $h \in L^2(0, T; V'_{div})$ for all $T > 0$.

Some further regularity properties of the weak solution can be established by using the following assumption

(H7) $F \in C_{loc}^{2,1}(\mathbb{R})$ and there exist $c_5 > 0$, $c_6 > 0$ and $p > 2$ such that

$$F''(s) + a(x) \geq c_5 |s|^{p-2} - c_6, \quad \forall s \in \mathbb{R}, \quad \text{a.e. } x \in \Omega.$$

Remark 1. A well known example of potential F and kernel J which satisfy the above conditions is given by $F(s) = (s^2 - 1)^2$ and $J(x) = j_3 |x|^{-1}$, if $d = 3$, and $J(x) = -j_2 \log |x|$, if $d = 2$, where j_2 and j_3 are positive constants.

Before stating the main result of this section, let us recall the definition of weak solution to system (1.5)–(1.10).

Definition 1. Let $u_0 \in G_{div}$, $\varphi_0 \in H$ such that $F(\varphi_0) \in L^1(\Omega)$, and $0 < T < \infty$ be given. Then, a couple $[u, \varphi]$ is a weak solution to (1.5)–(1.10) on $[0, T]$ if

$$u \in L^\infty(0, T; G_{div}) \cap L^2(0, T; V_{div}), \quad \varphi \in L^\infty(0, T; H) \cap L^2(0, T; V), \quad (3.1)$$

$$u_t \in L^{4/3}(0, T; V'_{div}), \quad \varphi_t \in L^{4/3}(0, T; V'), \quad \text{if } d = 3, \quad (3.2)$$

$$u_t \in L^{2-\gamma}(0, T; V'_{div}), \quad \varphi_t \in L^{2-\delta}(0, T; V') \quad \gamma, \delta \in (0, 1), \quad \text{if } d = 2, \quad (3.3)$$

$$\mu := a\varphi - J * \varphi + F'(\varphi) \in L^2(0, T; V), \quad (3.4)$$

and the following variational formulation is satisfied for almost any $t \in (0, T)$

$$\langle \varphi_t, \psi \rangle + (m(\varphi) \nabla \mu, \nabla \psi) = (u \varphi, \nabla \psi), \quad \forall \psi \in V, \quad (3.5)$$

$$\langle u_t, v \rangle + \nu(\nabla u, \nabla v) + b(u, u, v) = -(\varphi \nabla \mu, v) + \langle h, v \rangle, \quad \forall v \in V_{div}, \quad (3.6)$$

together with the initial conditions (1.10).

Remark 2. It is easy to see that $u \in C_w([0, T]; G_{div})$ and $\varphi \in C_w([0, T]; H)$. Hence, the initial conditions (1.10) make sense.

Theorem 1. Let $u_0 \in G_{div}$, $\varphi_0 \in H$ such that $F(\varphi_0) \in L^1(\Omega)$, and suppose that (H1)–(H6) are satisfied. Then, for every given $T > 0$, there exists a weak solution $[u, \varphi]$ to (1.5)–(1.10) in the sense of Definition 1 satisfying the energy inequality

$$\mathcal{E}(u(t), \varphi(t)) + \int_0^t \left(\nu \|\nabla u\|^2 + \|\sqrt{m(\varphi)} \nabla \mu\|^2 \right) d\tau \leq \mathcal{E}(u_0, \varphi_0) + \int_0^t \langle h(\tau), u \rangle d\tau, \quad (3.7)$$

for every $t > 0$, where we have set

$$\mathcal{E}(u(t), \varphi(t)) = \frac{1}{2} \|u(t)\|^2 + \frac{1}{4} \int_{\Omega} \int_{\Omega} J(x-y) (\varphi(x, t) - \varphi(y, t))^2 dx dy + \int_{\Omega} F(\varphi(t)). \quad (3.8)$$

Furthermore, assume that assumption (H4) is replaced by (H7). Then, for every $T > 0$ there exists a weak solution $[u, \varphi]$ to (1.5)–(1.10) on $[0, T]$ corresponding to $[u_0, \varphi_0]$ (in the sense of Definition 1) satisfying (3.1), (3.2) and also

$$\varphi \in L^\infty(0, T; L^p(\Omega)), \quad (3.9)$$

$$\varphi_t \in L^2(0, T; V'), \quad \text{if } d = 2 \quad \text{or} \quad (d = 3 \text{ and } p \geq 3), \quad (3.10)$$

$$u_t \in L^2(0, T; V'_{div}), \quad \text{if } d = 2. \quad (3.11)$$

Finally, assume that $d = 2$ and (H4) is replaced by (H7). Then,

1. any weak solution satisfies the energy identity

$$\frac{d}{dt} \mathcal{E}(u, \varphi) + \nu \|\nabla u\|^2 + \|\sqrt{m(\varphi)} \nabla \mu\|^2 = \langle h(t), u \rangle, \quad t > 0. \quad (3.12)$$

In particular we have $u \in C([0, \infty); G_{div})$, $\varphi \in C([0, \infty); H)$ and $\int_{\Omega} F(\varphi) \in C([0, \infty))$.

2. If in addition $h \in L^2_{tb}(0, \infty; V'_{div})$, then any weak solution satisfies also the dissipative estimate

$$\mathcal{E}(u(t), \varphi(t)) \leq \mathcal{E}(u_0, \varphi_0) e^{-kt} + F(m_0) |\Omega| + K, \quad \forall t \geq 0, \quad (3.13)$$

where $m_0 = (\varphi_0, 1)$ and k, K are two positive constants which are independent of the initial data, with K depending on Ω, ν, J, F and $\|h\|_{L^2_{tb}(0, \infty; V'_{div})}$.

Proof. The argument follows the lines of [14, Proof of Theorem 1] and is based on a Faedo-Galerkin approximation scheme. For the reader's convenience, we give a sketch of it. Let us first assume that $\varphi_0 \in D(B)$, where the operator B is given by $B = -\Delta + I$ with homogeneous Neumann boundary condition. We introduce the family $\{w_j\}_{j \geq 1}$ of the

eigenfunctions of the Stokes operator S as a Galerkin base in V_{div} and the family $\{\psi_j\}_{j \geq 1}$ of the eigenfunctions of B as a Galerkin base in V . We define the n -dimensional subspaces $\mathcal{W}_n := \langle w_1, \dots, w_n \rangle$ and $\Psi_n := \langle \psi_1, \dots, \psi_n \rangle$ and consider the orthogonal projectors on these subspaces in G_{div} and H , respectively, i.e., $\tilde{P}_n := P_{\mathcal{W}_n}$ and $P_n := P_{\Psi_n}$. We then look for three functions of the form

$$u_n(t) = \sum_{k=1}^n \alpha_k^{(n)}(t) w_k, \quad \varphi_n(t) = \sum_{k=1}^n \beta_k^{(n)}(t) \psi_k, \quad \mu_n(t) = \sum_{k=1}^n \gamma_k^{(n)}(t) \psi_k$$

which solve the following approximating problem

$$(\varphi'_n, \psi) + (m(\varphi_n) \nabla \mu_n, \nabla \psi) = (u_n \varphi_n, \nabla \psi) \quad (3.14)$$

$$(u'_n, v) + \nu(\nabla u_n, \nabla v) + b(u_n, u_n, v) = -(\varphi_n \nabla \mu_n, v) + (h_n, v) \quad (3.15)$$

$$\mu_n = P_n(a\varphi_n - J * \varphi_n + F'(\varphi_n)) \quad (3.16)$$

$$\varphi_n(0) = \varphi_{0n}, \quad u_n(0) = u_{0n}, \quad (3.17)$$

for every $\psi \in \Psi_n$ and every $v \in \mathcal{W}_n$, where $\varphi_{0n} = P_n \varphi_0$ and $u_{0n} = \tilde{P}_n u_0$ (primes denote derivatives with respect to time). In (3.15) $\{h_n\}$ is a sequence in $C^0([0, T]; G_{div})$ such that $h_n \rightarrow h$ in $L^2(0, T; V'_{div})$. It is easy to see that this approximating problem is equivalent to solve a Cauchy problem for a system of ODEs in the $2n$ unknowns $\alpha_i^{(n)}, \beta_i^{(n)}$ ($\gamma_i^{(n)}$ can be deduced from (3.16)). Since F'' and m are locally Lipschitz on \mathbb{R} , the Cauchy-Lipschitz theorem ensures that there exists $T_n^* \in (0, +\infty]$ such that system (3.14)-(3.17) has a unique maximal solution $\mathbf{a}^{(n)} := (\alpha_1^{(n)}, \dots, \alpha_n^{(n)})$, $\mathbf{b}^{(n)} := (\beta_1^{(n)}, \dots, \beta_n^{(n)})$ on $[0, T_n^*)$ with $\mathbf{a}^{(n)}, \mathbf{b}^{(n)} \in C^1([0, T_n^*]; \mathbb{R}^n)$.

By taking $\psi = \mu_n$ and $v = u_n$ in (3.14) and (3.15), respectively, and adding the resulting identities together, we get

$$\frac{d}{dt} \mathcal{E}(u_n, \varphi_n) + \nu \|\nabla u_n\|^2 + \|\sqrt{m(\varphi_n)} \nabla \mu_n\|^2 = (h_n, u_n),$$

where \mathcal{E} is defined as in (3.8). Integrating this identity between 0 and t

$$\mathcal{E}(u_n(t), \varphi_n(t)) + \int_0^t \left(\nu \|\nabla u_n\|^2 + \|\sqrt{m(\varphi_n)} \nabla \mu_n\|^2 \right) d\tau = \mathcal{E}(u_{0n}, \varphi_{0n}) + \int_0^t \langle h_n(\tau), u_n \rangle d\tau. \quad (3.18)$$

Observe now that

$$\int_0^t \langle h_n(\tau), u_n \rangle d\tau \leq \frac{\nu}{2} \int_0^t \|\nabla u_n\|^2 d\tau + \frac{1}{2\nu} \int_0^t \|h_n\|_{V'_{div}}^2 d\tau.$$

On the other hand, taking (H2) and (H4) into account, we get

$$\mathcal{E}(u_n, \varphi_n) = \frac{1}{2} \|u_n\|^2 + \frac{1}{2} \|\sqrt{a} \varphi_n\|^2 - \frac{1}{2} (\varphi_n, J * \varphi_n) + \int_{\Omega} F(\varphi_n)$$

$$\begin{aligned}
&\geq \frac{1}{2}\|u_n\|^2 + \frac{1}{2} \int_{\Omega} a\varphi_n^2 - \frac{1}{2}\|\varphi_n\| \|J * \varphi_n\| + \int_{\Omega} (c_1\varphi_n^2 - c_2) \\
&\geq \frac{1}{2}\|u_n\|^2 + c'_1\|\varphi_n\|^2 - c'_2,
\end{aligned}$$

where $c'_1 = (a_* - a^*)/2 + c_1 > 0$ and $c'_2 = c_2|\Omega|$. Using also the convergence assumption for $\{u_{0n}\}$, $\{h_n\}$, the fact that $\varphi_{0n} \rightarrow \varphi_0$ in $H^2(\Omega)$ (since $\varphi_0 \in D(B)$) and the lower bound $m_1 > 0$ for the mobility m (cf. (H1)), we first deduce that $T_n^* = +\infty$ for every $n \geq 1$ (notice that $|\mathbf{a}^{(n)}(t)| = \|u_n(t)\|$ and $|\mathbf{b}^{(n)}(t)| = \|\varphi_n(t)\|$) and furthermore we get the following estimates which hold for any given $0 < T < +\infty$

$$\|u_n\|_{L^\infty(0,T;G_{div}) \cap L^2(0,T;V_{div})} \leq C, \quad (3.19)$$

$$\|\varphi_n\|_{L^\infty(0,T;H)} \leq C, \quad (3.20)$$

$$\|F(\varphi_n)\|_{L^\infty(0,T;L^1(\Omega))} \leq C, \quad (3.21)$$

$$\|\nabla\mu_n\|_{L^2(0,T;H)} \leq C, \quad (3.22)$$

with C independent of n . Henceforth we shall denote by C a positive constant which depends at most on $\|u_0\|$, $\|\varphi_0\|$, $\int_{\Omega} F(\varphi_0)$, $\|h\|_{L^2(0,T;V'_{div})}$ and on J , F , ν , m_1 , Ω , but they are independent of n . Instead, c will stand for a generic positive constant depending on the parameters of the problem only, i.e. on J , F , ν , m_1 and Ω , but it is independent of n . The values of both C and c may possibly vary even within the same line.

Let us now recall the estimate

$$\|\nabla\mu_n\|^2 \geq \frac{c_0^2}{4}\|\nabla\varphi_n\|^2 - c\|\varphi_n\|^2, \quad (3.23)$$

which can be deduced as in [14, Proof of Theorem 1] by multiplying μ_n by $-\Delta\varphi_n$ in L^2 and integrating by parts. By means of (3.23), from (3.20) and (3.22) we get

$$\|\varphi_n\|_{L^2(0,T;V)} \leq C. \quad (3.24)$$

Due to (H5), which in particular implies that $|F'(s)| \leq c|F(s)| + c$, we have

$$|\bar{\mu}_n| = \left| \int_{\Omega} F'(\varphi_n) \right| \leq c\|F(\varphi_n)\|_{L^1(\Omega)} + c \leq C,$$

and therefore, thanks to (3.21), (3.22) and to the Poincaré-Wirtinger inequality, we deduce

$$\|\mu_n\|_{L^2(0,T;V)} \leq C. \quad (3.25)$$

In addition, (H5) and (3.21) yield the following control

$$\|F'(\varphi_n)\|_{L^\infty(0,T;L^r(\Omega))} \leq C. \quad (3.26)$$

Let us now derive the estimates for the sequences of time derivatives $\{u'_n\}$ and $\{\varphi'_n\}$. By taking $\psi_I := P_n\psi$, for $\psi \in V$ arbitrary, as test function in (3.14) we obtain

$$\langle \varphi'_n, \psi \rangle = (\varphi'_n, \psi_I) = -(m(\varphi_n)\nabla\mu_n, \nabla\psi_I) + (u_n\varphi_n, \nabla\psi_I).$$

Assume first $d = 3$. We have

$$\begin{aligned} |\langle \varphi'_n, \psi \rangle| &\leq m_2 \|\nabla\mu_n\| \|\nabla\psi_I\| + \|u_n\|_{L^6(\Omega)^3} \|\varphi_n\|_{L^3(\Omega)} \|\nabla\psi_I\| \\ &\leq c(\|\nabla\mu_n\| + \|\nabla u_n\| \|\varphi_n\|^{1/2} \|\varphi_n\|_{L^6(\Omega)}^{1/2}) \|\nabla\psi\| \\ &\leq C(\|\nabla\mu_n\| + \|\nabla u_n\| \|\varphi_n\|_V^{1/2}) \|\nabla\psi\|, \quad \forall \psi \in V \end{aligned} \quad (3.27)$$

where in the last estimate we have used (3.20). Then, by (3.19), (3.22) and (3.24) from (3.27) we get

$$\|\varphi'_n\|_{L^{4/3}(0,T;V')} \leq C. \quad (3.28)$$

For $d = 2$, by arguing as above it is not difficult to infer the following bound

$$\|\varphi'_n\|_{L^{2-\delta}(0,T;V')} \leq C. \quad (3.29)$$

As far as the sequence $\{u'_n\}$ is concerned, by arguing exactly as in [14, Proof of Theorem 1] we can deduce the bounds

$$\|u'_n\|_{L^{4/3}(0,T;V'_{div})} \leq C, \quad \text{if } d = 3, \quad (3.30)$$

$$\|u'_n\|_{L^{2-\gamma}(0,T;V'_{div})} \leq C, \quad \text{if } d = 2. \quad (3.31)$$

From (3.19)–(3.20), (3.24)–(3.26) and (3.28)–(3.31), using compactness results, we obtain for a not relabeled subsequence

$$u_n \rightharpoonup u \quad \text{weakly}^* \text{ in } L^\infty(0, T; G_{div}), \quad \text{weakly in } L^2(0, T; V_{div}), \quad (3.32)$$

$$u_n \rightarrow u \quad \text{strongly in } L^2(0, T; G_{div}), \quad \text{a.e. in } Q_T, \quad (3.33)$$

$$u'_n \rightharpoonup u_t \quad \text{weakly in } L^{4/3}(0, T; V'_{div}), \quad d = 3, \quad (3.34)$$

$$u'_n \rightharpoonup u_t \quad \text{weakly in } L^{2-\gamma}(0, T; V'_{div}), \quad d = 2, \quad (3.35)$$

$$\varphi_n \rightharpoonup \varphi \quad \text{weakly}^* \text{ in } L^\infty(0, T; H), \quad \text{weakly in } L^2(0, T; V), \quad (3.36)$$

$$\varphi_n \rightarrow \varphi \quad \text{strongly in } L^2(0, T; H), \quad \text{a.e. in } Q_T, \quad (3.37)$$

$$\varphi'_n \rightarrow \varphi_t \quad \text{weakly in } L^{2-\delta}(0, T; V'), \quad (3.38)$$

$$F'(\varphi_n) \rightharpoonup F^* \quad \text{weakly}^* \text{ in } L^\infty(0, T; L^r(\Omega)), \quad (3.39)$$

$$\mu_n \rightharpoonup \mu \quad \text{weakly in } L^2(0, T; V). \quad (3.40)$$

where $Q_T := \Omega \times (0, T)$. The pointwise convergence (3.37), the weak * convergence (3.39) and the continuity of F' yield $F^* = F'(\varphi)$. By means of (3.32)–(3.40) we can now pass

to the limit in the approximate problem (3.14)–(3.17) as done in [14, Proof of Theorem 1] and obtain the variational formulation of system (1.5)–(1.10) for functions u , φ and μ . In particular notice that, fixing n arbitrarily, for every $\chi \in C_0^\infty(0, T)$ and every $\psi \in \Psi_n$ we have

$$\int_0^T (m(\varphi_k) \nabla \mu_k, \nabla \psi) \chi d\tau \rightarrow \int_0^T (m(\varphi) \nabla \mu, \nabla \psi) \chi d\tau, \quad \text{as } k \rightarrow \infty,$$

as a consequence of the weak convergence (3.40) and of the convergence

$$m(\varphi_k) \rightarrow m(\varphi) \quad \text{strongly in } L^r(Q_T), \quad \text{for all } r \in [1, \infty), \quad (3.41)$$

ensured by (3.37) and by Lebesgue's theorem.

As far as the energy inequality is concerned, let us observe that (3.40) and (3.41) imply that

$$\sqrt{m(\varphi_n)} \nabla \mu_n \rightharpoonup \sqrt{m(\varphi)} \nabla \mu \quad \text{weakly in } L^s(Q_T), \quad \text{for all } s \in [1, 2). \quad (3.42)$$

But (3.18) yields

$$\|\sqrt{m(\varphi_n)} \nabla \mu_n\|_{L^2(Q_T)} \leq C,$$

and hence (3.42) holds also for $s = 2$. Therefore

$$\int_0^t \|\sqrt{m(\varphi)} \nabla \mu\|^2 d\tau \leq \liminf_{n \rightarrow \infty} \int_0^t \|\sqrt{m(\varphi_n)} \nabla \mu_n\|^2 d\tau.$$

This fact and the above convergences allow us to pass to the inferior limit in (3.18) and deduce (3.7).

Let us now suppose that assumption (H4) is replaced by (H7). Then, the energy of the approximate solution can be controlled from below by

$$\mathcal{E}(u_n, \varphi_n) \geq c(\|u_n\|^2 + \|\varphi_n\|_{L^p(\Omega)}^p) - c.$$

From (3.18) we can then improve (3.20), that is

$$\|\varphi_n\|_{L^\infty(0, T; L^p(\Omega))} \leq C, \quad (3.43)$$

which yields (3.9). The regularity properties (3.10) and (3.11) follow by a comparison argument exactly as in [14, Proof of Corollary 1] with reference to the weak formulation (3.5), (3.6) and taking the improved regularity for φ and (H1) into account.

Furthermore, if $d = 2$ and (H4) is replaced by (H7), we can take μ and u as test functions in the variational formulation (3.5), (3.6) and argue as in [14, Proof of Corollary 2] in order to deduce the energy identity (3.12). As far as the the dissipative estimate

(3.13) is concerned, this can be deduced without difficulties by adapting the argument of [14, Proof of Corollary 2] and taking (H1) into account.

Finally, if $\varphi_0 \in H$ with $F(\varphi_0) \in L^1(\Omega)$, we can approximate φ_0 with $\varphi_{0m} \in D(B)$ given by $\varphi_{0m} := (I + B/m)^{-1}\varphi_0$. This sequence satisfies $\varphi_{0m} \rightarrow \varphi$ in H and by exploiting assumption (H3) and the argument of [14, Proof of Theorem 1] we can easily recover the existence of a weak solution, the energy inequality (3.7) and, for $d = 2$, the energy identity (3.12) as well as the dissipative estimate (3.13). \square

It may be interesting to observe that another energy identity for $d = 2$ (inequality for $d = 3$) is satisfied by the weak solution of Theorem 1. This energy identity will turn to be useful especially in the degenerate case in order to establish the existence of the global attractor in 2D.

Corollary 1. *Let the assumptions of Theorem 1 be satisfied with (H4) replaced by (H7). Then, if $d = 2$ the weak solution $z = [u, \varphi]$ constructed in Theorem 1 satisfies the following energy identity*

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} (\|u\|^2 + \|\varphi\|^2) + \int_{\Omega} m(\varphi) F''(\varphi) |\nabla \varphi|^2 + \int_{\Omega} am(\varphi) |\nabla \varphi|^2 + \nu \|\nabla u\|^2 \\ & = \int_{\Omega} m(\varphi) (\nabla J * \varphi - \varphi \nabla a) \cdot \nabla \varphi + \int_{\Omega} (a\varphi - J * \varphi) u \cdot \nabla \varphi + \langle h, u \rangle, \end{aligned} \quad (3.44)$$

for almost any $t > 0$. Furthermore, if $d = 3$ and if (H7) is satisfied with $p \geq 3$, the weak solution z satisfies the following energy inequality

$$\begin{aligned} & \frac{1}{2} (\|u(t)\|^2 + \|\varphi(t)\|^2) + \int_0^t \int_{\Omega} m(\varphi) F''(\varphi) |\nabla \varphi|^2 + \int_0^t \int_{\Omega} am(\varphi) |\nabla \varphi|^2 \\ & + \nu \int_0^t \|\nabla u\|^2 \leq \frac{1}{2} (\|u_0\|^2 + \|\varphi_0\|^2) + \int_0^t \int_{\Omega} m(\varphi) (\nabla J * \varphi - \varphi \nabla a) \cdot \nabla \varphi \\ & + \int_0^t \int_{\Omega} (a\varphi - J * \varphi) u \cdot \nabla \varphi + \int_0^t \langle h, u \rangle d\tau, \quad \forall t > 0. \end{aligned} \quad (3.45)$$

Proof. If $d = 2$, we can take $\psi = \varphi$ and $v = u$ as test functions in the weak formulation (3.5) and (3.6), respectively. This is allowed due to the regularity properties (3.10) and (3.11). Then, we add the resulting identities and we observe that, since by an Alikakos' iteration technique as in [10, Theorem 2.1] it can be shown that $\varphi \in L^\infty(\Omega)$ and since $F \in C^2(\mathbb{R})$, we can write $\nabla F'(\varphi) = F''(\varphi) \nabla \varphi$. Then, on account of this identity and of (3.4), the second term on the left hand side of (3.5) can be rewritten and (3.44) immediately follows.

Let $d = 3$ and assume that (H7) holds with $p \geq 3$. Then, due to the second of (3.10) we can take $\psi = \varphi$ as test function in (3.5) and get

$$\frac{1}{2} \frac{d}{dt} \|\varphi\|^2 + (m(\varphi) \nabla \mu, \nabla \varphi) = 0.$$

By integrating this identity between 0 and t and by rewriting the term $(m(\varphi)\nabla\mu, \nabla\varphi)$ as done above in the case $d = 2$ we obtain

$$\begin{aligned} & \frac{1}{2}\|\varphi(t)\|^2 + \int_0^t \int_{\Omega} m(\varphi)F''(\varphi)|\nabla\varphi|^2 + \int_0^t \int_{\Omega} am(\varphi)|\nabla\varphi|^2 \\ & + \int_0^t \int_{\Omega} m(\varphi)(\varphi\nabla a - \nabla J * \varphi) \cdot \nabla\varphi = \frac{1}{2}\|\varphi_0\|^2. \end{aligned} \quad (3.46)$$

On the other hand we take $v = u_n$ in the second equation (3.15) of the Faedo-Galerkin approximating problem and integrate the resulting identity to get

$$\begin{aligned} & \frac{1}{2}\|u_n(t)\|^2 + \nu \int_0^t \|\nabla u_n\|^2 d\tau = \frac{1}{2}\|u_{0n}\|^2 - \int_0^t (\varphi_n \nabla \mu_n, u_n) d\tau + \int_0^t \langle h, u_n \rangle d\tau \\ & = \frac{1}{2}\|u_{0n}\|^2 + \int_0^t \int_{\Omega} (a\varphi_n - J * \varphi_n) \nabla \varphi_n \cdot u_n + \int_0^t \langle h, u_n \rangle d\tau. \end{aligned} \quad (3.47)$$

We can now pass to the limit in (3.47) and use the second weak convergence (3.32) and (3.33). In particular, observe that

$$\begin{aligned} & \int_0^t \int_{\Omega} (a\varphi_n - J * \varphi_n) \nabla \varphi_n \cdot u_n = - \int_{Q_t} \left(\frac{1}{2} \varphi_n^2 \nabla a - \varphi_n (\nabla J * \varphi_n) \right) \cdot u_n \\ & \rightarrow - \int_{Q_t} \left(\frac{1}{2} \varphi^2 \nabla a - \varphi (\nabla J * \varphi) \right) \cdot u = \int_0^t \int_{\Omega} (a\varphi - J * \varphi) \nabla \varphi \cdot u, \end{aligned}$$

where this last convergence is a consequence of the fact that, due to (3.32), (3.33), (3.36), (3.37) and interpolation, we have $\varphi_n \rightarrow \varphi$ strongly in $L^3(Q_t)$ and $u_n \rightarrow u$ strongly in $L^3(Q_t)^3$ (recall that $\nabla a \in L^\infty$). Hence we get

$$\frac{1}{2}\|u(t)\|^2 + \nu \int_0^t \|\nabla u\|^2 \leq \frac{1}{2}\|u_0\|^2 + \int_0^t \int_{\Omega} (a\varphi - J * \varphi) u \cdot \nabla \varphi + \int_0^t \langle h, u \rangle d\tau. \quad (3.48)$$

Summing (3.46) and (3.48) we deduce (3.45).

Remark 3. The theorems proven in this section still hold when the viscosity smoothly depends on φ (see [19]). Moreover, they allow us to generalize to the variable (non-degenerate) mobility the results obtained in [19] on the longtime behavior. More precisely, the existence of the global attractor and the existence of a trajectory attractor in the cases $d = 2$ and $d = 3$, respectively.

4 Degenerate mobility

In this section we consider a mobility m which degenerates at ± 1 and we assume that the double-well potential F is singular (e.g. logarithmic like) and defined in $(-1, 1)$. More precisely, we assume that $m \in C^1([-1, 1])$, $m \geq 0$ and that $m(s) = 0$ if and only if $s = -1$ or $s = 1$. Furthermore, we suppose that m and F fulfill the condition

(A1) $F \in C^2(-1, 1)$ and

$$mF'' \in C([-1, 1]).$$

We point out that (A1) is a typical condition which arises in the Cahn-Hilliard equation with degenerate mobility (see [16, 27, 28, 26]).

As far as F is concerned we assume that it can be written in the following form

$$F = F_1 + F_2,$$

where the singular component F_1 and the regular component $F_2 \in C^2([-1, 1])$ satisfy the following assumptions.

(A2) There exist $a_2 > 4(a^* - a_* - b_2)$, where $b_2 := \min_{[-1, 1]} F_2''$, and $\epsilon_0 > 0$ such that

$$F_1''(s) \geq a_2, \quad \forall s \in (-1, -1 + \epsilon_0] \cup [1 - \epsilon_0, 1).$$

(A3) There exists $\epsilon_0 > 0$ such that F_1'' is non-decreasing in $[1 - \epsilon_0, 1)$ and non-increasing in $(-1, -1 + \epsilon_0]$.

(A4) There exists $c_0 > 0$ such that

$$F''(s) + a(x) \geq c_0, \quad \forall s \in (-1, 1), \quad \text{a.a. } x \in \Omega. \quad (4.1)$$

The constants a^* and a_* are given in (H2) and (H4), respectively, and the assumption on the external force h is still (H6).

Remark 4. It is easy to see that (A1)–(A4) are satisfied in the physically relevant case where the mobility and the double-well potential are given by

$$m(s) = k_1(1 - s^2), \quad F(s) = -\frac{\theta_c}{2}s^2 + \frac{\theta}{2}((1 + s)\log(1 + s) + (1 - s)\log(1 - s)), \quad (4.2)$$

where $0 < \theta < \theta_c$. Indeed, setting $F_1(s) := (\theta/2)((1 + s)\log(1 + s) + (1 - s)\log(1 - s))$ and $F_2(s) = -(\theta_c/2)s^2$, then we have $mF_1'' = k_1\theta > 0$ and so (A1) is fulfilled. Moreover F_1 satisfies also (A2) and (A3), while (A4) holds if and only if $\inf_{\Omega} a > \theta_c - \theta$. Another example is given by taking

$$m(s) = k(s)(1 - s^2)^m, \quad F(s) = -k_2s^2 + F_1(s)$$

where $k \in C^1([-1, 1])$ such that $0 < k_3 \leq k(s) \leq k_4$ for all $s \in [-1, 1]$, and F_1 is a $C^2(-1, 1)$ convex function such that

$$F_1''(s) = l(s)(1 - s^2)^{-m}, \quad \forall s \in (-1, 1),$$

where $m \geq 1$ and $l \in C^1([-1, 1])$.

Remark 5. Note that (A4), which is equivalent to the condition $\inf_{\Omega} a > -\inf_{(-1,1)} F''$, is more general than [20, (A6)]. More precisely, [20, (A6)] implies (A4), while (A4) implies [20, (A6)] provided we have $\inf_{(-1,1)}(F_1'' + F_2'') = \inf_{(-1,1)} F_1'' + \min_{[-1,1]} F_2''$. For example, consider the following double-well potential

$$F(s) = -\frac{\theta_c}{2}s^2 - \frac{\theta_2}{12}s^4 + \frac{\theta}{2}((1+s)\log(1+s) + (1-s)\log(1-s)),$$

where $0 < \theta < \theta_c$ and $0 < \theta < \theta_2$. Then, it is easy to see that [20, (A6)] is satisfied iff $\inf_{\Omega} a > \theta_c + \theta_2 - \theta$, while (A4) requires the weaker condition $\inf_{\Omega} a > \theta_c + \theta_2 - 2\sqrt{\theta\theta_2}$.

If the mobility degenerates we are no longer able to control the gradient of the chemical potential μ in some L^p space. For this reason, and also in order to pass to the limit in the approximate problem considered in the proof of the existence result, we shall have to suitably reformulate the definition of weak solution in such a way that μ does not appear anymore (cf. [16]).

Definition 2. Let $u_0 \in G_{div}$, $\varphi_0 \in H$ with $F(\varphi_0) \in L^1(\Omega)$ and $0 < T < +\infty$ be given. A couple $[u, \varphi]$ is a weak solution to (1.5)-(1.10) on $[0, T]$ corresponding to $[u_0, \varphi_0]$ if

- u, φ satisfy

$$u \in L^\infty(0, T; G_{div}) \cap L^2(0, T; V_{div}), \quad (4.3)$$

$$u_t \in L^{4/3}(0, T; V'_{div}), \quad \text{if } d = 3, \quad (4.4)$$

$$u_t \in L^2(0, T; V'_{div}), \quad \text{if } d = 2, \quad (4.5)$$

$$\varphi \in L^\infty(0, T; H) \cap L^2(0, T; V), \quad (4.6)$$

$$\varphi_t \in L^2(0, T; V'), \quad (4.7)$$

and

$$\varphi \in L^\infty(Q_T), \quad |\varphi(x, t)| \leq 1 \quad \text{a.e. } (x, t) \in Q_T := \Omega \times (0, T); \quad (4.8)$$

- for every $\psi \in V$, every $v \in V_{div}$ and for almost any $t \in (0, T)$ we have

$$\begin{aligned} & \langle \varphi_t, \psi \rangle + \int_{\Omega} m(\varphi) F''(\varphi) \nabla \varphi \cdot \nabla \psi + \int_{\Omega} m(\varphi) a \nabla \varphi \cdot \nabla \psi \\ & + \int_{\Omega} m(\varphi) (\varphi \nabla a - \nabla J * \varphi) \cdot \nabla \psi = (u \varphi, \nabla \psi), \end{aligned} \quad (4.9)$$

$$\langle u_t, v \rangle + \nu(\nabla u, \nabla v) + b(u, u, v) = ((a\varphi - J * \varphi) \nabla \varphi, v) + \langle h, v \rangle; \quad (4.10)$$

- the initial conditions $u(0) = u_0$, $\varphi(0) = \varphi_0$ hold (cf. Remark 2).

We now state the existence result for the degenerate mobility case. To this aim we need to introduce the *entropy* function $M \in C^2(-1, 1)$ defined by

$$m(s)M''(s) = 1, \quad M(0) = M'(0) = 0.$$

Theorem 2. *Assume that (A1)-(A4) and (H2), (H6) are satisfied. Let $u_0 \in G_{div}$, $\varphi_0 \in L^\infty(\Omega)$ such that $F(\varphi_0) \in L^1(\Omega)$ and $M(\varphi_0) \in L^1(\Omega)$. Then, for every $T > 0$ there exists a weak solution $z := [u, \varphi]$ to (1.5)-(1.10) on $[0, T]$ corresponding to $[u_0, \varphi_0]$ in the sense of Definition 2 such that $\overline{\varphi}(t) = \overline{\varphi_0}$ for all $t \in [0, T]$ and*

$$\varphi \in L^\infty(0, T; L^p(\Omega)), \quad (4.11)$$

where $p \leq 6$ for $d = 3$ and $2 \leq p < \infty$ for $d = 2$. In addition, if $d = 2$, the weak solution $z := [u, \varphi]$ satisfies the energy equation (3.44), and, if $d = 3$, z satisfies the energy inequality (3.45).

Remark 6. The potential F and the entropy M are not independent. Indeed, assumption (A1) implies that there exists a constant $\gamma_0 > 0$ such that $|m(s)F''(s)| \leq \gamma_0$, for all $s \in [-1, 1]$. Combining this estimate with the definition of M we get $|F''(s)| \leq \gamma_0 M''(s)$, for all $s \in (-1, 1)$. Thus we get

$$|F(s)| \leq |F(0)| + |F'(0)||s| + \gamma_0 M(s), \quad \forall s \in (-1, 1). \quad (4.12)$$

Therefore, in the statement of Theorem 2, condition $F(\varphi_0) \in L^1(\Omega)$ is actually a consequence of the entropy assumption $M(\varphi_0) \in L^1(\Omega)$.

Proof. Let us consider the approximate problem P_ϵ : find a weak solution $[u_\epsilon, \varphi_\epsilon]$ to

$$\varphi'_\epsilon + u_\epsilon \cdot \nabla \varphi_\epsilon = \operatorname{div}(m_\epsilon(\varphi_\epsilon) \nabla \mu_\epsilon), \quad (4.13)$$

$$u'_\epsilon - \nu \Delta u_\epsilon + (u_\epsilon \cdot \nabla) u_\epsilon + \nabla \pi_\epsilon = \mu_\epsilon \nabla \varphi_\epsilon + h, \quad (4.14)$$

$$\mu_\epsilon = a\varphi_\epsilon - J * \varphi_\epsilon + F'_\epsilon(\varphi_\epsilon), \quad (4.15)$$

$$\operatorname{div}(u_\epsilon) = 0, \quad (4.16)$$

$$\frac{\partial \mu_\epsilon}{\partial n} = 0, \quad u_\epsilon = 0, \quad \text{on } \partial\Omega, \quad (4.17)$$

$$u_\epsilon(0) = u_0, \quad \varphi_\epsilon(0) = \varphi_0, \quad \text{in } \Omega. \quad (4.18)$$

Problem P_ϵ is obtained from (1.5)-(1.10) by replacing the singular potential F with a smooth potential F_ϵ and the degenerate mobility m with a non-degenerate one m_ϵ . In particular, F_ϵ is represented by

$$F_\epsilon = F_{1\epsilon} + F_{2\epsilon},$$

where $F_{1\epsilon}$ and $F_{2\epsilon}$ are defined by

$$F_{1\epsilon}''(s) = \begin{cases} F_1''(1 - \epsilon), & s \geq 1 - \epsilon \\ F_1''(s), & |s| \leq 1 - \epsilon \\ F_1''(-1 + \epsilon), & s \leq -1 + \epsilon \end{cases} \quad (4.19)$$

$$F_{2\epsilon}''(s) = \begin{cases} F_2''(1 - \epsilon), & s \geq 1 - \epsilon \\ F_2''(s), & |s| \leq 1 - \epsilon \\ F_2''(-1 + \epsilon), & s \leq -1 + \epsilon, \end{cases} \quad (4.20)$$

with $F_{1\epsilon}(0) = F_1(0)$, $F_{1\epsilon}'(0) = F_1'(0)$, $F_{2\epsilon}(0) = F_2(0)$, $F_{2\epsilon}'(0) = F_2'(0)$.

The approximate non-degenerate mobility is given by

$$m_\epsilon(s) = \begin{cases} m(1 - \epsilon), & s \geq 1 - \epsilon \\ m(s), & |s| \leq 1 - \epsilon \\ m(-1 + \epsilon), & s \leq -1 + \epsilon. \end{cases} \quad (4.21)$$

Assumption (A3) implies that

$$F_{1\epsilon}(s) \leq F_1(s), \quad \forall s \in (-1, 1), \quad \forall \epsilon \in (0, \epsilon_0]. \quad (4.22)$$

On the other hand, from the definition of $F_{2\epsilon}$ we have, for $s > 1 - \epsilon$,

$$F_{2\epsilon}(s) = F_2(1 - \epsilon) + F_2'(1 - \epsilon)(s - (1 - \epsilon)) + \frac{1}{2}F_2''(1 - \epsilon)(s - (1 - \epsilon))^2, \quad (4.23)$$

(a similar expression holds for $s < -1 + \epsilon$) and $F_{2\epsilon}(s) = F_2(s)$ for $|s| \leq 1 - \epsilon$. Since $F_2 \in C^2([-1, 1])$, then we deduce that there exist two positive constants L_1, L_2 , which are independent of ϵ , such that

$$|F_{2\epsilon}(s)| \leq L_1 s^2 + L_2, \quad \forall s \in \mathbb{R}. \quad (4.24)$$

Therefore, by using the assumption on the initial datum φ_0 and (4.22), (4.24) we have

$$\int_{\Omega} F_\epsilon(\varphi_0) \leq \int_{\Omega} F_1(\varphi_0) + L_1 \|\varphi_0\|^2 + L_2 < \infty, \quad \forall \epsilon \in (0, \epsilon_0]. \quad (4.25)$$

Furthermore, we can see that by assumption (A2) there is a $\delta_0 \in \mathbb{R}$ such that

$$F_\epsilon(s) \geq \delta s^2 - \delta_0, \quad \forall s \in \mathbb{R}, \quad \forall \epsilon \in (0, \epsilon_0], \quad (4.26)$$

where $0 < \delta < a_2/8 + b_2/2$. Indeed, (A2) implies that there exist $a_1^+, a_1^-, a_0 \in \mathbb{R}$ such that $F_1'(s) \geq a_1^+$ for all $s \in [1 - \epsilon_0, 1)$, $F_1'(s) \leq a_1^-$ for all $s \in (-1, -1 + \epsilon_0]$, and $F_1(s) \geq a_0$ for all $s \in (-1, 1)$. Hence, by using definition (4.19), for $s \geq 1 - \epsilon$ we get

$$F_{1\epsilon}(s) = F_1(1 - \epsilon) + F_1'(1 - \epsilon)(s - (1 - \epsilon)) + \frac{1}{2}F_1''(1 - \epsilon)(s - (1 - \epsilon))^2$$

$$\geq a'_0 + \frac{a_2}{4}(s - (1 - \epsilon))^2 \geq \frac{a_2}{8}s^2 + a'_0 - \frac{a_2}{2}, \quad (4.27)$$

provided that $\epsilon \in (0, \epsilon_0]$. For $|s| \leq 1 - \epsilon$ we have $F_{1\epsilon}(s) = F_1(s) \geq a_0 \geq (a_2/8)s^2 - a_2/8 + a_0$. For $s \leq -1 + \epsilon$ we can argue as in (4.27). On the other hand, from (4.23) we have

$$F_{2\epsilon}(s) \geq \left(\delta - \frac{a_2}{8}\right)s^2 - \delta_1, \quad \forall s \in \mathbb{R}, \quad \forall \epsilon \in (0, 1), \quad (4.28)$$

where δ is taken as above and δ_1 is a nonnegative constant (depending on $b_0 := \min_{[-1,1]} F_2$, $b_1 := \min_{[-1,1]} F'_2$ and b_2). Combining (4.27) with (4.28) we deduce (4.26).

Moreover, (A4) immediately implies that, for $\epsilon \in (0, \epsilon_0]$, there holds

$$F'_\epsilon(s) + a(x) \geq c_0, \quad \forall s \in \mathbb{R}, \quad \text{a.e. } x \in \Omega. \quad (4.29)$$

Indeed, recall that for $s \geq \pm 1 \mp \epsilon$ we have $F''_{1\epsilon}(s) = F''_1(\pm 1 \mp \epsilon)$, $F''_{2\epsilon}(s) = F''_2(\pm 1 \mp \epsilon)$, and for $|s| \leq 1 - \epsilon$ we have $F''_{1\epsilon}(s) = F''_1(s)$, $F''_{2\epsilon}(s) = F''_2(s)$.

We can now check that assumptions (H1)–(H6) of Theorem 1 for Problem P_ϵ (with a fixed $\epsilon \in (0, \epsilon_0]$) are satisfied. In particular, due to (A2), we can choose δ such that $(a^* - a_*)/2 < \delta < a_2/8 + b_2/2$ and so (H4) is ensured by (4.26), while (H5) is satisfied with $r = 2$. Theorem 1 therefore entails that Problem P_ϵ admits a weak solution $z_\epsilon = [u_\epsilon, \varphi_\epsilon]$ satisfying (3.1)–(3.3) and the energy inequality

$$\mathcal{E}_\epsilon(z_\epsilon(t)) + \int_0^t \left(\nu \|\nabla u_\epsilon\|^2 + \|\sqrt{m_\epsilon(\varphi_\epsilon)} \nabla \mu_\epsilon\|^2 \right) d\tau \leq \mathcal{E}_\epsilon(z_0) + \int_0^t \langle h(\tau), u_\epsilon(\tau) \rangle d\tau, \quad (4.30)$$

for every $t > 0$, where

$$\mathcal{E}_\epsilon(z_\epsilon) := \frac{1}{2} \|u_\epsilon\|^2 + \frac{1}{2} \|\sqrt{a} \varphi_\epsilon\|^2 - \frac{1}{2} (\varphi_\epsilon, J * \varphi_\epsilon) + \int_\Omega F_\epsilon(\varphi_\epsilon).$$

Now, using (4.25) and (4.26), from (4.30) we immediately obtain the following uniform (with respect to $\epsilon \in (0, \epsilon_0]$) estimates

$$\|u_\epsilon\|_{L^\infty(0,T;G_{div}) \cap L^2(0,T;V_{div})} \leq C, \quad (4.31)$$

$$\|\varphi_\epsilon\|_{L^\infty(0,T;H)} \leq C, \quad (4.32)$$

$$\|\sqrt{m_\epsilon(\varphi_\epsilon)} \nabla \mu_\epsilon\|_{L^2(0,T;H)} \leq C, \quad (4.33)$$

where the positive constant C can possibly change from line to line but is always independent of ϵ . Now, test (4.13) by $\psi = M'_\epsilon(\varphi_\epsilon)$, where M_ϵ is a C^2 function such that $m_\epsilon(s)M''_\epsilon(s) = 1$ and $M_\epsilon(0) = M'_\epsilon(0) = 0$. We get

$$\frac{d}{dt} \int_\Omega M_\epsilon(\varphi_\epsilon) + \int_\Omega m_\epsilon(\varphi_\epsilon) \nabla \mu_\epsilon \cdot M''_\epsilon(\varphi_\epsilon) \nabla \varphi_\epsilon = \int_\Omega u_\epsilon \varphi_\epsilon \cdot M''_\epsilon(\varphi_\epsilon) \nabla \varphi_\epsilon = 0, \quad (4.34)$$

where the last identity is due to (4.16). Therefore (4.34) yields

$$\frac{d}{dt} \int_{\Omega} M_{\epsilon}(\varphi_{\epsilon}) + \int_{\Omega} \nabla \mu_{\epsilon} \cdot \nabla \varphi_{\epsilon} = 0.$$

On account of (4.15), we obtain

$$\frac{d}{dt} \int_{\Omega} M_{\epsilon}(\varphi_{\epsilon}) + \int_{\Omega} \left((a + F_{\epsilon}''(\varphi_{\epsilon})) |\nabla \varphi_{\epsilon}|^2 + \varphi_{\epsilon} \nabla a \cdot \nabla \varphi_{\epsilon} - \nabla J * \varphi_{\epsilon} \cdot \nabla \varphi_{\epsilon} \right) = 0. \quad (4.35)$$

By using (4.29) and (4.32), from (4.35) we hence infer

$$\frac{d}{dt} \int_{\Omega} M_{\epsilon}(\varphi_{\epsilon}) + \frac{c_0}{2} \|\nabla \varphi_{\epsilon}\|^2 \leq c_J \|\varphi_{\epsilon}\|^2 \leq C,$$

where c_J is a positive constant depending on J only. Therefore, on account of the fact that for ϵ small enough we have $M_{\epsilon}(s) \leq M(s)$ for all $s \in (-1, 1)$, and recalling that $\int_{\Omega} M(\varphi_0) < \infty$, we deduce the bounds (see also (4.32))

$$\|\varphi_{\epsilon}\|_{L^2(0,T;V)} \leq C, \quad (4.36)$$

$$\|M_{\epsilon}(\varphi_{\epsilon})\|_{L^{\infty}(0,T;L^1(\Omega))} \leq C. \quad (4.37)$$

Let us now establish the estimates for φ'_{ϵ} and u'_{ϵ} . Let us first consider the case $d = 3$ and start from φ'_{ϵ} . From the variational formulation of (4.13) we get

$$\begin{aligned} |\langle \varphi'_{\epsilon}, \psi \rangle| &\leq \|m_{\epsilon}(\varphi_{\epsilon}) \nabla \mu_{\epsilon}\| \|\nabla \psi\| + \|u_{\epsilon}\|_{L^6(\Omega)} \|\varphi_{\epsilon}\|_{L^3(\Omega)} \|\nabla \psi\| \\ &\leq c \|\sqrt{m_{\epsilon}(\varphi_{\epsilon})} \nabla \mu_{\epsilon}\| \|\nabla \psi\| + c \|\nabla u_{\epsilon}\| \|\varphi_{\epsilon}\|^{1/2} \|\varphi_{\epsilon}\|_{L^6(\Omega)}^{1/2} \|\nabla \psi\| \\ &\leq C (\|\sqrt{m_{\epsilon}(\varphi_{\epsilon})} \nabla \mu_{\epsilon}\| + \|\nabla u_{\epsilon}\| \|\varphi_{\epsilon}\|_V^{1/2}) \|\nabla \psi\|, \quad \forall \psi \in V, \end{aligned} \quad (4.38)$$

where we have used (4.32). Then, on account of (4.31), (4.33) and (4.36), from (4.38) we obtain

$$\|\varphi'_{\epsilon}\|_{L^{4/3}(0,T;V')} \leq C. \quad (4.39)$$

In order to deduce a bound for u'_{ϵ} , observe first that we have

$$\langle \mu_{\epsilon} \nabla \varphi_{\epsilon}, v \rangle = \langle (a \varphi_{\epsilon} - J * \varphi_{\epsilon}) \nabla \varphi_{\epsilon}, v \rangle, \quad \forall v \in V_{div}.$$

Thus, by (4.32), we have that

$$\begin{aligned} |\langle \mu_{\epsilon} \nabla \varphi_{\epsilon}, v \rangle| &\leq \|a \varphi_{\epsilon} - J * \varphi_{\epsilon}\|_{L^3(\Omega)} \|\nabla \varphi_{\epsilon}\| \|v\|_{L^6(\Omega)} \\ &\leq c_J \|\varphi_{\epsilon}\|_{L^3(\Omega)} \|\varphi_{\epsilon}\|_V \|v\|_{V_{div}} \leq c_J \|\varphi_{\epsilon}\|^{1/2} \|\varphi_{\epsilon}\|_{L^6(\Omega)}^{1/2} \|\varphi_{\epsilon}\|_V \|v\|_{V_{div}} \\ &\leq C \|\varphi_{\epsilon}\|_V^{3/2} \|v\|_{V_{div}}, \end{aligned} \quad (4.40)$$

Hence, (4.36) entails the estimate for the Korteweg force in (4.14)

$$\|\mu_\epsilon \nabla \varphi_\epsilon\|_{L^{4/3}(0,T;V'_{div})} \leq C.$$

The estimates for the other terms in the variational formulation of (4.14) can be deduced by standard arguments for the 3D Navier-Stokes and finally we obtain

$$\|u'_\epsilon\|_{L^{4/3}(0,T;V'_{div})} \leq C. \quad (4.41)$$

In the case $d = 2$, by arguing as in (4.38) and (4.40) and again using (4.31)–(4.33) and (4.36), it not difficult to see that the following estimates hold

$$\|u'_\epsilon\|_{L^{2-\gamma}(0,T;V'_{div})} \leq C, \quad \|\varphi'_\epsilon\|_{L^{2-\delta}(0,T;V')} \leq C, \quad \forall \gamma, \delta \in (0, 1). \quad (4.42)$$

From the estimates above and using standard compactness results we deduce that there exist $u \in L^\infty(0, T; G_{div}) \cap L^2(0, T; V_{div})$ and $\varphi \in L^\infty(0, T; H) \cap L^2(0, T; V)$ such that, up to a (not relabeled) subsequence we have

$$u_\epsilon \rightharpoonup u \quad \text{weakly}^* \text{ in } L^\infty(0, T; G_{div}), \quad \text{weakly in } L^2(0, T; V_{div}), \quad (4.43)$$

$$u_\epsilon \rightarrow u \quad \text{strongly in } L^2(0, T; G_{div}), \quad \text{a.e. in } Q_T, \quad (4.44)$$

$$u'_\epsilon \rightharpoonup u_t \quad \text{weakly in } L^{4/3}(0, T; V'_{div}), \quad \text{if } d = 3, \quad (4.45)$$

$$u'_\epsilon \rightharpoonup u_t \quad \text{weakly in } L^{2-\gamma}(0, T; V'_{div}), \quad \gamma \in (0, 1), \quad \text{if } d = 2, \quad (4.46)$$

$$\varphi_\epsilon \rightharpoonup \varphi \quad \text{weakly}^* \text{ in } L^\infty(0, T; H), \quad \text{weakly in } L^2(0, T; V), \quad (4.47)$$

$$\varphi_\epsilon \rightarrow \varphi \quad \text{strongly in } L^2(0, T; H), \quad \text{a.e. in } Q_T, \quad (4.48)$$

$$\varphi'_\epsilon \rightarrow \varphi_t \quad \text{weakly in } L^{4/3}(0, T; V'), \quad \text{if } d = 3, \quad (4.49)$$

$$\varphi'_\epsilon \rightarrow \varphi_t \quad \text{weakly in } L^{2-\delta}(0, T; V'), \quad \delta \in (0, 1), \quad \text{if } d = 2. \quad (4.50)$$

Let us now show that $|\varphi| \leq 1$ almost everywhere in Q_T . In order to do that we can argue as in [16, Proof of Theorem 1] (see also [12, Proof of Theorem 2.3]). More precisely, in [16, Proof of Theorem 1] the following estimates are established

$$M_\epsilon(s) \geq \frac{1}{2m(1-\epsilon)}(s-1)^2, \quad \forall s > 1,$$

$$M_\epsilon(s) \geq \frac{1}{2m(-1+\epsilon)}(s+1)^2, \quad \forall s < -1.$$

Therefore

$$\begin{aligned} \int_\Omega (|\varphi_\epsilon| - 1)_+^2 &= \int_{\{\varphi_\epsilon > 1\}} (\varphi_\epsilon - 1)^2 + \int_{\{\varphi_\epsilon < -1\}} (\varphi_\epsilon + 1)^2 \\ &\leq 2m(1-\epsilon) \int_{\{\varphi_\epsilon > 1\}} M_\epsilon(\varphi_\epsilon) + 2m(-1+\epsilon) \int_{\{\varphi_\epsilon < -1\}} M_\epsilon(\varphi_\epsilon) \end{aligned}$$

$$\leq 2 \max(m(1 - \epsilon), m(-1 + \epsilon)) \int_{\Omega} M_{\epsilon}(\varphi_{\epsilon}).$$

Now, by using (4.37), the fact that $m(\pm 1 \mp \epsilon) \rightarrow 0$ as $\epsilon \rightarrow 0$ and the generalized Lebesgue theorem, we obtain

$$\int_{\Omega} (|\varphi| - 1)_+^2 = 0 \quad \text{for a.a. } t \in (0, T),$$

so that

$$|\varphi(x, t)| \leq 1, \quad \text{a.e. in } \Omega, \quad \text{a.a. } t \in (0, T). \quad (4.51)$$

We now have to pass to the limit in the variational formulation of the approximate problem (4.13)–(4.18) in order to show that $[u, \varphi]$ is a weak solution to (1.5)–(1.10) according to Definition 2. It is easy that the variational formulation of (4.13)–(4.18) gives

$$\begin{aligned} \langle \varphi'_{\epsilon}, \psi \rangle + \int_{\Omega} m_{\epsilon}(\varphi_{\epsilon}) F'_{\epsilon}(\varphi_{\epsilon}) \nabla \varphi_{\epsilon} \cdot \nabla \psi + \int_{\Omega} m_{\epsilon}(\varphi_{\epsilon}) a \nabla \varphi_{\epsilon} \cdot \nabla \psi \\ + \int_{\Omega} m_{\epsilon}(\varphi_{\epsilon}) (\varphi_{\epsilon} \nabla a - \nabla J * \varphi_{\epsilon}) \cdot \nabla \psi = (u_{\epsilon} \varphi_{\epsilon}, \nabla \psi), \end{aligned} \quad (4.52)$$

$$\langle u'_{\epsilon}, v \rangle + \nu (\nabla u_{\epsilon}, \nabla v) + b(u_{\epsilon}, u_{\epsilon}, v) = ((a \varphi_{\epsilon} - J * \varphi_{\epsilon}) \nabla \varphi_{\epsilon}, v) + \langle h, v \rangle, \quad (4.53)$$

for every $\psi \in V$, every $v \in V_{div}$ and for almost any $t \in (0, T)$.

On account of (A1) we have $|m(s)F''(s)| \leq a$ for every $s \in [-1, 1]$ and for some positive constant a . This immediately implies that

$$|m_{\epsilon}(s)F''_{\epsilon}(s)| \leq a, \quad \forall s \in \mathbb{R}, \quad \forall \epsilon \in (0, 1). \quad (4.54)$$

Furthermore, by using the continuity of mF'' on $[-1, 1]$ (cf. (A1)) and (4.48), it is not difficult to see that

$$m_{\epsilon}(\varphi_{\epsilon})F''_{\epsilon}(\varphi_{\epsilon}) \rightarrow m(\varphi)F''(\varphi), \quad \text{a.e. in } Q_T.$$

Therefore, we obtain

$$m_{\epsilon}(\varphi_{\epsilon})F''_{\epsilon}(\varphi_{\epsilon}) \rightarrow m(\varphi)F''(\varphi), \quad \text{strongly in } L^r(Q_T), \quad \forall r \in [2, \infty). \quad (4.55)$$

Notice also that (4.47), the continuous embedding $L^{\infty}(0, T; L^2(\Omega)) \cap L^2(0, T; L^6(\Omega)) \hookrightarrow L^{10/3}(Q_T)$ and (4.48) imply that, for $d = 3$, we have

$$\varphi_{\epsilon} \rightarrow \varphi \quad \text{strongly in } L^s(Q_T), \quad \text{for all } 2 \leq s < 10/3. \quad (4.56)$$

For $d = 2$, by the Gagliardo-Nirenberg inequality the continuous embedding $L^{\infty}(0, T; L^2(\Omega)) \cap L^2(0, T; V) \hookrightarrow L^4(Q_T)$ holds, thus we get

$$\varphi_{\epsilon} \rightarrow \varphi \quad \text{strongly in } L^s(Q_T), \quad \text{for all } 2 \leq s < 4. \quad (4.57)$$

Furthermore, by Lebesgue's theorem we also deduce

$$m_\epsilon(\varphi_\epsilon) \rightarrow m(\varphi) \quad \text{strongly in } L^s(Q_T), \quad \text{for all } 2 \leq s < \infty. \quad (4.58)$$

We can now multiply (4.52), (4.53) by $\chi, \omega \in C_0^\infty(0, T)$, respectively, and integrate the resulting identities with respect to time from 0 and T . The convergences (4.43)–(4.50) and (4.55)–(4.58) are enough to pass to the limit and to deduce that u and φ satisfy the variational formulation (4.9) and (4.10). Observe that in passing to the limit we can initially take ψ in $C^1(\overline{\Omega})$ and then prove by density that (4.9) holds also for all ψ in V .

In order to prove that $z := [u, \varphi]$ is a weak solution according to Definition 2, we are only left to show that (4.5) and (4.7) hold. To this aim let us first see that φ satisfies (4.11). Indeed, this is a consequence of (4.51) and of the fact that, since $\varphi \in L^2(0, T; V)$, φ is measurable with values in $L^6(\Omega)$ when $d = 3$, and measurable with values in $L^p(\Omega)$, for all $p \in [2, \infty)$, when $d = 2$. Recalling the improved regularity (4.11), and also on account of (4.51), it is now immediate to get (4.5) and (4.7) by a comparison argument in the weak formulation (4.9), (4.10). In particular, notice that the contribution from the convective term in (4.9) can now be estimated as follows

$$|(u\varphi, \nabla\psi)| \leq \|u\|_{L^3(\Omega)} \|\varphi\|_{L^6(\Omega)} \|\nabla\psi\| \leq C \|\nabla u\| \|\nabla\psi\|, \quad \forall \psi \in V.$$

Finally, if $d = 2$, the energy identity (3.44) can be obtained by taking $\psi = \varphi$ and $v = u$ (see (4.7) and (4.5)) as test functions in the weak formulation (4.9) and (4.10), respectively, and adding the resulting identities. In the case $d = 3$, we choose first $\psi = \varphi$ as test function in (4.9) (cf. (4.7)) and integrate the resulting identity to get (3.46). Then we observe that, due to the last part of the proof of Corollary 1, the approximate solution $z_\epsilon := [u_\epsilon, \varphi_\epsilon]$ satisfies

$$\frac{1}{2} \|u_\epsilon(t)\|^2 + \nu \int_0^t \|\nabla u_\epsilon\|^2 \leq \frac{1}{2} \|u_0\|^2 + \int_0^t \int_\Omega (a\varphi_\epsilon - J * \varphi_\epsilon) u_\epsilon \cdot \nabla \varphi_\epsilon + \int_0^t \langle h, u_\epsilon \rangle d\tau.$$

Passing to the limit as $\epsilon \rightarrow 0$ in this last inequality and arguing as at the end of the proof of Corollary 1, taking advantage of (4.43), (4.44), (4.47) and (4.48), we get (3.48). The energy inequality (3.45) is then obtained by adding together (3.46) and (3.48).

It is worth noting that if the mobility degenerates slightly stronger, then the weak solution of Theorem 2 satisfies a physically more relevant energy inequality with respect to the energy identity (3.44). This is proven in the following

Corollary 2. *Let all assumptions of Theorem 2 be satisfied and $d = 2, 3$. In addition, assume that $m(\pm 1) = 0$ with order ≥ 2 . Then the weak solution $z = [u, \varphi]$ given by Theorem 2 also fulfills the following integral inequality*

$$\mathcal{E}(z(t)) + \int_0^t \left(\nu \|\nabla u\|^2 + \left\| \frac{\mathcal{J}}{\sqrt{m(\varphi)}} \right\|^2 \right) d\tau \leq \mathcal{E}(z_0) + \int_0^t \langle h, u \rangle d\tau, \quad (4.59)$$

for all $t > 0$, where the mass flux \mathcal{J} given by

$$\mathcal{J} = -m(\varphi)\nabla(a\varphi - J * \varphi) - m(\varphi)F''(\varphi)\nabla\varphi$$

is such that

$$\mathcal{J} \in L^2(Q_T), \quad \frac{\mathcal{J}}{\sqrt{m(\varphi)}} \in L^2(Q_T).$$

Proof. Setting $\widehat{\mathcal{J}}_\epsilon := -\sqrt{m_\epsilon(\varphi_\epsilon)}\nabla\mu_\epsilon$, from (4.33) we have $\|\widehat{\mathcal{J}}_\epsilon\|_{L^2(Q_T)} \leq C$ and therefore there exists $\widehat{\mathcal{J}} \in L^2(Q_T)$ such that

$$\widehat{\mathcal{J}}_\epsilon \rightharpoonup \widehat{\mathcal{J}} \quad \text{weakly in } L^2(Q_T). \quad (4.60)$$

On the other hand, setting $\mathcal{J}_\epsilon := -m_\epsilon(\varphi_\epsilon)\nabla\mu_\epsilon = \sqrt{m_\epsilon(\varphi_\epsilon)}\widehat{\mathcal{J}}_\epsilon$, we have also $\|\mathcal{J}_\epsilon\|_{L^2(Q_T)} \leq C$. Thus there exists $\mathcal{J} \in L^2(Q_T)$ such that

$$\mathcal{J}_\epsilon \rightharpoonup \mathcal{J} \quad \text{weakly in } L^2(Q_T). \quad (4.61)$$

Since $m_\epsilon(\varphi_\epsilon) \rightarrow m(\varphi)$ strongly in $L^r(Q_T)$ for every $r \in [1, \infty)$, from (4.60) we have $\sqrt{m_\epsilon(\varphi_\epsilon)}\widehat{\mathcal{J}}_\epsilon \rightharpoonup \sqrt{m(\varphi)}\widehat{\mathcal{J}}$ weakly in $L^{2-\gamma}(Q_T)$ for every $\gamma \in (0, 1]$ and therefore by comparison with (4.61) we get

$$\mathcal{J} := \sqrt{m(\varphi)}\widehat{\mathcal{J}}.$$

Let us observe now that inequality (4.30) can be rewritten in the form

$$\mathcal{E}_\epsilon(z_\epsilon(t)) + \int_0^t \left(\nu \|\nabla u_\epsilon\|^2 + \left\| \frac{\mathcal{J}_\epsilon}{\sqrt{m_\epsilon(\varphi_\epsilon)}} \right\|^2 \right) d\tau \leq \mathcal{E}(z_0) + \int_0^t \langle h, u_\epsilon \rangle d\tau, \quad \forall t > 0, \quad (4.62)$$

where \mathcal{J}_ϵ is defined in terms of φ_ϵ , that is,

$$\mathcal{J}_\epsilon = -m_\epsilon(\varphi_\epsilon)\nabla\mu_\epsilon = -m_\epsilon(\varphi_\epsilon)\nabla(a\varphi_\epsilon - J * \varphi_\epsilon) - m_\epsilon(\varphi_\epsilon)F''(\varphi_\epsilon)\nabla\varphi_\epsilon.$$

Using (4.47), (4.55) and (4.58) it is easy to see that

$$\mathcal{J}_\epsilon \rightharpoonup -m(\varphi)\nabla(a\varphi - J * \varphi) - m(\varphi)F''(\varphi)\nabla\varphi \quad \text{weakly in } L^{2-\eta}(Q_T), \quad \forall \eta \in (0, 1).$$

On account of (4.61), we therefore see that \mathcal{J} is given by

$$\mathcal{J} = -m(\varphi)\nabla(a\varphi - J * \varphi) - m(\varphi)F''(\varphi)\nabla\varphi, \quad (4.63)$$

and that the last convergence is also in $L^2(Q_T)$. Hence, taking the inferior limit of the third term on the left hand side of (4.62), we have

$$\int_0^t \left\| \frac{\mathcal{J}}{\sqrt{m(\varphi)}} \right\|^2 d\tau = \int_0^t \|\widehat{\mathcal{J}}\|^2 d\tau \leq \liminf_{\epsilon \rightarrow 0} \int_0^t \|\widehat{\mathcal{J}}_\epsilon\|^2 d\tau = \liminf_{\epsilon \rightarrow 0} \int_0^t \left\| \frac{\mathcal{J}_\epsilon}{\sqrt{m_\epsilon(\varphi_\epsilon)}} \right\|^2 d\tau. \quad (4.64)$$

On the other hand, the strong degeneracy condition $m(\pm 1) = 0$ with order ≥ 2 implies (see Remark 8 here below) that the set $\{x \in \Omega : |\varphi(x, t)| = 1\}$ has zero measure for almost any $t \in (0, T)$. Hence we have $F_\epsilon(\varphi_\epsilon) \rightarrow F(\varphi)$ almost everywhere in Q_T , and Fatou's lemma yields

$$\int_{\Omega} F(\varphi) \leq \liminf_{\epsilon \rightarrow 0} \int_{\Omega} F_\epsilon(\varphi_\epsilon). \quad (4.65)$$

Finally, (4.59) follows immediately from (4.62) on account of (4.64), (4.65), (4.63) and the second weak convergence in (4.43). \square

Remark 7. In comparison with the analogous result for the case of constant mobility (see [20, Theorem 1]), Theorem 2 does not require the condition $|\bar{\varphi}_0| < 1$ (the assumptions on φ_0 imply only that $|\bar{\varphi}_0| \leq 1$). This is essentially due to the fact that here we are dealing with a different weak formulation. Therefore, if F is bounded (e.g. F is given by (4.2)) and at $t = 0$ the fluid is in a pure phase, e.g., $\varphi_0 = 1$ almost everywhere in Ω , and furthermore $u_0 = u(0)$ is given in G_{div} , then we can immediately check that the couple $[u, \varphi]$ given by

$$u = u(\cdot, t), \quad \varphi = \varphi(\cdot, t) = 1, \quad \text{a.e. in } \Omega, \quad \text{a.a. } t \geq 0,$$

where u is solution of the Navier-Stokes equations with non-slip boundary condition, initial velocity field u_0 and external force h , explicitly satisfies the weak formulation (4.9), (4.10). Hence, the nonlocal Cahn-Hilliard-Navier-Stokes model with degenerate mobility and bounded double-well potentials allows pure phases at any time $t \geq 0$. This possibility is excluded in the model with constant mobility since in such model the chemical potential μ (and hence $F'(\varphi)$) appears explicitly.

Remark 8. If $m(\pm 1) = 0$ with order strictly less than 2, then, as a consequence of (A1) and of the definition of the function M , both F and M are bounded in $[-1, 1]$. This can be seen, for instance, by writing M as follows

$$M(s) = \int_0^s \frac{s-t}{m(t)} dt, \quad \forall s \in (-1, 1).$$

In this case, the conditions $F(\varphi_0) \in L^1(\Omega)$ and $M(\varphi_0) \in L^1(\Omega)$ of Theorem 2 are satisfied by every initial datum φ_0 such that $|\varphi_0| \leq 1$ in Ω . Therefore the existence of pure phases is allowed. A relevant example for this case is given by (4.2). On the other hand, if $m(\pm 1) = 0$ with order greater than or equal 2 (in this case we say that m is *strongly degenerate*), then it can be proved (cf. [16, Corollary] and also [12, Theorem 2.3]) that the conditions $F(\varphi_0) \in L^1(\Omega)$ and $M(\varphi_0) \in L^1(\Omega)$ imply that the sets $\{x \in \Omega : \varphi_0(x) = 1\}$ and $\{x \in \Omega : \varphi_0(x) = -1\}$ have both measure zero. Hence, in this case we have obviously

$|\bar{\varphi}_0| < 1$ and furthermore it can be seen that also the sets $\{x \in \Omega : \varphi(x, t) = 1\}$ and $\{x \in \Omega : \varphi(x, t) = -1\}$ have both measure zero for almost any $t > 0$. Therefore, in the case of strongly degenerating mobility the presence of pure phases is not allowed (even on subsets of Ω of positive measure), and so, from this point of view the situation is more similar to the case of constant mobility. Summing up, as far as the possibility of existence of pure phases is concerned, the difference between the cases of constant and degenerate mobility is more relevant when the mobility is degenerate, but not strongly degenerate (on this issue see also [48]).

If, in addition to the assumptions of Theorem 2, we suppose that the initial chemical potential $\mu(0) =: \mu_0$ belongs to H , then we can improve the regularity of μ . In particular, we obtain $\mu \in L^2(0, T; V)$. This is shown in the next theorem, which also requires some further reasonable conditions on the singular part F_1 of the potential F (the potential (4.2) is however included) and whose proof is based on a suitable choice of the test function (see [26]). We point out that, while in [26] the improved regularity for the chemical potential is justified only by formal computations, the argument in the proof of Theorem 3 is rigorous.

Theorem 3. *Let all assumptions of Theorem 2 be satisfied. In addition, assume that $F_1 \in C^3(-1, 1)$ and that there exist some constants $\alpha_0, \beta_0 > 0$ and $\rho \in [0, 1)$ such that the following conditions are fulfilled*

$$\rho F_1''(s) + F_2''(s) + a(x) \geq 0, \quad \forall s \in (-1, 1), \quad a.e. \text{ in } \Omega, \quad (4.66)$$

$$m(s)F_1''(s) \geq \alpha_0, \quad \forall s \in [-1, 1], \quad (4.67)$$

$$|m^2(s)F_1'''(s)| \leq \beta_0, \quad \forall s \in [-1, 1], \quad (4.68)$$

$$F_1'(s)F_1'''(s) \geq 0, \quad \forall s \in (-1, 1). \quad (4.69)$$

Let φ_0 be such that

$$F'(\varphi_0) \in H. \quad (4.70)$$

Then, the weak solution $z = [u, \varphi]$ given by Theorem 2 fulfills

$$\mu \in L^\infty(0, T; H), \quad \nabla \mu \in L^2(0, T; H). \quad (4.71)$$

As a consequence, $z = [u, \varphi]$ also satisfies the weak formulation (3.5) and (3.6), the energy inequality (3.7) and, for $d = 2$, the energy identity (3.12).

Remark 9. Note that in Theorem 3 assumptions (A2) and (A4) can be omitted. Indeed, (A2) follows from (4.67) and (A1), while (A4) follows from (4.66) and (4.67). Observe moreover that the standard potential F and the mobility m in (4.2) comply with the assumptions of Theorem 3.

Remark 10. Assumption (4.67) together with (A1) imply that the two conditions $F(\varphi_0) \in L^1(\Omega)$ and $M(\varphi_0) \in L^1(\Omega)$ are equivalent. Indeed, by combining (4.67) with the definition of function M we get $\alpha_0 M''(s) \leq F_1''(s)$, for all $s \in (-1, 1)$. Integrating this inequality we obtain

$$\alpha_0 M(s) \leq F_1(s) - F_1(0) - F_1'(0)s, \quad \forall s \in (-1, 1),$$

and, on account of (4.12), we get the above equivalence.

Proof. Let us write the weak formulation of (4.13) as

$$\langle \varphi'_\epsilon, \psi \rangle + (m_\epsilon(\varphi_\epsilon) \nabla \tilde{\mu}_\epsilon, \nabla \psi) + (m_\epsilon(\varphi_\epsilon) \nabla (a\varphi_\epsilon + F'_{2\epsilon}(\varphi_\epsilon)), \nabla \psi) = (u_\epsilon \varphi_\epsilon, \nabla \psi), \quad (4.72)$$

for all $\psi \in V$, where we have set

$$\tilde{\mu}_\epsilon := w_\epsilon + F'_{1\epsilon}(\varphi_\epsilon), \quad w_\epsilon := -J * \varphi_\epsilon,$$

and where $F_{1\epsilon}$ is defined as in (4.19). Take $\psi = F'_{1\epsilon}(\varphi_\epsilon) F''_{1\epsilon}(\varphi_\epsilon)$ as test function in (4.72) and notice that $\psi \in V$. Indeed, $F'_{1\epsilon}(\varphi_\epsilon) \in V$ and, since by (4.19) $F''_{1\epsilon}$ is globally Lipschitz on \mathbb{R} , we have also $F''_{1\epsilon}(\varphi_\epsilon) \in V$. On the other hand, by applying an Alikakos' iteration argument as in [10, Theorem 2.1], we can prove that the family of approximate solutions φ_ϵ is uniformly bounded in $L^\infty(\Omega)$. To see this, let us take $\psi = |\varphi_\epsilon|^{p-1} \varphi_\epsilon$ as test function in (4.72), where $p > 1$. Then we get the following differential identity

$$\begin{aligned} & \frac{1}{p+1} \frac{d}{dt} \|\varphi_\epsilon\|_{L^{p+1}(\Omega)}^{p+1} + p \int_{\Omega} m_\epsilon(\varphi_\epsilon) F''_{1\epsilon}(\varphi_\epsilon) |\nabla \varphi_\epsilon|^2 |\varphi_\epsilon|^{p-1} + p \int_{\Omega} m_\epsilon(\varphi_\epsilon) \nabla w_\epsilon \cdot \nabla \varphi_\epsilon |\varphi_\epsilon|^{p-1} \\ & + p \int_{\Omega} m_\epsilon(\varphi_\epsilon) \nabla (a\varphi_\epsilon + F'_{2\epsilon}(\varphi_\epsilon)) \cdot \nabla \varphi_\epsilon |\varphi_\epsilon|^{p-1} = 0. \end{aligned} \quad (4.73)$$

Actually, the above choice of test function would not be generally admissible. Nevertheless, the argument can be made rigorous by means of a density procedure, e.g., by first truncating the test function $|\varphi_\epsilon|^{p-1} \varphi_\epsilon$ and then passing to the limit with respect to the truncation parameter.

By using (4.66) we obtain

$$\begin{aligned} & p \int_{\Omega} m_\epsilon(\varphi_\epsilon) F''_{1\epsilon}(\varphi_\epsilon) |\nabla \varphi_\epsilon|^2 |\varphi_\epsilon|^{p-1} + p \int_{\Omega} m_\epsilon(\varphi_\epsilon) \nabla (a\varphi_\epsilon + F'_{2\epsilon}(\varphi_\epsilon)) \cdot \nabla \varphi_\epsilon |\varphi_\epsilon|^{p-1} \\ & = p \int_{\Omega} m_\epsilon(\varphi_\epsilon) F''_{1\epsilon}(\varphi_\epsilon) |\nabla \varphi_\epsilon|^2 |\varphi_\epsilon|^{p-1} + p \int_{\Omega} m_\epsilon(\varphi_\epsilon) (a + F''_{2\epsilon}(\varphi_\epsilon)) |\nabla \varphi_\epsilon|^2 |\varphi_\epsilon|^{p-1} \\ & + p \int_{\Omega} m_\epsilon(\varphi_\epsilon) \varphi_\epsilon \nabla a \cdot \nabla \varphi_\epsilon |\varphi_\epsilon|^{p-1} \geq (1 - \rho) p \int_{\Omega} m_\epsilon(\varphi_\epsilon) F''_{1\epsilon}(\varphi_\epsilon) |\nabla \varphi_\epsilon|^2 |\varphi_\epsilon|^{p-1} \\ & + p \int_{\Omega} m_\epsilon(\varphi_\epsilon) \varphi_\epsilon \nabla a \cdot \nabla \varphi_\epsilon |\varphi_\epsilon|^{p-1} \geq \frac{4\alpha_0 p (1 - \rho)}{(p+1)^2} \int_{\Omega} \left| \nabla |\varphi_\epsilon|^{\frac{p+1}{2}} \right|^2 \end{aligned}$$

$$+ p \int_{\Omega} m_{\epsilon}(\varphi_{\epsilon}) \varphi_{\epsilon} \nabla a \cdot \nabla \varphi_{\epsilon} |\varphi_{\epsilon}|^{p-1}. \quad (4.74)$$

Therefore, by combining (4.73) with (4.74), we deduce

$$\begin{aligned} & \frac{1}{p+1} \frac{d}{dt} \|\varphi_{\epsilon}\|_{L^{p+1}(\Omega)}^{p+1} + \frac{4\alpha_0 p(1-\rho)}{(p+1)^2} \int_{\Omega} \left| \nabla |\varphi_{\epsilon}|^{\frac{p+1}{2}} \right|^2 + p \int_{\Omega} m_{\epsilon}(\varphi_{\epsilon}) \nabla w_{\epsilon} \cdot \nabla \varphi_{\epsilon} |\varphi_{\epsilon}|^{p-1} \\ & + p \int_{\Omega} m_{\epsilon}(\varphi_{\epsilon}) \varphi_{\epsilon} \nabla a \cdot \nabla \varphi_{\epsilon} |\varphi_{\epsilon}|^{p-1} \leq 0. \end{aligned} \quad (4.75)$$

Starting from (4.75) and using the fact that $m_{\epsilon}(\varphi_{\epsilon})$ is uniformly bounded with respect to ϵ , we can argue exactly as in [10, Proof of Theorem 2.1] in order to conclude that

$$\|\varphi_{\epsilon}\|_{L^{\infty}(\Omega)} \leq C, \quad (4.76)$$

where the positive constant C depends on $\|\varphi_0\|_{L^{\infty}(\Omega)}$ and on F, J, Ω , but is independent of ϵ .

Hence, since $\varphi_{\epsilon} \in L^{\infty}(\Omega)$, then $F'_{1\epsilon}(\varphi_{\epsilon})$ and $F''_{1\epsilon}(\varphi_{\epsilon})$ are in $V \cap L^{\infty}(\Omega)$. This yields $\psi = F'_{1\epsilon}(\varphi_{\epsilon}) F''_{1\epsilon}(\varphi_{\epsilon}) \in V \cap L^{\infty}(\Omega)$. Using this test function we first see that the contribution of the convective term vanishes. Indeed, by the incompressibility condition (1.8) we deduce

$$\int_{\Omega} (u_{\epsilon} \cdot \nabla \varphi_{\epsilon}) F'_{1\epsilon}(\varphi_{\epsilon}) F''_{1\epsilon}(\varphi_{\epsilon}) = \int_{\Omega} u_{\epsilon} \cdot \nabla \left(\frac{F'^2_{1\epsilon}(\varphi_{\epsilon})}{2} \right) = 0.$$

Furthermore, we can see that we have

$$\langle \varphi'_{\epsilon}, F'_{1\epsilon}(\varphi_{\epsilon}) F''_{1\epsilon}(\varphi_{\epsilon}) \rangle = \frac{1}{2} \frac{d}{dt} \int_{\Omega} |F'_{1\epsilon}(\varphi_{\epsilon})|^2. \quad (4.77)$$

Indeed, setting $G_{\epsilon}(s) := F'^2_{1\epsilon}(s)/2$, we find $G''_{\epsilon} = F'_{1\epsilon} F'''_{1\epsilon} + F'^2_{1\epsilon} \geq 0$ almost everywhere in \mathbb{R} , due to (4.69), i.e., G_{ϵ} is convex on \mathbb{R} . Hence, (4.77) follows as an easy application of the chain rule result proved in [15, Proposition 4.2].

We are therefore led to the following identity

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \|F'_{1\epsilon}(\varphi_{\epsilon})\|^2 + \int_{\Omega} m_{\epsilon}(\varphi_{\epsilon}) F''_{1\epsilon}(\varphi_{\epsilon}) \nabla \tilde{\mu}_{\epsilon} \cdot \nabla F'_{1\epsilon}(\varphi_{\epsilon}) \\ & + \int_{\Omega} m_{\epsilon}(\varphi_{\epsilon}) F'_{1\epsilon}(\varphi_{\epsilon}) F'''_{1\epsilon}(\varphi_{\epsilon}) \nabla \tilde{\mu}_{\epsilon} \cdot \nabla \varphi_{\epsilon} + \int_{\Omega} m_{\epsilon}(\varphi_{\epsilon}) \varphi_{\epsilon} \nabla a \cdot \nabla (F'_{1\epsilon}(\varphi_{\epsilon}) F''_{1\epsilon}(\varphi_{\epsilon})) \\ & + \int_{\Omega} m_{\epsilon}(\varphi_{\epsilon}) (a + F''_{2\epsilon}(\varphi_{\epsilon})) \nabla \varphi_{\epsilon} \cdot \nabla (F'_{1\epsilon}(\varphi_{\epsilon}) F''_{1\epsilon}(\varphi_{\epsilon})) = 0, \end{aligned} \quad (4.78)$$

which can be rewritten as

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \|\tilde{\mu}_{\epsilon} - w_{\epsilon}\|^2 + \int_{\Omega} m_{\epsilon}(\varphi_{\epsilon}) F''_{1\epsilon}(\varphi_{\epsilon}) |\nabla(\tilde{\mu}_{\epsilon} - w_{\epsilon})|^2 + \int_{\Omega} m_{\epsilon}(\varphi_{\epsilon}) F''_{1\epsilon}(\varphi_{\epsilon}) \nabla w_{\epsilon} \cdot \nabla(\tilde{\mu}_{\epsilon} - w_{\epsilon}) \\ & + \int_{\Omega} m_{\epsilon}(\varphi_{\epsilon}) (\tilde{\mu}_{\epsilon} - w_{\epsilon}) F'''_{1\epsilon}(\varphi_{\epsilon}) \nabla \tilde{\mu}_{\epsilon} \cdot \nabla \varphi_{\epsilon} + \int_{\Omega} m_{\epsilon}(\varphi_{\epsilon}) \varphi_{\epsilon} F''_{1\epsilon}(\varphi_{\epsilon}) \nabla a \cdot \nabla F'_{1\epsilon}(\varphi_{\epsilon}) \end{aligned}$$

$$\begin{aligned}
& + \int_{\Omega} m_{\epsilon}(\varphi_{\epsilon}) \varphi_{\epsilon} F'_{1\epsilon}(\varphi_{\epsilon}) F'''_{1\epsilon}(\varphi_{\epsilon}) \nabla a \cdot \nabla \varphi_{\epsilon} + \int_{\Omega} m_{\epsilon}(\varphi_{\epsilon}) (a + F''_{2\epsilon}(\varphi_{\epsilon})) F''_{1\epsilon}{}^2(\varphi_{\epsilon}) |\nabla \varphi_{\epsilon}|^2 \\
& + \int_{\Omega} m_{\epsilon}(\varphi_{\epsilon}) (a + F''_{2\epsilon}(\varphi_{\epsilon})) F'_{1\epsilon}(\varphi_{\epsilon}) F'''_{1\epsilon}(\varphi_{\epsilon}) |\nabla \varphi_{\epsilon}|^2 = 0.
\end{aligned} \tag{4.79}$$

Condition (4.66) implies that

$$\rho F''_{1\epsilon}(s) + F''_{2\epsilon}(s) + a(x) \geq 0, \quad \forall s \in \mathbb{R}, \quad \text{a.a } x \in \Omega, \quad \forall \epsilon \in (0, 1). \tag{4.80}$$

Then we get

$$\begin{aligned}
& \int_{\Omega} m_{\epsilon}(\varphi_{\epsilon}) F''_{1\epsilon}(\varphi_{\epsilon}) |\nabla(\tilde{\mu}_{\epsilon} - w_{\epsilon})|^2 + \int_{\Omega} m_{\epsilon}(\varphi_{\epsilon}) (a + F''_{2\epsilon}(\varphi_{\epsilon})) F''_{1\epsilon}{}^2(\varphi_{\epsilon}) |\nabla \varphi_{\epsilon}|^2 \\
& \geq (1 - \rho) \int_{\Omega} m_{\epsilon}(\varphi_{\epsilon}) F''_{1\epsilon}(\varphi_{\epsilon}) |\nabla(\tilde{\mu}_{\epsilon} - w)|^2.
\end{aligned} \tag{4.81}$$

On the other hand, condition (4.69) implies that, for all $\epsilon \in (0, 1)$, $F'_{1\epsilon} F'''_{1\epsilon} \geq 0$ on \mathbb{R} . Thus using again (4.80), we have

$$\begin{aligned}
& \int_{\Omega} m_{\epsilon}(\varphi_{\epsilon}) (\tilde{\mu}_{\epsilon} - w_{\epsilon}) F'''_{1\epsilon}(\varphi_{\epsilon}) \nabla \tilde{\mu}_{\epsilon} \cdot \nabla \varphi_{\epsilon} + \int_{\Omega} m_{\epsilon}(\varphi_{\epsilon}) (a + F''_{2\epsilon}(\varphi_{\epsilon})) F'_{1\epsilon}(\varphi_{\epsilon}) F'''_{1\epsilon}(\varphi_{\epsilon}) |\nabla \varphi_{\epsilon}|^2 \\
& \geq \int_{\Omega} m_{\epsilon}(\varphi_{\epsilon}) F'_{1\epsilon}(\varphi_{\epsilon}) F'''_{1\epsilon}(\varphi_{\epsilon}) \nabla \tilde{\mu}_{\epsilon} \cdot \nabla \varphi_{\epsilon} - \rho \int_{\Omega} m_{\epsilon}(\varphi_{\epsilon}) F''_{1\epsilon}(\varphi_{\epsilon}) F'_{1\epsilon}(\varphi_{\epsilon}) F'''_{1\epsilon}(\varphi_{\epsilon}) |\nabla \varphi_{\epsilon}|^2 \\
& = \int_{\Omega} m_{\epsilon}(\varphi_{\epsilon}) F'_{1\epsilon}(\varphi_{\epsilon}) F'''_{1\epsilon}(\varphi_{\epsilon}) F''_{1\epsilon}(\varphi_{\epsilon}) |\nabla \varphi_{\epsilon}|^2 + \int_{\Omega} m_{\epsilon}(\varphi_{\epsilon}) F'_{1\epsilon}(\varphi_{\epsilon}) F'''_{1\epsilon}(\varphi_{\epsilon}) \nabla w_{\epsilon} \cdot \nabla \varphi_{\epsilon} \\
& - \rho \int_{\Omega} m_{\epsilon}(\varphi_{\epsilon}) F'_{1\epsilon}(\varphi_{\epsilon}) F'''_{1\epsilon}(\varphi_{\epsilon}) F''_{1\epsilon}(\varphi_{\epsilon}) |\nabla \varphi_{\epsilon}|^2 \\
& = (1 - \rho) \int_{\Omega} m_{\epsilon}(\varphi_{\epsilon}) F'_{1\epsilon}(\varphi_{\epsilon}) F'''_{1\epsilon}(\varphi_{\epsilon}) \nabla(\tilde{\mu}_{\epsilon} - w_{\epsilon}) \cdot \nabla \varphi_{\epsilon} + \int_{\Omega} m_{\epsilon}(\varphi_{\epsilon}) F'_{1\epsilon}(\varphi_{\epsilon}) F'''_{1\epsilon}(\varphi_{\epsilon}) \nabla w_{\epsilon} \cdot \nabla \varphi_{\epsilon} \\
& = (1 - \rho) \int_{\Omega} m_{\epsilon}(\varphi_{\epsilon}) F'_{1\epsilon}(\varphi_{\epsilon}) F'''_{1\epsilon}(\varphi_{\epsilon}) \nabla \tilde{\mu}_{\epsilon} \cdot \nabla \varphi_{\epsilon} + \rho \int_{\Omega} m_{\epsilon}(\varphi_{\epsilon}) F'_{1\epsilon}(\varphi_{\epsilon}) F'''_{1\epsilon}(\varphi_{\epsilon}) \nabla w_{\epsilon} \cdot \nabla \varphi_{\epsilon}.
\end{aligned} \tag{4.82}$$

Furthermore, observe that

$$m_{\epsilon}(\varphi_{\epsilon}) F'''_{1\epsilon}(\varphi_{\epsilon}) \nabla \varphi_{\epsilon} = \frac{m_{\epsilon}^2(\varphi_{\epsilon}) F'''_{1\epsilon}(\varphi_{\epsilon})}{m_{\epsilon}(\varphi_{\epsilon}) F''_{1\epsilon}(\varphi_{\epsilon})} \nabla F'_{1\epsilon}(\varphi_{\epsilon}) = \gamma_{\epsilon}(\varphi_{\epsilon}) \nabla F'_{1\epsilon}(\varphi_{\epsilon}), \tag{4.83}$$

where

$$\gamma_{\epsilon}(s) := \frac{m_{\epsilon}^2(s) F'''_{1\epsilon}(s)}{m_{\epsilon}(s) F''_{1\epsilon}(s)}, \quad \forall s \in \mathbb{R}.$$

Therefore the following identity holds

$$\int_{\Omega} m_{\epsilon}(\varphi_{\epsilon}) \varphi_{\epsilon} F'_{1\epsilon}(\varphi_{\epsilon}) F'''_{1\epsilon}(\varphi_{\epsilon}) \nabla a \cdot \nabla \varphi_{\epsilon} = \int_{\Omega} \gamma_{\epsilon}(\varphi_{\epsilon}) (\tilde{\mu}_{\epsilon} - w_{\epsilon}) \varphi_{\epsilon} \nabla a \cdot \nabla(\tilde{\mu}_{\epsilon} - w_{\epsilon}). \tag{4.84}$$

By plugging (4.81), (4.82) and (4.84) into (4.79) and using (4.83) once more, we get

$$\begin{aligned}
& \frac{1}{2} \frac{d}{dt} \|\tilde{\mu}_\epsilon - w_\epsilon\|^2 + (1 - \rho) \int_{\Omega} m_\epsilon(\varphi_\epsilon) F_{1\epsilon}''(\varphi_\epsilon) |\nabla(\tilde{\mu}_\epsilon - w_\epsilon)|^2 \\
& + \int_{\Omega} m_\epsilon(\varphi_\epsilon) F_{1\epsilon}''(\varphi_\epsilon) \nabla w_\epsilon \cdot \nabla(\tilde{\mu}_\epsilon - w_\epsilon) + (1 - \rho) \int_{\Omega} \gamma_\epsilon(\varphi_\epsilon) (\tilde{\mu}_\epsilon - w_\epsilon) |\nabla(\tilde{\mu}_\epsilon - w_\epsilon)|^2 \\
& + \int_{\Omega} \gamma_\epsilon(\varphi_\epsilon) (\tilde{\mu}_\epsilon - w_\epsilon) \nabla(\tilde{\mu}_\epsilon - w_\epsilon) \cdot \nabla w_\epsilon + \int_{\Omega} m_\epsilon(\varphi_\epsilon) F_{1\epsilon}''(\varphi_\epsilon) \varphi_\epsilon \nabla a \cdot \nabla(\tilde{\mu}_\epsilon - w_\epsilon) \\
& + \int_{\Omega} \gamma_\epsilon(\varphi_\epsilon) (\tilde{\mu}_\epsilon - w_\epsilon) \varphi_\epsilon \nabla a \cdot \nabla(\tilde{\mu}_\epsilon - w_\epsilon) \leq 0. \tag{4.85}
\end{aligned}$$

Notice that the fourth term on the left hand side is nonnegative since there holds

$$\gamma_\epsilon(\varphi_\epsilon) (\tilde{\mu}_\epsilon - w_\epsilon) = \gamma(\varphi_\epsilon) F_{1\epsilon}'(\varphi_\epsilon) = \frac{m_\epsilon^2(\varphi_\epsilon) F_{1\epsilon}'''(\varphi_\epsilon)}{m_\epsilon(\varphi_\epsilon) F_{1\epsilon}''(\varphi_\epsilon)} F_{1\epsilon}'(\varphi_\epsilon) \geq 0.$$

On the other hand, condition (4.67) implies that

$$m_\epsilon(s) F_{1\epsilon}''(s) \geq \alpha_0, \quad \forall s \in \mathbb{R}, \quad \forall \epsilon \in (0, 1), \tag{4.86}$$

and, using (H2), we have

$$\begin{aligned}
& \int_{\Omega} \gamma_\epsilon(\varphi_\epsilon) (\tilde{\mu}_\epsilon - w_\epsilon) \nabla(\tilde{\mu}_\epsilon - w_\epsilon) \cdot \nabla w_\epsilon \geq -(1 - \rho) \frac{\alpha_0}{8} \|\nabla(\tilde{\mu}_\epsilon - w_\epsilon)\|^2 \\
& - C_1 \int_{\Omega} \gamma_\epsilon^2(\varphi_\epsilon) |\nabla w_\epsilon|^2 |\tilde{\mu}_\epsilon - w_\epsilon|^2, \tag{4.87}
\end{aligned}$$

$$\begin{aligned}
& \int_{\Omega} \gamma_\epsilon(\varphi_\epsilon) (\tilde{\mu}_\epsilon - w_\epsilon) \varphi_\epsilon \nabla a \cdot \nabla(\tilde{\mu}_\epsilon - w_\epsilon) \geq -(1 - \rho) \frac{\alpha_0}{8} \|\nabla(\tilde{\mu}_\epsilon - w_\epsilon)\|^2 \\
& - C_1 \int_{\Omega} \gamma_\epsilon^2(\varphi_\epsilon) \varphi_\epsilon^2 |\nabla a|^2 |\tilde{\mu}_\epsilon - w_\epsilon|^2, \tag{4.88}
\end{aligned}$$

$$\int_{\Omega} m_\epsilon(\varphi_\epsilon) F_{1\epsilon}''(\varphi_\epsilon) \varphi_\epsilon \nabla a \cdot \nabla(\tilde{\mu}_\epsilon - w_\epsilon) \geq -(1 - \rho) \frac{\alpha_0}{8} \|\nabla(\tilde{\mu}_\epsilon - w_\epsilon)\|^2 - C_2 \|\varphi_\epsilon\|^2, \tag{4.89}$$

$$\int_{\Omega} m_\epsilon(\varphi_\epsilon) F_{1\epsilon}''(\varphi_\epsilon) \nabla w_\epsilon \cdot \nabla(\tilde{\mu}_\epsilon - w_\epsilon) \geq -(1 - \rho) \frac{\alpha_0}{8} \|\nabla(\tilde{\mu}_\epsilon - w_\epsilon)\|^2 - C_3 \|\nabla w_\epsilon\|^2. \tag{4.90}$$

In (4.89) and (4.90) we have used the fact that (4.54) holds as a consequence of the boundedness of mF'' on $[-1, 1]$.

Recall that $\gamma_\epsilon(s) = 0$ for all $|s| > 1 - \epsilon$, and $\gamma_\epsilon(s) = m^2(s) F_1'''(s) / m(s) F_1''(s)$ for $|s| \leq 1 - \epsilon$. Hence, on account of (4.67) and (4.68), we have

$$|\gamma_\epsilon(s)| \leq \gamma_0, \quad \forall s \in \mathbb{R}, \quad \forall \epsilon \in (0, 1), \tag{4.91}$$

with $\gamma_0 := \beta_0 / \alpha_0$. Thus the second term on the right hand side of (4.88) can be estimated as follows

$$\int_{\{\varphi_\epsilon \leq 1 - \epsilon\}} \gamma_\epsilon^2(\varphi_\epsilon) \varphi_\epsilon^2 |\nabla a|^2 |\tilde{\mu}_\epsilon - w_\epsilon|^2 \leq \gamma_0^2 \|\nabla a\|_{L^\infty(\Omega)}^2 \|\tilde{\mu}_\epsilon - w_\epsilon\|^2. \tag{4.92}$$

By means of (4.76) and (4.86)–(4.92), we deduce from (4.85) the inequality

$$\frac{d}{dt} \|\tilde{\mu}_\epsilon - w_\epsilon\|^2 + \alpha_0(1 - \rho) \|\nabla(\tilde{\mu}_\epsilon - w_\epsilon)\|^2 \leq C_4 + C_5 \|\tilde{\mu}_\epsilon - w_\epsilon\|^2. \quad (4.93)$$

It is easy to see that

$$|F'_{1\epsilon}(s)| \leq |F'_1(s)|, \quad \forall s \in (-1, 1), \quad \forall \epsilon \in (0, \epsilon_0], \quad (4.94)$$

with $\epsilon_0 > 0$ small enough. Therefore, condition (4.70) entails

$$\|\tilde{\mu}_\epsilon(0) - w_\epsilon(0)\| = \|F'_{1\epsilon}(\varphi_0)\| \leq \|F'_1(\varphi_0)\|,$$

and by applying Gronwall's lemma to (4.93) and using the fact that w_ϵ is bounded independently of ϵ in $L^\infty(0, T; H)$ as well as its gradient (cf. (H2) and (4.6)), we immediately get

$$\tilde{\mu} \in L^\infty(0, T; H), \quad \nabla \tilde{\mu} \in L^2(0, T; H), \quad (4.95)$$

where $\tilde{\mu} := w + F'_1(\varphi)$ and $w := -J * \varphi$. Furthermore, due to (4.20), we have (see also (4.24))

$$|F'_{2\epsilon}(s)| \leq L_3|s| + L_4, \quad |F''_{2\epsilon}(s)| \leq L_5, \quad \forall s \in \mathbb{R},$$

for some nonnegative constants L_3, L_4, L_5 which are independent of ϵ . By using these estimates, (4.32) and (4.36) we also obtain

$$F'_2(\varphi) \in L^\infty(0, T; H), \quad \nabla F'_2(\varphi) \in L^2(0, T; H). \quad (4.96)$$

From (4.95) and (4.96) we immediately get (4.71).

In order to see that (3.5) and (3.6) are satisfied, consider the weak formulation (4.52), (4.53) of the approximate problem P_ϵ , write $F''_\epsilon(\varphi_\epsilon) \nabla \varphi_\epsilon = \nabla F'_\epsilon(\varphi_\epsilon)$, rearrange the terms on the left hand side of (4.52), take the definition of μ_ϵ into account and pass to the limit as $\epsilon \rightarrow 0$.

The energy inequality (3.7) can be deduced by passing to the limit in (4.30) and by observing that

$$\int_{\Omega} F_\epsilon(\varphi_0) \leq \int_{\Omega} F_1(\varphi_0) + \int_{\Omega} F_{2\epsilon}(\varphi_0),$$

and that $\int_{\Omega} F_{2\epsilon}(\varphi_0) \rightarrow \int_{\Omega} F_2(\varphi_0)$ by Lebesgue's theorem (see (4.24)).

Finally, the energy identity (3.12) for $d = 2$ is obtained by choosing $\psi = \mu$ as test function in (3.5). \square

The argument of Theorem 3 can be exploited to prove rigorously the existence of a weak solution satisfying (4.11) with $6 < p < \infty$, for $d = 3$. We point out that the $L^\infty(L^p)$ regularity of φ follows only formally from the condition $|\varphi| \leq 1$ proved in Theorem 2. Indeed, in the case $d = 3$ it is not known that the φ component of the weak solution of Theorem 2 is measurable with values in L^p if $p > 6$. Instead, a rigorous way to deduce such regularity is by means of an approximation argument which makes use of an approximating potential with polynomial growth of order p .

Corollary 3. *Let all the assumptions of Theorem 2 and Theorem 3 be satisfied and $d = 3$. In particular let φ_0 be such that (4.70) holds. In addition, assume that there exists $\epsilon_0 > 0$ such that the following assumptions are satisfied*

$$F_1'''(s) \geq 1, \quad \forall s \in [1 - \epsilon_0, 1) \quad \text{and} \quad F_1'''(s) \leq -1, \quad \forall s \in (-1, -1 + \epsilon_0], \quad (4.97)$$

$$F_1'(s) \geq 0, \quad \forall s \in [1 - \epsilon_0, 1) \quad \text{and} \quad F_1'(s) \leq 0, \quad \forall s \in (-1, -1 + \epsilon_0]. \quad (4.98)$$

Let $p \in (6, \infty)$ be fixed arbitrarily. Then, for every $T > 0$, there exists a weak solution $z = [u, \varphi]$ to (1.5)–(1.10) on $[0, T]$ in the sense of Definition 2, satisfying the weak formulation (3.5), (3.6) and such that

$$\varphi \in L^\infty(0, T; L^p(\Omega)). \quad (4.99)$$

Proof. We argue as in the proofs of Theorem 2 and Theorem 3, but we employ a different approximation of the singular part $F_{1\epsilon}$ of the double-well potential. Namely, here we consider the approximate problem (4.13)–(4.18) with the following choice for $F_{1\epsilon}$

$$F_{1\epsilon}(s) = \begin{cases} F_1(1 - \epsilon) + F_1'(1 - \epsilon)(s - (1 - \epsilon)) + \frac{1}{2}F_1''(1 - \epsilon)(s - (1 - \epsilon))^2 \\ \quad + \frac{1}{6}(s - (1 - \epsilon))^p, & s \geq 1 - \epsilon \\ F_1(s), & |s| \leq 1 - \epsilon \\ F_1(-1 + \epsilon) + F_1'(-1 + \epsilon)(s - (-1 + \epsilon)) + \frac{1}{2}F_1''(-1 + \epsilon)(s - (-1 + \epsilon))^2 \\ \quad + \frac{1}{6}|s - (-1 + \epsilon)|^p, & s \leq -1 + \epsilon. \end{cases}$$

Let us argue as in Theorem 2 to deduce the estimates for φ_ϵ and u_ϵ (and their time derivatives). Instead, the passage to the limit will be performed differently with respect to Theorem 2.

We first check that (4.22) is still true for this new approximate potential. Indeed, for $1 - \epsilon \leq s < 1$, thanks to (4.97) we have, for some $\xi \in (1 - \epsilon, s)$

$$\begin{aligned} F_1(s) &= F_1(1 - \epsilon) + F_1'(1 - \epsilon)(s - (1 - \epsilon)) + \frac{1}{2}F_1''(1 - \epsilon)(s - (1 - \epsilon))^2 \\ &\quad + \frac{1}{6}F_1'''(\xi)(s - (1 - \epsilon))^3 \end{aligned}$$

$$\begin{aligned} &\geq F_1(1 - \epsilon) + F_1'(1 - \epsilon)(s - (1 - \epsilon)) + \frac{1}{2}F_1''(1 - \epsilon)(s - (1 - \epsilon))^2 \\ &+ \frac{1}{6}(s - (1 - \epsilon))^p = F_{1\epsilon}(s), \end{aligned}$$

provided that $\epsilon \in (0, \epsilon_0]$, with ϵ_0 as in (4.97). For $-1 < s \leq -1 + \epsilon$ we argue similarly.

It is easy to see that the uniform (with respect to ϵ) coercivity condition (4.26) will now be replaced by (cf. also (A2))

$$F_\epsilon(s) \geq \frac{1}{24}|s|^p - c_p, \quad \forall s \in \mathbb{R}, \quad \forall \epsilon \in (0, \epsilon_0], \quad (4.100)$$

where the constant c_p depends on p but is independent of ϵ . It is also immediate to check that (4.29) and (H7) hold for the new approximate potential. Therefore, since we have also $F_\epsilon \in C_{loc}^{2,1}(\mathbb{R})$, assumptions (H1)–(H6) of Theorem 1, with (H4) replaced by (H7), are satisfied. Hence, for every fixed $\epsilon \in (0, \epsilon_0]$, Problem P_ϵ admits a weak solution satisfying (3.1), the first of (3.2), (3.9), (3.10), and the energy inequality (4.30). By means of (4.25), (4.29) and (4.100), from the approximate energy inequality we can still recover estimates (4.31), (4.33), (4.36) and (4.37), while estimate (4.32) will now be substituted with the stronger

$$\|\varphi_\epsilon\|_{L^\infty(0,T;L^p(\Omega))} \leq C. \quad (4.101)$$

The estimates for u'_ϵ and φ'_ϵ can be improved as well. Indeed, instead of (4.39) and (4.42) we now find

$$\|\varphi'_\epsilon\|_{L^2(0,T;V')} \leq C, \quad d = 3, \quad (4.102)$$

and

$$\|u'_\epsilon\|_{L^2(0,T;V'_{div})} \leq C, \quad \|\varphi'_\epsilon\|_{L^2(0,T;V')} \leq C, \quad d = 2, \quad (4.103)$$

respectively, while (4.41) still holds. By compactness results we can still deduce that there exists a couple $z := [u, \varphi]$ and a subsequence such that (4.43)–(4.50) hold. Instead, in (4.46), (4.49) and (4.50) we shall have L^2 in place of $L^{2-\gamma}$, $L^{4/3}$ and $L^{2-\delta}$, respectively, and

$$\varphi_\epsilon \rightharpoonup \varphi \quad \text{weakly}^* \text{ in } L^\infty(0, T; L^p(\Omega)), \quad \text{weakly in } L^2(0, T; V), \quad (4.104)$$

in place of (4.47).

As far as the passage to the limit is concerned, the situation is now a bit different. Indeed, due to the polynomial growth of $F_{1\epsilon}$, a strong convergence for the term $m_\epsilon(\varphi_\epsilon)F_{1\epsilon}''(\varphi_\epsilon)$ in some $L^s(Q_T)$ space with $s \geq 2$ is no longer available in order to deduce the weak formulation (4.9) for φ . Therefore, the idea is to use (4.70), argue as in

Theorem 3, get a control for μ_ϵ in $L^2(0, T; V)$ and pass to the limit to deduce the weak formulation (3.5) for φ . Of course, we have to check that the argument of Theorem 3 still applies with this different choice of $F_{1\epsilon}$. Now, it is easy to verify that the test function $\psi = F'_{1\epsilon}(\varphi_\epsilon)F''_{1\epsilon}(\varphi_\epsilon)$ in the weak formulation of (4.13) is still in V . Indeed, $F''_{1\epsilon}$ is now locally Lipschitz on \mathbb{R} and $\varphi_\epsilon \in V \cap L^\infty(\Omega)$, which implies that $F''_{1\epsilon}(\varphi_\epsilon) \in V \cap L^\infty(\Omega)$. Thus $\psi \in V$.

By repeating the calculations in the first part of the proof of Theorem 3 we are then led again to the differential identity (4.79). It is immediate to check that (4.86) still holds. As far as (4.91) is concerned, notice that in this case we have for, e.g., $s > 1 - \epsilon$

$$\begin{aligned} \gamma_\epsilon(s) &= \frac{m^2(1-\epsilon)\frac{1}{6}p(p-1)(p-2)(s-(1-\epsilon))^{p-3}}{m(1-\epsilon)\left(F''_1(1-\epsilon) + \frac{1}{6}p(p-1)(s-(1-\epsilon))^{p-2}\right)} \\ &\leq \frac{c_{1,p}m^2(1-\epsilon)(s-(1-\epsilon))^{p-3}}{\alpha_0 + c_{2,p}m(1-\epsilon)(s-(1-\epsilon))^{p-2}}, \end{aligned}$$

where (4.67) has been taken into account. Now, if we consider the function

$$\psi_\delta(\tau) := \frac{\delta^2\tau^{p-3}}{1 + \delta\tau^{p-2}}, \quad \forall \tau \geq 0,$$

by an elementary calculation we can see that ψ_δ has its maximum at $\tau^* := ((p-3)/\delta)^{\frac{1}{p-2}}$ and that $\psi_\delta(\tau) \leq \psi_\delta(\tau^*) = c_p\delta^{(p-1)/(p-2)} \leq c_p$, for all $\tau \geq 0$ and for all $\delta \in (0, 1)$, where c_p is a positive constant depending only on p . Hence we deduce that $|\gamma_\epsilon(s)| \leq c'_p$ for all $|s| > 1 - \epsilon$ and for all $\epsilon \in (0, 1)$. Using also the fact that $\gamma_\epsilon(s) = m^2(s)F'''_1(s)/m(s)F''_1(s)$ for $|s| \leq 1 - \epsilon$, and taking (4.67) and (4.68) into account, we can therefore conclude that

$$|\gamma_\epsilon(s)| \leq \gamma_{0,p}, \quad \forall s \in \mathbb{R}, \quad \forall \epsilon \in (0, 1), \quad (4.105)$$

with $\gamma_{0,p}$ given by $\gamma_{0,p} = \max(c'_p, \beta_0/\alpha_0)$ and independent of ϵ . Hence, (4.91) is still satisfied.

On the other hand, using (A2) and (4.98), it is easy to check that also conditions

$$F'_{1\epsilon}F'''_{1\epsilon} \geq 0 \quad \text{on } \mathbb{R}, \quad \forall \epsilon \in (0, 1),$$

and (4.94) still hold. We can therefore conclude that the argument of Theorem 3 applies and that, due also to (4.70), the following estimates hold (cf. (??))

$$\|\mu_\epsilon\|_{L^\infty(0,T;H)} \leq C, \quad \|\nabla\mu_\epsilon\|_{L^2(0,T;H)} \leq C, \quad (4.106)$$

where the constant C depends on $\|\mu_0\|$ or, equivalently, on $\|F'(\varphi_0)\|$.

Finally, let us consider the weak formulation (3.5) and (3.6) for the approximate problem (4.13)–(4.18). By passing to the limit as $\epsilon \rightarrow 0$ and by means of (4.106) and the

convergences provided in the proof of Theorem 2, it is not difficult to show that $z = [u, \varphi]$, which in particular satisfies (4.99) due to the first of (4.104), satisfies the weak formulation (3.5) and (3.6). \square

5 The global attractor in 2D

In this section we consider system (1.5)–(1.10) for $d = 2$ and we suppose that the external force h is time-independent, namely,

(A5) $h \in V'_{div}$.

Let us introduce the set \mathcal{G}_{m_0} of all weak solutions to (1.5)–(1.10) (in the sense of Definition 2) corresponding to all initial data $z_0 = [u_0, \varphi_0] \in \mathcal{X}_{m_0}$, where the phase space \mathcal{X}_{m_0} is the metric space defined by

$$\mathcal{X}_{m_0} := G_{div} \times \mathcal{Y}_{m_0},$$

with \mathcal{Y}_{m_0} given by

$$\mathcal{Y}_{m_0} := \{\varphi \in L^\infty(\Omega) : |\varphi| \leq 1 \text{ a.e. in } \Omega, \quad F(\varphi), M(\varphi) \in L^1(\Omega), \quad |\overline{\varphi}| \leq m_0\}, \quad (5.1)$$

and $m_0 \in [0, 1]$ is fixed. The metric on \mathcal{X}_{m_0} is

$$\mathbf{d}(z_2, z_1) := \|u_2 - u_1\| + \|\varphi_2 - \varphi_1\|,$$

for every $z_1 := [u_1, \varphi_1]$ and $z_2 := [u_2, \varphi_2]$ in \mathcal{X}_{m_0} .

In the next proposition we shall prove that the set of all weak solution is a generalized semiflow in the sense of J.M. Ball (cf. [8]).

Proposition 1. *Let $d = 2$ and suppose that the conditions of Theorem 2 are satisfied. Assume also that (4.66), (4.67) and that (A5) hold. Then \mathcal{G}_{m_0} is a generalized semiflow on \mathcal{X}_{m_0} .*

Proof. It is immediate to check that hypotheses (H1), (H2) and (H3) of the definition of generalized semiflow [8, Definition 2.1] are satisfied. It remains to prove the upper semicontinuity with respect to initial data, i.e., that \mathcal{G}_{m_0} satisfies (H4) of [8, Definition 2.1]. Take then $z_j = [u_j, \varphi_j] \in \mathcal{G}_{m_0}$ such that $z_j(0) \rightarrow z_0$ in \mathcal{X}_{m_0} . Our aim is to prove that there exist $z \in \mathcal{G}_{m_0}$ with $z(0) = z_0$ and a subsequence $\{z_{j_k}\}$ such that $z_{j_k}(t) \rightarrow z(t)$ in \mathcal{X}_{m_0} for all $t \geq 0$. Now, each weak solution $z_j = [u_j, \varphi_j]$ satisfies the regularity properties (4.3)–(4.8), (4.11) and the energy equation (3.44) which, on account of (4.66), (4.67) and of the fact that $|\varphi_j| \leq 1$, implies

$$\frac{d}{dt} \left(\|u_j\|^2 + \|\varphi_j\|^2 \right) + (1 - \rho)\alpha_0 \|\nabla \varphi_j\|^2 + \nu \|\nabla u_j\|^2 \leq c + c\|u_j\|^2 + \frac{1}{2\nu} \|h\|_{V'_{div}}^2,$$

where the positive constant c depends on J and on m . By integrating this inequality between 0 and t , using the fact that $z_j(0) \rightarrow z_0$ in \mathcal{X}_{m_0} and Gronwall's lemma, from the above differential inequality we get

$$\|u_j\|_{L^\infty(0,T;G_{div}) \cap L^2(0,T;V_{div})} \leq C, \quad \|\varphi_j\|_{L^\infty(0,T;H) \cap L^2(0,T;V)} \leq C.$$

By comparison in the variational formulation (4.9) and (4.10) written for each weak solution $z_j = [u_j, \varphi_j]$ we also obtain the estimates for the time derivatives u'_j and φ'_j

$$\|u'_j\|_{L^2(0,T;V'_{div})} \leq C, \quad \|\varphi'_j\|_{L^2(0,T;V')} \leq C.$$

Therefore, by standard compactness results, we deduce that there exist two functions u and φ such that, for a not relabeled subsequence, we have

$$u_j \rightharpoonup u \quad \text{weakly}^* \text{ in } L^\infty(0, T; G_{div}), \quad \text{weakly in } L^2(0, T; V_{div}), \quad (5.2)$$

$$u_j \rightarrow u \quad \text{strongly in } L^2(0, T; G_{div}), \quad \text{a.e. in } Q_T, \quad (5.3)$$

$$u'_j \rightharpoonup u_t \quad \text{weakly in } L^2(0, T; V'_{div}), \quad (5.4)$$

$$\varphi_j \rightharpoonup \varphi \quad \text{weakly}^* \text{ in } L^\infty(0, T; H), \quad \text{weakly in } L^2(0, T; V), \quad (5.5)$$

$$\varphi_j \rightarrow \varphi \quad \text{strongly in } L^2(0, T; H), \quad \text{a.e. in } Q_T, \quad (5.6)$$

$$\varphi'_j \rightharpoonup \varphi_t \quad \text{weakly in } L^2(0, T; V'). \quad (5.7)$$

Passing to the limit in the variational formulation for $z_j = [u_j, \varphi_j]$ and using the weak/strong convergences (5.2)–(5.7) and the fact that $|\varphi_j| \leq 1$, we get that $z := [u, \varphi] \in \mathcal{G}_{m_0}$ (the argument is similar to the passage to the limit in (4.52) and (4.53) in the proof of Theorem 2). Furthermore we have $z(0) = z_0$, since, as a consequence of (5.2), (5.4), (5.5) and (5.7), we have, for all $t \geq 0$

$$u_j(t) \rightharpoonup u(t) \quad \text{weakly in } G_{div}, \quad (5.8)$$

$$\varphi_j(t) \rightharpoonup \varphi(t) \quad \text{weakly in } H. \quad (5.9)$$

Let us now see that $z_j(t) \rightarrow z(t)$ in \mathcal{X}_{m_0} for all $t \geq 0$. First observe that the energy equation (3.44) can be written in the form

$$\frac{d}{dt} \tilde{E}(z_j(t)) + \int_{\Omega} m(\varphi_j) F''(\varphi_j) |\nabla \varphi_j|^2 + \int_{\Omega} a m(\varphi_j) |\nabla \varphi_j|^2 + \nu \|\nabla u_j\|^2 = 0, \quad (5.10)$$

where

$$\begin{aligned} \tilde{E}(z_j(t)) &:= E(z_j(t)) - \int_0^t \int_{\Omega} m(\varphi_j) (\nabla J * \varphi_j - \varphi_j \nabla a) \cdot \nabla \varphi_j \\ &\quad - \int_0^t \int_{\Omega} (a \varphi_j - J * \varphi_j) u_j \cdot \nabla \varphi_j - \int_0^t \langle h, u_j \rangle d\tau, \end{aligned}$$

and

$$E(z_j(t)) := \frac{1}{2}(\|u_j(t)\|^2 + \|\varphi_j(t)\|^2).$$

On the other hand, from (5.3) and (5.6) we have, for almost any $t > 0$,

$$\begin{aligned} u_j(t) &\rightarrow u(t) && \text{strongly in } G_{div}, \\ \varphi_j(t) &\rightarrow \varphi(t) && \text{strongly in } H, \end{aligned}$$

and hence $E(z_j(t)) \rightarrow E(z(t))$ for almost any $t > 0$. Moreover, due to the condition $|\varphi_j| \leq 1$ for each j , to the pointwise convergence (5.6) and Lebesgue's theorem we have

$$\varphi_j \rightarrow \varphi, \quad \text{strongly in } L^s(Q_T), \quad \forall s \in [2, \infty).$$

Therefore, recalling that $m(\varphi_j) \rightarrow m(\varphi)$ strongly in $L^s(Q_T)$ for all $s \in [2, \infty)$ and using the second weak convergence in (5.5), we have, for all $t \geq 0$,

$$\int_0^t \int_{\Omega} m(\varphi_j)(\nabla J * \varphi_j - \varphi_j \nabla a) \cdot \nabla \varphi_j \rightarrow \int_0^t \int_{\Omega} m(\varphi)(\nabla J * \varphi - \varphi \nabla a) \cdot \nabla \varphi.$$

Furthermore, since $a\varphi_j - J * \varphi_j \rightarrow a\varphi - J * \varphi$ strongly in $L^s(Q_T)$ for all $s \in [2, \infty)$ and, by (5.2), $u_j \rightarrow u$ strongly in $L^s(Q_T)^2$ for all $s \in [2, 4)$, then we also have, for all $t \geq 0$

$$\int_0^t \int_{\Omega} (a\varphi_j - J * \varphi_j)u_j \cdot \nabla \varphi_j \rightarrow \int_0^t \int_{\Omega} (a\varphi - J * \varphi)u \cdot \nabla \varphi.$$

Therefore

$$\tilde{E}(z_j(t)) \rightarrow \tilde{E}(z(t)), \quad \text{a.a. } t > 0.$$

Since, due to (5.10), $\tilde{E}(z_j(\cdot))$ is non increasing in $[0, \infty)$ for every j and $\tilde{E}(z_j(\cdot)), \tilde{E}(z(\cdot)) \in C([0, \infty))$, we infer

$$\tilde{E}(z_j(t)) \rightarrow \tilde{E}(z(t)), \quad \forall t \geq 0,$$

and hence also $E(z_j(t)) \rightarrow E(z(t))$ for all $t \geq 0$. This last convergence together with (5.8) and (5.9) yield $u_j(t) \rightarrow u(t)$ strongly in G_{div} and $\varphi_j(t) \rightarrow \varphi(t)$ strongly in H , and hence $z_j(t) \rightarrow z(t)$ in \mathcal{X}_{m_0} , for all $t \geq 0$. \square

Proposition 2. *Let $d = 2$ and suppose that the conditions of Theorem 2 are satisfied. Assume also that (4.66), (4.67) and (A5) hold. Then \mathcal{G}_{m_0} is point dissipative and eventually bounded.*

Proof. Let us estimate the second term on the right hand side of (3.44) as follows

$$\begin{aligned} \left| \int_{\Omega} (a\varphi - J * \varphi) u \cdot \nabla \varphi \right| &= \left| \int_{\Omega} \left(\frac{1}{2} \varphi^2 \nabla a - \varphi (\nabla J * \varphi) \right) \cdot u \right| \\ &\leq \frac{3}{2} b |\Omega|^{1/2} \|u\| \leq \frac{\nu}{4} \|\nabla u\|^2 + C_0, \end{aligned}$$

where $C_0 = 9b^2|\Omega|/4\nu\lambda_1$ with the constant b defined as in (H2). Moreover, the first term on the right hand side of (3.44) can be controlled in the following way

$$\left| \int_{\Omega} m(\varphi) (\nabla J * \varphi - \varphi \nabla a) \cdot \nabla \varphi \right| \leq 2m_0 |\Omega|^{1/2} b \|\nabla \varphi\| \leq (1 - \rho) \frac{\alpha_0}{2} \|\nabla \varphi\|^2 + C_1,$$

where $C_1 = 2m_*^2|\Omega|b^2/\alpha_0$, and $m_* = \max_{s \in [-1,1]} m(s)$. Then, by taking (4.66) and (4.67) into account, we get the differential inequality

$$\frac{d}{dt} (\|u\|^2 + \|\varphi\|^2) + (1 - \rho) \alpha_0 \|\nabla \varphi\|^2 + \nu \|\nabla u\|^2 \leq C_2 + \frac{1}{\nu} \|h\|_{V'_{div}}^2,$$

where $C_2 = 2(C_0 + C_1)$. By using the identity $\|\varphi\|^2 = \|\varphi - \bar{\varphi}_0\|^2 + \|\bar{\varphi}_0\|^2$, we obtain

$$\frac{d}{dt} (\|u\|^2 + \|\varphi - \bar{\varphi}_0\|^2) + (1 - \rho) \alpha_0 C_P \|\varphi - \bar{\varphi}_0\|^2 + \nu \lambda_1 \|u\|^2 \leq C_2 + \frac{1}{\nu} \|h\|_{V'_{div}}^2,$$

where C_P is the constant appearing in the Poincaré-Wirtinger inequality. Therefore we get

$$\frac{d}{dt} E(\tilde{z}(t)) + \eta E(\tilde{z}(t)) \leq C_3, \quad (5.11)$$

where $\tilde{z} = [u, \varphi - \bar{\varphi}_0]$ and the constants η and C_3 are given by $\eta = \min((1 - \rho) \alpha_0 C_P, \nu \lambda_1)/2$, and $C_3 = C_2/2 + \|h\|_{V'_{div}}^2/2\nu$, respectively. By Gronwall's lemma from (5.11) we have

$$E(\tilde{z}(t)) \leq E(\tilde{z}(0)) e^{-\eta t} + \frac{C_3}{\eta}, \quad \forall t \geq 0.$$

This estimate easily yields

$$\mathbf{d}^2(z(t), 0) \leq \mathbf{d}^2(z_0, 0) e^{-\eta t} + \frac{2C_3}{\eta} + |\bar{\varphi}_0|^2 |\Omega|, \quad \forall t \geq 0,$$

which entails both the point dissipativity and the eventual boundedness of \mathcal{G}_{m_0} .

We can now state the result on the existence of the global attractor.

Proposition 3. *Let the assumptions of Proposition 2 hold. Then \mathcal{G}_{m_0} possesses a global attractor.*

Proof. Thanks to Proposition 2 and by [8, Proposition 3.2] and [8, Theorem 3.3] we only need to show that \mathcal{G}_{m_0} is compact. Let $\{z_j\} \subset \mathcal{G}$ be a sequence with $\{z_j(0)\}$ bounded in \mathcal{X}_{m_0} . We claim that there exists a subsequence $\{z_{j_k}\}$ such that $z_{j_k}(t)$ converges in \mathcal{X}_{m_0} for every $t > 0$. Indeed, the energy equation (3.44) entails the existence of a subsequence (not relabeled) such that (cf. the proof of Proposition 1), for almost all $t > 0$,

$$u_j(t) \rightarrow u(t) \quad \text{strongly in } G_{div}, \quad \varphi_j(t) \rightarrow \varphi(t) \quad \text{strongly in } H \text{ and a.e. in } \Omega,$$

where $z = [u, \varphi]$ is a weak solution. By arguing as in the proof of Proposition 1 we infer that $\tilde{E}(z_j(t)) \rightarrow \tilde{E}(z(t))$ for all $t \geq 0$. Thus $z_j(t) \rightarrow z(t)$ in \mathcal{X}_{m_0} for $t > 0$, which yields the compactness of \mathcal{G}_{m_0} . \square

Remark 11. We point out that the existence of the global attractor has been established without the restriction $|\bar{\varphi}| < 1$ on the generalized semiflow (compare with Remark 7). In particular, this result does not require the separation property. It is also worth observing that in dimension three the trajectory attractor approach used in [19, 20] might be extended to the present case.

Remark 12. *As mentioned in the Introduction, uniqueness of weak solutions in two dimensions has been recently proved in [18]. As a consequence, the generalized semiflow becomes a semigroup and the global attractor is connected (see [18, Section 5]).*

6 The convective nonlocal Cahn-Hilliard equation with degenerate mobility

By relying on the results of the previous sections we can prove similar results for the convective nonlocal Cahn-Hilliard equation with degenerate mobility and with a given velocity field, for $d = 2, 3$. In particular, from Theorem 2 it is straightforward to deduce the following

Theorem 4. *Assume that (A1)-(A4) and (H2) hold. Let $u \in L^2_{loc}([0, \infty); V_{div} \cap L^\infty(\Omega)^d)$ be given and let $\varphi_0 \in L^\infty(\Omega)$ be such that $F(\varphi_0) \in L^1(\Omega)$ and $M(\varphi_0) \in L^1(\Omega)$. Then, for every $T > 0$, there exists a weak solution φ to (1.5), (1.6), (1.9)₁ on $[0, T]$ corresponding to φ_0 fulfilling (4.6)-(4.8), the weak formulation (4.9), and such that $\bar{\varphi}(t) = \bar{\varphi}_0$ for all $t \in [0, T]$. Furthermore, φ satisfies*

$$\varphi \in L^\infty(0, T; L^p(\Omega)), \tag{6.1}$$

where $p \leq 6$ for $d = 3$ and $2 \leq p < \infty$ for $d = 2$. In addition, for almost any $t > 0$, the following energy identity holds

$$\frac{1}{2} \frac{d}{dt} \|\varphi\|^2 + \int_{\Omega} m(\varphi) F''(\varphi) |\nabla \varphi|^2 + \int_{\Omega} a m(\varphi) |\nabla \varphi|^2 + \int_{\Omega} m(\varphi) (\varphi \nabla a - \nabla J * \varphi) \cdot \nabla \varphi = 0. \quad (6.2)$$

Remark 13. The basic estimates in the proof of Theorem 4 are now obtained from (6.2) (written for the approximate solution φ_ϵ and integrated between 0 and t), in place of (4.30). In particular, (6.2) still yields (4.32), (4.36) and (4.37), so that we can argue as in the proof of Theorem 3.

For the convective nonlocal Cahn-Hilliard equation with degenerate mobility we can also prove a uniqueness result (see [28] for more restrictive assumptions). We remind that uniqueness of solutions is an open issue for the local case (see [16]) as well as for the complete system (1.5)-(1.10) even in dimension two.

Proposition 4. *Let all the assumptions of Theorem 4 and (4.66), (4.67) be satisfied. Then, the weak solution to (1.5), (1.6), (1.9)₁, (1.10)₂ is unique.*

Proof. Following [28], let us introduce

$$\Lambda_1(s) := \int_0^s m(\sigma) F_1''(\sigma) d\sigma, \quad \Lambda_2(s) := \int_0^s m(\sigma) F_2''(\sigma) d\sigma, \quad \Gamma(s) := \int_0^s m(\sigma) d\sigma,$$

for all $s \in [-1, 1]$. Due to (A1) and (4.67) we have $\Lambda_1 \in C^1([-1, 1])$ and $0 < \alpha_0 \leq \Lambda_1'(s) \leq a$ for some positive constant a . Then, it is easy to see that the weak formulation (4.9) can be rewritten as follows

$$\langle \varphi_t, \psi \rangle + (\nabla \Lambda(\cdot, \varphi), \nabla \psi) - (\Gamma(\varphi) \nabla a, \nabla \psi) + (m(\varphi) (\varphi \nabla a - \nabla J * \varphi), \nabla \psi) = (u\varphi, \nabla \psi), \quad (6.3)$$

for all $\psi \in V$, where

$$\Lambda(x, s) := \Lambda_1(s) + \Lambda_2(s) + a(x)\Gamma(s).$$

Consider now (6.3) for two weak solutions φ_1 and φ_2 corresponding to the same initial datum. Let us take the difference between the two identities, set $\varphi := \varphi_1 - \varphi_2$ and choose $\psi = \mathcal{N}\varphi$ as test function in the resulting identity (notice that $\bar{\varphi} = 0$). This yields

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \|\mathcal{N}^{1/2} \varphi\|^2 + (\Lambda(\varphi_2) - \Lambda(\varphi_1), \varphi) - ((\Gamma(\varphi_2) - \Gamma(\varphi_1)) \nabla a, \nabla \mathcal{N}\varphi) \\ & + ((m(\varphi_2) - m(\varphi_1)) (\varphi_2 \nabla a - \nabla J * \varphi_2) + m(\varphi_1) (\varphi \nabla a - \nabla J * \varphi), \nabla \mathcal{N}\varphi) \\ & = (u\varphi, \nabla \mathcal{N}\varphi). \end{aligned} \quad (6.4)$$

On account of (4.66) and (4.67), we find

$$\begin{aligned} (\Lambda(\cdot, \varphi_2) - \Lambda(\cdot, \varphi_1), \varphi) &\geq (1 - \rho) \int_{\Omega} m(\theta\varphi_2 + (1 - \theta)\varphi_1) F_1''(\theta\varphi_2 + (1 - \theta)\varphi_1) \varphi^2 \\ &\geq (1 - \rho) \alpha_0 \|\varphi\|^2. \end{aligned} \quad (6.5)$$

Furthermore, since $|\varphi_1|, |\varphi_2| \leq 1$, then we have

$$|((\Gamma(\varphi_2) - \Gamma(\varphi_1))\nabla a, \nabla \mathcal{N}\varphi)| \leq m_* b \|\varphi\| \|\nabla \mathcal{N}\varphi\| \leq \frac{1}{8} (1 - \rho) \alpha_0 \|\varphi\|^2 + C_1 \|\nabla \mathcal{N}\varphi\|^2, \quad (6.6)$$

$$\begin{aligned} |(m(\varphi_2) - m(\varphi_1))(\varphi_2 \nabla a - \nabla J * \varphi_2), \nabla \mathcal{N}\varphi| &\leq 2m_{**} b \|\varphi\| \|\nabla \mathcal{N}\varphi\| \\ &\leq \frac{1}{8} (1 - \rho) \alpha_0 \|\varphi\|^2 + C_2 \|\nabla \mathcal{N}\varphi\|^2, \end{aligned} \quad (6.7)$$

$$\begin{aligned} |(m(\varphi_1)(\varphi \nabla a - \nabla J * \varphi), \nabla \mathcal{N}\varphi)| &\leq 2m_* b \|\varphi\| \|\nabla \mathcal{N}\varphi\| \\ &\leq \frac{1}{8} (1 - \rho) \alpha_0 \|\varphi\|^2 + C_3 \|\nabla \mathcal{N}\varphi\|^2, \end{aligned} \quad (6.8)$$

$$\begin{aligned} |(u\varphi, \nabla \mathcal{N}\varphi)| &\leq \|u\|_{L^\infty(\Omega)^d} \|\varphi\| \|\mathcal{N}^{1/2}\varphi\| \\ &\leq \frac{1}{8} (1 - \rho) \alpha_0 \|\varphi\|^2 + C_4 \|u\|_{L^\infty(\Omega)^d}^2 \|\mathcal{N}^{1/2}\varphi\|^2, \end{aligned} \quad (6.9)$$

where b is given as in (H2), m_* as in the proof of Proposition 2, $m_{**} := \max_{s \in [-1, 1]} m'(s)$ and the positive constants C_1, \dots, C_4 depend on $\alpha_0, \rho, m_*, m_{**}$. By plugging (6.5)–(6.9) into (6.4) we get

$$\frac{d}{dt} \|\mathcal{N}^{1/2}\varphi\|^2 + (1 - \rho) \alpha_0 \|\varphi\|^2 \leq C_5 (1 + \|u\|_{L^\infty(\Omega)^d}^2) \|\mathcal{N}^{1/2}\varphi\|^2 \quad (6.10)$$

and Gronwall's lemma applies.

As a consequence of Theorem 4 and of Proposition 4 we can define a semiflow $S(t)$ on \mathcal{Y}_{m_0} (cf. (5.1)), $m_0 \in [0, 1]$, endowed with the metric induced by the L^2 -norm. It is then immediate to check that the arguments used in the proofs of Proposition 1, Proposition 2 and Proposition 5 can be adapted to the present situation. This gives

Theorem 5. *Let (A1)–(A4), (4.66) and (4.67) hold. Suppose that $u \in L^\infty(\Omega)^d \cap V_{div}$ is given and independent of time. Then the dynamical system $(\mathcal{Y}_{m_0}, S(t))$ possesses a connected global attractor.*

Remark 14. The fact u must be divergence-free is not necessary (see also [19, 20]). In particular, the convective term can be of the form $\nabla \cdot (u\varphi)$.

Remark 15. In [25, Section 4] the case $u = 0$ was considered. In particular, the existence of an exponential attractor was proven and, as a consequence, the existence of a global attractor of finite fractal dimension. However, contrary to the present case, in [25] pure phases were a priori excluded from the phase space (cf. also Remark 11).

Acknowledgments. The authors would like to thank Ciprian G. Gal for a useful remark concerning the first part of the proof of Theorem 3. The authors are members of the Gruppo Nazionale per l'Analisi Matematica, la Probabilità e le loro Applicazioni (GNAMPA) of the Istituto Nazionale di Alta Matematica (INdAM).

References

- [1] H. Abels, *On a diffusive interface model for two-phase flows of viscous, incompressible fluids with matched densities*, Arch. Ration. Mech. Anal. **194** (2009), 463-506.
- [2] H. Abels, *Longtime behavior of solutions of a Navier-Stokes/Cahn-Hilliard system*, Proceedings of the Conference “Nonlocal and Abstract Parabolic Equations and their Applications”, Bedlewo, Banach Center Publ. **86** (2009), 9-19.
- [3] H. Abels, D. Depner, H. Garcke, *Existence of weak solutions for a diffuse interface model for two-phase flows of incompressible fluids with different densities*, J. Math. Fluid Mech. **15** (2013), 453-480.
- [4] H. Abels, D. Depner, H. Garcke, *On an incompressible Navier-Stokes/Cahn-Hilliard system with degenerate mobility*, Ann. Inst. H. Poincaré Anal. Non Linéaire **30** (2013), 1175-1190.
- [5] H. Abels, L. Diening, Y. Terasawa, *Existence of Weak Solutions for a Diffuse Interface Model of Non-Newtonian Two-Phase Flows*, Nonlinear Anal. Real World Appl. **15** (2014), 149-157.
- [6] H. Abels, E. Feireisl, *On a diffuse interface model for a two-phase flow of compressible viscous fluids*, Indiana Univ. Math. J. **57** (2008), 659–698.
- [7] D.M. Anderson, G.B. McFadden, A.A. Wheeler, *Diffuse-interface methods in fluid mechanics*, Annu. Rev. Fluid Mech. **30**, Annual Reviews, Palo Alto, CA, 1998, 139-165.
- [8] J.M. Ball, *Continuity properties and global attractors of generalized semiflows and the Navier-Stokes equation*, J. Nonlinear Sci. **7** (1997), 475-502 (Erratum, J. Nonlinear Sci. **8** (1998), 233).
- [9] J.W. Barrett, J.F. Blowey, *Finite element approximation of the Cahn-Hilliard equation with concentration dependent mobility*, Math. Comp. **68** (1999), 487-517.
- [10] P.W. Bates, J. Han, *The Neumann boundary problem for a nonlocal Cahn-Hilliard equation*, J. Differential Equations **212** (2005), 235-277.

- [11] S. Bosia, *Analysis of a Cahn-Hilliard-Ladyzhenskaya system with singular potential*, J. Math. Anal. Appl. **397** (2013), 307-321.
- [12] F. Boyer, *Mathematical study of multi-phase flow under shear through order parameter formulation*, Asymptot. Anal. **20** (1999), 175-212.
- [13] C. Cao, C.G. Gal, *Global solutions for the 2D NS-CH model for a two-phase flow of viscous, incompressible fluids with mixed partial viscosity and mobility*, Nonlinearity **25** (2012), 3211-3234.
- [14] P. Colli, S. Frigeri, M. Grasselli, *Global existence of weak solutions to a nonlocal Cahn-Hilliard-Navier-Stokes system*, J. Math. Anal. Appl. **386** (2012), 428-444.
- [15] P. Colli, P. Krejčí, E. Rocca, J. Sprekels, *Nonlinear evolution inclusions arising from phase change models*, Czechoslovak Math. J. **57** (2007), 1067-1098.
- [16] C.M. Elliott, H. Garcke, *On the Cahn-Hilliard equation with degenerate mobility*, SIAM J. Math. Anal. **27** (1996), 404-423.
- [17] H. Emmerich, *The diffuse interface approach in materials science*, Springer, Berlin Heidelberg, 2003.
- [18] S. Frigeri, C.G. Gal, M. Grasselli, *On nonlocal Cahn-Hilliard-Navier-Stokes systems in two dimensions*, submitted. Wias Preprint **1923**, (2014), 34 pp.
- [19] S. Frigeri, M. Grasselli, *Global and trajectory attractors for a nonlocal Cahn-Hilliard-Navier-Stokes system*, J. Dynam. Differential Equations **24** (2012), 827-856.
- [20] S. Frigeri, M. Grasselli, *Nonlocal Cahn-Hilliard-Navier-Stokes systems with singular potentials*, Dyn. Partial Differ. Equ. **9** (2012), 273-304.
- [21] S. Frigeri, M. Grasselli, P. Krejčí, *Strong solutions for two-dimensional nonlocal Cahn-Hilliard-Navier-Stokes systems*, J. Differential Equations **255** (2013), 2587-2614.
- [22] C.G. Gal, M. Grasselli, *Asymptotic behavior of a Cahn-Hilliard-Navier-Stokes system in 2D*, Ann. Inst. H. Poincaré Anal. Non Linéaire **27** (2010), 401-436.
- [23] C.G. Gal, M. Grasselli, *Trajectory attractors for binary fluid mixtures in 3D*, Chinese Ann. Math. Ser. B **31** (2010), 655-678.
- [24] C.G. Gal, M. Grasselli, *Instability of two-phase flows: a lower bound on the dimension of the global attractor of the Cahn-Hilliard-Navier-Stokes system*, Phys. D **240** (2011), 629-635.

- [25] C.G. Gal, M. Grasselli, *Longtime behavior of nonlocal Cahn-Hilliard equations*, Discrete Contin. Dyn. Syst. Ser. A **34** (2014), 145-179.
- [26] H. Gajewski, K. Zacharias, *On a nonlocal phase separation model*, J. Math. Anal. Appl. **286** (2003), 11-31.
- [27] G. Giacomin, J.L. Lebowitz, *Phase segregation dynamics in particle systems with long range interactions. I. Macroscopic limits*, J. Statist. Phys. **87** (1997), 37-61.
- [28] G. Giacomin, J.L. Lebowitz, *Phase segregation dynamics in particle systems with long range interactions. II. Phase motion*, SIAM J. Appl. Math. **58** (1998), 1707-1729.
- [29] G. Giacomin, J.L. Lebowitz, R. Marra, *Macroscopic evolution of particle systems with short- and long-range interactions*, Nonlinearity **13** (2000), 2143-2162.
- [30] M. Grasselli, D. Pražák, *Longtime behavior of a diffuse interface model for binary fluid mixtures with shear dependent viscosity*, Interfaces Free Bound. **13** (2011), 507-530.
- [31] M.E. Gurtin, D. Polignone, J. Viñals, *Two-phase binary fluids and immiscible fluids described by an order parameter*, Math. Models Meth. Appl. Sci. **6** (1996), 8-15.
- [32] M. Heida, J. Málek, K.R. Rajagopal, *On the development and generalizations of Cahn-Hilliard equations within a thermodynamic framework*, Z. Angew. Math. Phys. **63** (2012), 145-169.
- [33] P.C. Hohenberg, B.I. Halperin, *Theory of dynamical critical phenomena*, Rev. Mod. Phys. **49** (1977), 435-479.
- [34] D. Jasnow, J. Viñals, *Coarse-grained description of thermo-capillary flow*, Phys. Fluids **8** (1996), 660-669.
- [35] J.S. Kim, *Phase-field models for multi-component fluid flows*, Commun. Comput. Phys., **12** (2012), 613-661.
- [36] N. Kim, L. Consiglieri, J.F. Rodrigues, *On non-Newtonian incompressible fluids with phase transitions*, Math. Methods Appl. Sci. **29** (2006), 1523-1541.
- [37] P. Krejčí, E. Rocca, J. Sprekels, *A nonlocal phase-field model with nonconstant specific heat*, Interfaces Free Bound., **9** (2007), 285-306.
- [38] P. Krejčí, E. Rocca, J. Sprekels, *Non-local temperature dependent phase-field model for non-isothermal phase transitions*, J. London Math. Soc., **76** (2007), 197-210.

- [39] A.G. Lamorgese, D. Molin, R. Mauri, *Phase field approach to multiphase flow modeling*, Milan J. Math. **79** (2011), 597-642.
- [40] S. Lisini, D. Matthes, G. Savaré, *Cahn-Hilliard and thin film equations with nonlinear mobility as gradient flows in weighted-Wasserstein metrics*, J. Differential Equations **253** (2012), 814-850.
- [41] C. Liu, J. Shen, *A phase field model for the mixture of two incompressible fluids and its approximation by a Fourier spectral method*, Phys. D **179** (2003), 211-228.
- [42] S.-O. Londen, H. Petzeltová, *Convergence of solutions of a non-local phase-field system*, Discrete Contin. Dyn. Syst. Ser. S **4** (2011), 653-670.
- [43] S.-O. Londen, H. Petzeltová, *Regularity and separation from potential barriers for a non-local phase-field system*, J. Math. Anal. Appl. **379** (2011), 724-735.
- [44] A. Morro, *Phase-field models of Cahn-Hilliard Fluids and extra fluxes*, Adv. Theor. Appl. Mech. **3** (2010), 409-424.
- [45] A. Novick-Cohen, *The Cahn-Hilliard equation*, Handbook of differential equations: evolutionary equations. Vol. IV, 201-228, Handb. Differ. Equ., Elsevier/North-Holland, Amsterdam, 2008.
- [46] J.S. Rowlinson, Translation of J.D. van der Waals, *The thermodynamic theory of capillarity under the hypothesis of a continuous variation of density*, J. Statist. Phys. **20** (1979), 197-244.
- [47] G. Schimperna, *Global attractors for Cahn-Hilliard equations with nonconstant mobility*, Nonlinearity **20** (2007), 2365-2387.
- [48] G. Schimperna, S. Zelik, *Existence of solutions and separation from singularities for a class of fourth order degenerate parabolic equations*, Trans. Amer. Math. Soc. **365** (2013), 3799-3829.
- [49] V.N. Starovoitov, *The dynamics of a two-component fluid in the presence of capillary forces*, Math. Notes **62** (1997), 244-254.
- [50] R. Temam, Navier-Stokes Equations. Theory and Numerical Analysis, North-Holland (Third edition), Oxford, 1984.
- [51] L. Zhao, H. Wu, H. Huang, *Convergence to equilibrium for a phase-field model for the mixture of two viscous incompressible fluids*, Commun. Math. Sci. **7** (2009), 939-962.
- [52] Y. Zhou, J. Fan, *The vanishing viscosity limit for a 2D Cahn-Hilliard-Navier-Stokes system with a slip boundary condition*, Nonlinear Anal. Real World Appl. **14** (2013), 1130-1134.