

FRONT PROPAGATION FOR NONLINEAR DIFFUSION EQUATIONS ON THE HYPERBOLIC SPACE

HIROSHI MATANO, FABIO PUNZO, AND ALBERTO TESEI

ABSTRACT. We study the Cauchy problem in the hyperbolic space \mathbb{H}^n ($n \geq 2$) for the semilinear heat equation with forcing term, which is either of KPP type or of Allen-Cahn type. Propagation and extinction of solutions, asymptotical speed of propagation and asymptotical symmetry of solutions are addressed. With respect to the corresponding problem in the Euclidean space \mathbb{R}^n new phenomena arise, which depend on the properties of the diffusion process in \mathbb{H}^n . We also investigate a family of travelling wave solutions, named horospheric waves, which have properties similar to those of plane waves in \mathbb{R}^n .

1. INTRODUCTION

In this paper we study the Cauchy problem for the semilinear parabolic equation

$$\boxed{\text{e1}} \quad (1.1) \quad \begin{cases} \frac{\partial u}{\partial t} = \Delta_H u + f(u) & \text{in } \mathbb{H}^n \times \mathbb{R}_+ \\ u = u_0 & \text{in } \mathbb{H}^n \times \{0\} \end{cases}$$

in the hyperbolic space \mathbb{H}^n ($n \geq 2$). Here Δ_H denotes the Laplace-Beltrami operator in \mathbb{H}^n and $\mathbb{R}_+ \equiv (0, \infty)$.

Concerning the Cauchy data function u_0 , the following assumption is always made in the sequel:

$$(A) \quad u_0 \text{ continuous in } \mathbb{H}^n, \quad 0 \leq u_0(x) \leq 1 \text{ for any } x \in \mathbb{H}^n.$$

As for the function f , we always assume:

$$(H_0) \quad f \in C^1([0, 1]), \quad f(0) = f(1) = 0.$$

Moreover, either

$$(H_1) \quad f'(0) > 0, f(u) > 0 \text{ for any } u \in (0, 1),$$

or

$$(H_2) \quad \begin{cases} (i) \text{ there exist } a \in (0, 1) \text{ such that} \\ \quad f(u) < 0 \text{ for any } u \in (0, a), f(u) > 0 \text{ for any } u \in (a, 1); \\ (ii) f'(0) < 0, \int_0^1 f(u) du > 0. \end{cases}$$

Following a common terminology, the function f is said to be "of KPP type" if it satisfies assumption (H_1) , or "of Allen-Cahn type" if (H_2) holds.

Date: June 27, 2012.

1991 Mathematics Subject Classification. 35K57, 35B40.

Key words and phrases. Semilinear parabolic equations; hyperbolic space; extinction and propagation; asymptotical symmetry of solutions; horospheric waves.

Solutions of problem (1.1) are always meant in the classical sense. By assumptions (A), (H_0) and comparison results [17], every solution u of problem (1.1) satisfies the inequality

$$\boxed{\text{estz}} \quad (1.2) \quad 0 \leq u(x, t) \leq 1 \quad \text{for any } (x, t) \in \mathbb{H}^n \times \mathbb{R}_+.$$

Under assumptions (A) and (H_0) a unique solution of problem (1.1) is easily seen to exist. In fact, existence follows by the a priori estimate (1.2) and standard compactness arguments, uniqueness by [17, Theorem 3.1].

The counterpart of problem (1.1) in \mathbb{R}^n , namely

$$\boxed{\text{e1i}} \quad (1.3) \quad \begin{cases} \frac{\partial u}{\partial t} = \Delta u + f(u) & \text{in } \mathbb{R}^n \times \mathbb{R}_+ \\ u = u_0 & \text{in } \mathbb{R}^n \times \{0\} \end{cases}$$

has been widely investigated (in particular, see [2, 3, 6, 9, 13]). Let us recall some well-known results (see [3] for details).

(a) If the forcing term f is of KPP type, then propagation always occurs - namely,

$$\boxed{\text{e2i}} \quad (1.4) \quad \lim_{t \rightarrow \infty} u(x, t) = 1 \quad \text{uniformly on compact subsets of } \mathbb{R}^n$$

for every solution $u \not\equiv 0$ of problem (1.3). This follows from the so-called "hair-trigger effect": if f satisfies (H_0) and

$$(HT) \quad \begin{cases} (i) \text{ there exists } a \in (0, 1] \text{ such that} \\ \quad f(u) > 0 \text{ for any } u \in (0, a); \\ (ii) \liminf_{u \rightarrow 0^+} u^{-(1+\frac{2}{n})} f(u) > 0, \end{cases}$$

then for every solution $u \not\equiv 0$ of problem (1.3)

$$\boxed{\text{propp}} \quad (1.5) \quad \liminf_{t \rightarrow \infty} u(x, t) \geq a \quad \text{uniformly on compact subsets of } \mathbb{R}^n.$$

Clearly, assumption (H_1) implies (HT) with $a = 1$, thus (1.4) follows. Observe that the exponent $p = 1 + \frac{2}{n}$ in (HT)-(ii) is the Fujita exponent of problem (1.3) with $f(u) = u^p$ (see [10]).

More generally, assumption (HT) is satisfied if (HT)-(i) holds and

$$\liminf_{u \rightarrow 0^+} u^{-p} f(u) > 0 \quad \text{for some } p \in \left(1, 1 + \frac{2}{n}\right),$$

thus (1.5) follows also in this case. On the other hand, if

$$\boxed{\text{exticondir}} \quad (1.6) \quad \limsup_{u \rightarrow 0^+} u^{-p} f(u) < \infty \quad \text{for some } p > 1 + \frac{2}{n}$$

and u_0 is small enough, then extinction occurs, namely

$$\boxed{\text{exti}} \quad (1.7) \quad \lim_{t \rightarrow \infty} u(x, t) = 0 \quad \text{uniformly in } \mathbb{R}^n.$$

(b) If f is of Allen-Cahn type, there is a "threshold effect". In fact, there is extinction (namely, equality (1.7) holds) if the initial data function u_0 is suitably small; instead, there is propagation (namely, equality (1.4) holds true) if u_0 is sufficiently large.

(c) Both for KPP and for Allen-Cahn there exists an "asymptotic speed of propagation" $c_0 > 0$, which is uniquely determined by the following properties:

(i) no solution with compact support of problem (1.3) can propagate with speed greater than c_0 . In fact, for any $c > c_0$ and $y \in \mathbb{R}^n$

$$\lim_{t \rightarrow \infty} \sup_{|x-y| > ct} u(x, t) = 0;$$

(ii) if a solution of problem (1.3) propagates, then its speed is no smaller than c_0 . In fact, if

$$\liminf_{t \rightarrow \infty} u(x, t) \geq a \quad \text{uniformly on compact subsets of } \mathbb{R}^n$$

for some $a \in (0, 1]$, then for any $c < c_0$ and $y \in \mathbb{R}^n$

$$\liminf_{t \rightarrow \infty} \inf_{|x-y| < ct} u(x, t) \geq a.$$

Observe that the asymptotic speed of propagation c_0 only depends on the forcing term f . In fact, its definition relies on a detailed investigation of the ordinary differential equation

$$\boxed{\text{e84}} \quad (1.8) \quad q'' + \kappa q' + f(q) = 0 \quad \text{in } \mathbb{R} \quad (\kappa \in \mathbb{R})$$

(in this connection, see Proposition 3.1 below). Equation (1.8) arises when seeking plane wave solutions of the equation

$$\boxed{\text{e92e}} \quad (1.9) \quad \frac{\partial u}{\partial t} = \Delta u + f(u) \quad \text{in } \mathbb{R}^n \times \mathbb{R}_+$$

- namely, solutions of equation (1.9) of the form

$$\boxed{\text{e7i}} \quad (1.10) \quad u(x, t) = q(\langle x, \nu \rangle - \kappa t) \quad (x \in \mathbb{R}^n, t > 0),$$

where q is a real function, $\nu \in \mathbb{R}^n$ is a fixed unit direction, $\kappa \in \mathbb{R}$ and $\langle x, \nu \rangle := \sum_{i=1}^n x_i \nu_i$.

Under the additional assumption

$$(H'_1) \quad \sup_{u \in (0,1]} \frac{f(u)}{u} = f'(0)$$

there holds $c_0 = 2\sqrt{f'(0)}$ (see [3, Proposition 4.2 and following remarks]).

In this paper we prove results for problem (1.1) which are analogous to those above for problem (1.3), yet exhibit remarkably novel features with respect to the Euclidean case. In fact, the following holds.

(a') If the function f is of KPP type, under the additional assumption (H'_1) we prove the existence of a new threshold effect. In fact, extinction prevails if

$$c_0 = 2\sqrt{f'(0)} < n - 1$$

and u_0 has compact support, whereas there is propagation if $c_0 > n - 1$ (see Theorem 3.2). Therefore, at variance from the Euclidean case, in the hyperbolic space we can have extinction even in the KPP case, depending on the sign of the difference $c_0 - (n - 1)$.

This implies that no hair-trigger effect holds in \mathbb{H}^n ; for, otherwise no extinction could arise for the KPP case, which is a contradiction when $c_0 < n - 1$. In this connection, observe that the proof of the hair-trigger effect in \mathbb{R}^n given in [3] relies on the Fujita phenomenon, whereas it is known that the Fujita exponent of problem (1.1) is $p = 1$ (see [4]) - namely, global solutions of problem (1.1) with $f(u) = u^p$

exist for any $p > 1$, if u_0 is sufficiently small. This remark also explains Theorem 3.3, which shows that every solution of problem (1.1) is extinct, if

$$\boxed{\text{exticondih}} \quad (1.11) \quad \limsup_{u \rightarrow 0^+} u^{-p} f(u) < \infty \quad \text{for some } p > 1$$

and u_0 is sufficiently small (condition (1.11) should be compared with (1.6) of the Euclidean case).

(*b'*) If f is of Allen-Cahn type, we prove extinction if the initial data function u_0 is sufficiently small, respectively propagation if u_0 is sufficiently large and $c_0 > n - 1$ (see Theorems 3.4 and 3.5). Hence the sign of the difference $c_0 - (n - 1)$ plays a role also in this case.

(*c'*) The role of the difference $c_0 - (n - 1)$ becomes clear by addressing the asymptotical speed of propagation of problem (1.1). In fact, in the hyperbolic space this speed (both for the KPP and the Allen-Cahn case) turns out to be $c_0 - (n - 1)$ (see Theorem 3.7).

A first heuristic explanation of the above differences can be given comparing the expression in polar coordinates (r, θ) in \mathbb{R}^n of equation (1.9) with that in polar geodesic coordinates (ρ, θ) in \mathbb{H}^n of the equation

$$\boxed{\text{e92}} \quad (1.12) \quad \frac{\partial u}{\partial t} = \Delta_H u + f(u) \quad \text{in } \mathbb{H}^n \times \mathbb{R}_+.$$

We have

$$\boxed{\text{e92re}} \quad (1.13) \quad \frac{\partial u}{\partial t} = \frac{\partial^2 u}{\partial r^2} + \frac{n-1}{r} \frac{\partial u}{\partial r} + \frac{1}{r^2} \Delta_\theta u + f(u),$$

respectively

$$\boxed{\text{e92rh}} \quad (1.14) \quad \frac{\partial u}{\partial t} = \frac{\partial^2 u}{\partial \rho^2} + (n-1) \coth \rho \frac{\partial u}{\partial \rho} + \frac{1}{(\sinh \rho)^2} \Delta_\theta u + f(u)$$

(see Section 2; here Δ_θ denotes the Laplace-Beltrami operator on the $(n - 1)$ -dimensional sphere of \mathbb{R}^n). While the coefficient of the first order term in the right-hand side of (1.13) tends to zero as $r \rightarrow \infty$, the corresponding term in (1.14) tends to $n - 1$ as $\rho \rightarrow \infty$. Therefore, in \mathbb{H}^n there is a "drift from infinity" which must be overcome to have propagation. If propagation prevails, the resulting speed is $c_0 - (n - 1)$, and not c_0 as in \mathbb{R}^n where the "drift from infinity" is absent. This explains why extinction can arise in \mathbb{H}^n even in the KPP case, and why the condition $c_0 > n - 1$ is needed to have propagation in the Allen-Cahn case.

A related viewpoint is that the different geometrical properties of \mathbb{H}^n affect diffusion through the spectral properties of the Laplace-Beltrami operator and the growth estimates of its heat kernel. In this connection, observe that by (H'_1)

$$\boxed{\text{e303}} \quad (1.15) \quad c_0 = 2\sqrt{f'(0)} > n - 1 \quad \Leftrightarrow \quad f'(0) > \frac{(n-1)^2}{4} =: \lambda_1,$$

where λ_1 denotes the infimum of the L^2 -spectrum of the operator $-\Delta_H$ in \mathbb{H}^n . Therefore, the important role of the difference $c_0 - (n - 1)$ is related to the fact that the spectrum has positive infimum. In general terms, the positivity of λ_1 makes diffusion "stronger in \mathbb{H}^n than in \mathbb{R}^n ". This explains why the Fujita exponent is $p = 1$ instead of $p = 1 + \frac{2}{n}$, and why extinction more easily prevails on propagation than in the Euclidean case.

Let us point out some technical points to illustrate the above remarks:

- the condition $f'(0) > \lambda_1$ plays a role when constructing a suitable family of subsolutions to problem (1.1), which is needed to prove propagation for the KPP case (see the proof of Theorem 3.2);
- estimates from above of the heat kernel in \mathbb{H}^n are used to exhibit a family of supersolutions to problem (1.1), which is used to prove extinction in the Allen-Cahn case (see the proof of Theorem 3.5);
- the existence of a ground state - namely, of a positive solution of the equation

eigenvalue (1.16)
$$\Delta_H \phi + \lambda_1 \phi = 0 \quad \text{in } \mathbb{H}^n,$$

which is radial and infinitesimal as $\rho \rightarrow \infty$, plays an important role to prove extinction when (1.11) holds (see the proof of Theorem 3.3).

Beside the results outlined in (a')-(c') above, we shall address the asymptotical symmetry of solutions to problem (1.1) (see Theorem 3.7), extending to the present situation some results in [13]. In doing so, we make use of the Alexandrov reflection method, already used in the hyperbolic space in [14, 15].

Finally, we introduce and investigate a family of travelling wave solutions of equation (1.12), named horospheric waves, which in some respects are analogous to plane waves in \mathbb{R}^n . In this connection, observe that in the Poincaré half-space model \mathbb{U}^n of the hyperbolic space horospheres take the form $x_n = \text{constant}$ ($(x_1, \dots, x_n) \in \mathbb{U}^n$).

Horospheric waves are defined as solutions of problem (1.1), whose level set is for any $t \in \mathbb{R}_+$ a horosphere of the form

?? (1.17)
$$\{x \in \mathbb{U}^n \mid x_n = Ke^{-\kappa t}\} \quad (K > 0, \kappa \in \mathbb{R})$$

(see Definition 3.1). In Theorem 3.9 we prove that:

- (i) a solution of this kind exists (both for KPP and for Allen-Cahn) if $c_0 > n - 1$;
- (ii) in \mathbb{U}^n it has the following representation, formally analogous to (1.10) with $\nu \equiv (0, \dots, 0, 1)$:

e8i (1.18)
$$u(x_1, \dots, x_n, t) = q^*(-\log x_n - c^*t),$$

where $c^* := c_0 - (n - 1)$ and q^* is the solution of the equation

eqn1 (1.19)
$$q'' + c_0 q' + f(q) = 0$$

such that

$$\lim_{\xi \rightarrow -\infty} q^*(\xi) = 1, \quad \lim_{\xi \rightarrow \infty} q^*(\xi) = 0.$$

Interestingly, horospheric waves give an estimate of any solutions to problem (1.1) (see Definition ?? and Theorem 3.10), as plane waves do in the Euclidean case. However, it is an open problem whether, under suitable assumptions, any solution is a horospheric wave (as it is the case for solutions of problem (1.3) and plane waves in \mathbb{R}^n ; see [6]).

2. GEOMETRICAL BACKGROUND

gb

By definition, the hyperbolic space \mathbb{H}^n is the unique simply connected, complete n -dimensional Riemannian manifold with sectional curvature equal to -1 . The hyperbolic distance between any two points $x, y \in \mathbb{H}^n$ will be denoted by $d(x, y)$. At

any fixed point $x_0 \in \mathbb{H}^n$, the inner Riemannian product $\langle \cdot, \cdot \rangle_H$ is given by

$$\boxed{\text{e13}} \quad (2.1) \quad \langle \xi, \eta \rangle_H := \sum_{i,j=1}^n g_{ij} \xi_i \eta_j,$$

where (g_{ij}) denotes the Riemannian metric of \mathbb{H}^n and the vectors $\xi \equiv (\xi_1, \xi_2, \dots, \xi_n)$, $\eta \equiv (\eta_1, \eta_2, \dots, \eta_n)$ belong to the tangent space T_{x_0} .

As is well-known, in \mathbb{H}^n every geodesic is defined on the whole real line; moreover, there exists exactly one geodesic passing through any two different points of \mathbb{H}^n . A subset $A \subseteq \mathbb{H}^n$ is a hyperbolic subspace if it contains the entire geodesic passing through any two of its points (a hyperbolic subspace of codimension 1 is called a hyperbolic hyperplane). Moreover, a subset $K \subseteq \mathbb{H}^n$ is convex if for any $x, y \in K$ the geodesic arc joining x to y lies in K . By definition, the convex hull of $A \subseteq \mathbb{H}^n$ is the smallest convex set of \mathbb{H}^n containing A .

Let us recall the definition of reflection through a hyperplane $\pi \subseteq \mathbb{H}^n$ (e.g., see [14]). First observe that for any given $x \in \mathbb{H}^n$ there exists a unique point $x^* \in \pi$ such that $d(x, x^*) = \min_{y \in \pi} d(x, y)$. Let $\gamma : \mathbb{R} \rightarrow \mathbb{H}^n$ be the unique entire geodesic joining x to x^* such that $\gamma(0) = x^*$, $\gamma(t_0) = x$ for some $t_0 > 0$. Then the reflection R_π with respect to the hyperplane π is defined as follows:

$$\boxed{\text{ref1}} \quad (2.2) \quad R_\pi : \mathbb{H}^n \rightarrow \mathbb{H}^n, \quad R_\pi(x) := \gamma(-t_0) \quad (x \in \mathbb{H}^n).$$

Denote by $\partial\mathbb{H}^n$ the boundary of the hyperbolic space (see [5]); the points of $\partial\mathbb{H}^n$ can be regarded as the points at infinity of \mathbb{H}^n . By definition, a horosphere centered at $x_0 \in \partial\mathbb{H}^n$ is a closed hypersurface of \mathbb{H}^n orthogonal to all geodesic lines with endpoint x_0 . Observe that, for any fixed $x_0 \in \partial\mathbb{H}^n$, \mathbb{H}^n is the disjoint union of the horospheres with center x_0 .

Let us recall for convenience of the reader some properties of different models of \mathbb{H}^n , which will be used in the following (e.g., see [5, 8, 12]).

(a) Set $B_r := \{x \in \mathbb{R}^n \mid |x| < r\}$ ($r > 0$), $|\cdot|$ denoting the euclidian norm of \mathbb{R}^n . The Poincaré disk model \mathbb{D}^n is the unitary ball B_1 endowed with the Riemannian metric

$$\boxed{\text{e1a}} \quad (2.3) \quad g_{ij} := \frac{4}{(1 - |x|^2)^2} \delta_{ij} \quad (x \in \mathbb{D}^n; i, j = 1, \dots, n).$$

Let us mention for further reference that:

- (i) geodesics in \mathbb{D}^n are either diameters of \mathbb{D}^n or circles orthogonal to $\partial\mathbb{D}^n$;
- (ii) a subset $\pi \subseteq \mathbb{D}^n$ is a hyperbolic hyperplane if and only if it is the intersection of \mathbb{D}^n either with a hyperplane of \mathbb{R}^n through the origin, or with an $(n-1)$ -dimensional sphere orthogonal to $\partial\mathbb{D}^n$ (see [5]);
- (iii) in \mathbb{D}^n , a horosphere with center $x_0 \in \partial\mathbb{H}^n$ is an $(n-1)$ -dimensional sphere of \mathbb{R}^n contained in \mathbb{D}^n and tangent in x_0 to $\partial\mathbb{D}^n$.

For any fixed point $O \in \mathbb{H}^n$ and $x \in \mathbb{H}^n$ we can consider the geodesic coordinates (ρ, θ) of x , namely $\rho \equiv \rho(x) := d(x, O)$. In \mathbb{D}^n , taking O at the origin we have

$$\boxed{\text{e5}} \quad (2.4) \quad \rho(x) = \int_0^{|x|} \frac{2}{1-s^2} ds = \log \left(\frac{1+|x|}{1-|x|} \right),$$

thus

$$\boxed{\text{e6}} \quad (2.5) \quad |x| = \tanh \frac{\rho(x)}{2}, \quad \frac{2}{1-|x|^2} = 2 \left[\cosh \frac{\rho(x)}{2} \right]^2 \quad (x \in \mathbb{D}^n).$$

Therefore, defining

$$\mathfrak{B}_r := \{x \in \mathbb{H}^n \mid \rho(x) < r\} \quad (r > 0),$$

in \mathbb{D}^n there holds for any $r \in (0, 1)$

$$B_r = \mathfrak{B}_{\log(\frac{1+r}{1-r})}.$$

Let us recall that in \mathbb{D}^n there is an isometry of the form (see [1]):

$$\boxed{\text{e61}} \quad (2.6) \quad \tau_y : \mathbb{D}^n \rightarrow \mathbb{D}^n, \quad \tau_y(x) := \frac{(1 - |y|^2)(x - y) - |x - y|^2 y}{(1 + |x|^2|y|^2 - 2 \sum_{i=1}^n x_i y_i)^2} \quad (x, y \in \mathbb{D}^n).$$

It is easily seen that $\tau_{-y} = \tau_y^{-1}$ ($y \in \mathbb{D}^n$). Moreover,

$$|\tau_y(x)| = \frac{|x - y|}{1 + |x|^2|y|^2 - 2 \sum_{i=1}^n x_i y_i} \quad (x, y \in \mathbb{D}^n).$$

For any $x, y \in \mathbb{D}^n$ we have

$$d(x, y) = d(\tau_x(x), \tau_x(y)) = d(0, \tau_x(y)) = \log \left(\frac{1 + |\tau_x(y)|}{1 - |\tau_x(y)|} \right).$$

Recall that the group of isometry of \mathbb{D}^n coincides with the group of conformal mappings of \mathbb{D}^n (see [5]). In particular, to prove Theorem 3.7 we shall use a particular class of isometries, namely the reflection through any hyperplane of \mathbb{H}^n .

The gradient $\nabla_H u \equiv ((\nabla_H u)_1, \dots, (\nabla_H u)_n)$ of a function $u \in C^1(\mathbb{H}^n)$ is given by

$$\boxed{\text{e11a}} \quad (2.7) \quad (\nabla_H u)_i := \sum_{j=1}^n g^{ij} \frac{\partial u}{\partial x_j} \quad (i = 1, \dots, n).$$

The Laplace-Beltrami operator in \mathbb{H}^n of a function $u \in C^2(\mathbb{H}^n)$ is

$$\boxed{\text{e2a}} \quad (2.8) \quad \Delta_H u \equiv \frac{1}{\sqrt{g}} \sum_{i=1}^n \frac{\partial}{\partial x_i} \left(\sqrt{g} \sum_{j=1}^n g^{ij} \frac{\partial u}{\partial x_j} \right),$$

where $g := \det(g_{ij})$, $g^{ij} := (g_{ij})^{-1}$. As already mentioned, the infimum of the L^2 -spectrum of the operator $-\Delta_H$ in \mathbb{H}^n is

$$\lambda_1 := \frac{(n-1)^2}{4}.$$

Moreover,

$$\boxed{\text{e28}} \quad (2.9) \quad \lambda_1(\mathfrak{B}_r) \searrow \lambda_1 \quad \text{as } r \rightarrow \infty$$

(see [11]), where $\lambda_1(\mathfrak{B}_r)$ is the first eigenvalue for $-\Delta_H$ in \mathfrak{B}_r with Dirichlet zero boundary conditions:

$$\boxed{\text{pre29}} \quad (2.10) \quad \begin{cases} \Delta_H \varphi_1 + \lambda_1(\mathfrak{B}_r) \varphi_1 = 0 & \text{in } \mathfrak{B}_r \\ \varphi_1 = 0 & \text{on } \partial \mathfrak{B}_r \quad (r > 0), \end{cases}$$

and $\varphi_1 = \varphi_1(\rho) > 0$ denotes the corresponding eigenfunction.

Let us recall that Δ_H commutes with isometries τ of \mathbb{H}^n (e.g., see [14]). Hence, if u is a solution of the equation

$$\frac{\partial u}{\partial t} = \Delta_H u + f(u) \quad \text{in } \mathbb{H}^n$$

then $v := u \circ \tau$ satisfies

$$\frac{\partial v}{\partial t} = \Delta_H v + f(v) \quad \text{in } \mathbb{H}^n.$$

By (2.3) and (2.8), the Laplace-Beltrami operator in \mathbb{D}^n has the expression

$$\boxed{\text{e5a}} \quad (2.11) \quad \Delta_H u = \frac{1}{4}(1-|x|^2)^2 \sum_{i=1}^n \frac{\partial^2 u}{\partial x_i^2} + \frac{n-2}{2}(1-|x|^2) \sum_{i=1}^n x_i \frac{\partial u}{\partial x_i},$$

thus it can be regarded as a linear elliptic operator on $B_1 \subseteq \mathbb{R}^n$ with bounded coefficients degenerating at the boundary ∂B_1 . Using polar geodesic coordinates (ρ, θ) in \mathbb{D}^n , thus

$$ds^2 := \sum_{i,j=1}^n g_{ij} dx_i dx_j = d\rho^2 + (\sinh \rho)^2 d\theta,$$

we obtain

$$\boxed{\text{e9}} \quad (2.12) \quad \Delta_H u = \frac{\partial^2 u}{\partial \rho^2} + (n-1) \coth \rho \frac{\partial u}{\partial \rho} + \frac{1}{(\sinh \rho)^2} \Delta_\theta u,$$

Δ_θ being the Laplace-Beltrami operator on the $(n-1)$ -dimensional sphere of \mathbb{R}^n .

In \mathbb{H}^n the heat kernel $G : \mathbb{H}^n \times \mathbb{H}^n \times \mathbb{R}_+ \rightarrow \mathbb{R}_+$ is well defined (see [7] and references therein for its explicit construction and main properties). Then the unique bounded solution of the Cauchy problem

$$\begin{cases} \frac{\partial u}{\partial t} = \Delta_H u & \text{in } \mathbb{H}^n \times \mathbb{R}_+ \\ u = u_0 & \text{in } \mathbb{H}^n \times \{0\} \end{cases}$$

with Cauchy data u_0 bounded in \mathbb{H}^n is

$$\boxed{\text{e70}} \quad (2.13) \quad u(x, t) := \int_{\mathbb{H}^n} G(x, y, t) u_0(y) d\mu_y \quad ((x, t) \in \mathbb{H}^n \times \mathbb{R}_+),$$

where $d\mu_y$ denotes the volume element of \mathbb{H}^n .

It is known [7] that there exists $c_n > 0$ such that for any $x, y \in \mathbb{H}^n$ and $t \in \mathbb{R}_+$

$$\boxed{\text{e71}} \quad (2.14) \quad \frac{1}{c_n} h_n(d(x, y), t) \leq G(x, y, t) \leq c_n h_n(d(x, y), t),$$

where

$$h_n(d, t) := (4\pi t)^{-\frac{n}{2}} (1+d)(1+d+t)^{\frac{n-3}{2}} e^{-\lambda_1 t - \frac{n-1}{2}d - \frac{d^2}{4t}} \quad (d \geq 0, t > 0).$$

Hence there exists $C_n > 0$ such that for any $t \in \mathbb{R}_+$

$$\boxed{\text{e72}} \quad (2.15) \quad \sup_{x, y \in \mathbb{H}^n} G(x, y, t) \leq C_n \frac{(1+t)^{\frac{n-3}{2}}}{t^{\frac{n}{2}}} e^{-\lambda_1 t}.$$

(b) The hyperboloid model of \mathbb{H}^n , denoted by \mathbb{I}^n , is given by the hyperboloid

$$\{(x_1, \dots, x_n, x_{n+1}) \in \mathbb{R}^{n+1} \mid x_1^2 + \dots + x_n^2 - x_{n+1}^2 = -1, x_{n+1} > 0\}$$

with the metric induced on its tangent bundle by the Euclidean metric of \mathbb{R}^{n+1} .

For any $x \in \mathbb{I}^n$ and y in the tangent space of \mathbb{I}^n at x , the geodesic starting from x with tangent vector y is given by

$$\gamma(t) := x \cosh t + y \sinh t \quad (t \in \mathbb{R}).$$

Observe that the map

$$\phi : \mathbb{D}^n \rightarrow \mathbb{I}^n, \quad \phi(x) := \frac{(2x, 1 + |x|^2)}{1 - |x|^2}$$

is bijective and isometric. Then the Klein model of \mathbb{H}^n , denoted by \mathbb{K}^n , is defined using the bijective map

$$\psi : B_1 \subseteq \mathbb{R}^n \rightarrow \mathbb{I}^n, \quad \psi(x) := \frac{(x, 1)}{\sqrt{1 - |x|^2}}$$

(from a geometrical viewpoint, the point $\psi(x) \in \mathbb{R}^{n+1}$, $x \in B_1$ is the intersection of \mathbb{I}^n with the line passing through the point $(x, 1) \in \mathbb{R}^{n+1}$ and the origin in \mathbb{R}^{n+1}). By definition, the Klein model \mathbb{K}^n is B_1 equipped with the metric obtained by transporting the hyperbolic metric of \mathbb{I}^n along ψ .

hvarie

Remark 2.1. Observe for further purposes that in \mathbb{K}^n :

- (i) geodesics are the traces of ordinary affine lines in \mathbb{K}^n (see [12]);
- (ii) for any convex subset A of \mathbb{K}^n and any point $x \in \mathbb{K}^n \setminus A$ there exists a hyperplane π such that $x \in \pi$ and $\pi \cap A = \emptyset$;
- (iii) for any convex subset A of \mathbb{K}^n and any entire geodesic $\gamma \subseteq \mathbb{K}^n$ which does not intersect A , there exists a hyperplane $\pi \subseteq \mathbb{K}^n$ such that $\gamma \subseteq \pi$ and $\pi \cap A = \emptyset$.

Furthermore, since the map $\phi^{-1} \circ \psi : \mathbb{K}^n \rightarrow \mathbb{D}^n$ is bijective and transforms geodesics of \mathbb{K}^n on those of \mathbb{D}^n , properties (ii)-(iii) hold true in \mathbb{D}^n , too.

(c) Finally, consider the Poincaré half-space model of the hyperbolic space, denoted by \mathbb{U}^n . This is the upper half-space $\{x \in \mathbb{R}^n \mid x_n > 0\}$ endowed with the metric

$$ds^2 := \frac{1}{x_n^2} \sum_{i=1}^n dx_i^2.$$

Observe that an inversion in a sphere with radius $\sqrt{2}$ and center in the south pole of \mathbb{D}^n maps \mathbb{U}^n onto \mathbb{D}^n (see [12]). Therefore \mathbb{U}^n is isometric to \mathbb{D}^n , thus to \mathbb{I}^n .

For any $\bar{x} \in \partial\mathbb{U}^n = \{x \in \mathbb{R}^n \mid x_n = 0\} \cup \{\infty\}$, the horospheres with center at \bar{x} are either hyperplanes of equation $x_n = \xi$ for some $\xi > 0$, if $\bar{x} = \{\infty\}$, or spheres of \mathbb{R}^n contained in \mathbb{U}^n tangent in \bar{x} to $\{x \in \mathbb{R}^n \mid x_n = 0\}$, if $\bar{x}_n = 0$ (e.g., see [5]). Clearly, for any $\bar{x} \in \partial\mathbb{U}^n$ with $\bar{x}_n = 0$, a horosphere with center at \bar{x} can be isometrically mapped into a horosphere with center at ∞ .

For any $x = (x_1, \dots, x_n), y = (y_1, \dots, y_n) \in \mathbb{U}^n$ there holds

distsp

$$(2.16) \quad d(x, y) = \cosh^{-1} \left[1 + \frac{(x_1 - y_1)^2 + \dots + (x_n - y_n)^2}{2x_n y_n} \right].$$

The Laplace-Beltrami operator on \mathbb{U}^n can be expressed as follows:

e727

$$(2.17) \quad \Delta_H u = x_n^2 \Delta u + (2 - n)x_n \frac{\partial u}{\partial x_n},$$

where Δ denotes the Laplacian in \mathbb{R}^n .

3. MAIN RESULTS

mr

Let us first recall the following result concerning equation (1.8) (see [3, Theorem 4.1 and Lemma 4.3]).

awresults

Proposition 3.1. *Let assumptions (H_0) and either (H_1) or (H_2) be satisfied. Then there exists $c_0 > 0$ with the following properties:*

- (i) for $\kappa = c_0$ equation (1.8) admits a decreasing solution q^* in \mathbb{R} satisfying

e85

$$(3.1) \quad \lim_{\xi \rightarrow -\infty} q^*(\xi) = 1, \quad \lim_{\xi \rightarrow \infty} q^*(\xi) = 0;$$

(ii) for any $\kappa \in (0, c_0)$ there exists $\gamma_\kappa \in (0, 1)$ such that: for any $\eta \in (\gamma_\kappa, 1)$ there exist $b = b(\kappa, \eta) > 0$ and a solution q to equation (1.8) satisfying

$$\boxed{\text{e86}} \quad (3.2) \quad q(0) = \eta, \quad q'(0) = 0, \quad q(b) = 0, \quad q' < 0 \text{ in } (0, b];$$

(iii) for any $\kappa \in (c_0, \infty)$ there exists a solution q to equation (1.8) in \mathbb{R}_+ such that

$$\boxed{\text{e87}} \quad (3.3) \quad q(0) = 1, \quad q' < 0 \text{ in } \mathbb{R}_+, \quad q(\xi) \rightarrow 0 \text{ as } \xi \rightarrow \infty.$$

3.1. Extinction versus propagation. If the function f is of KPP type, the following holds.

$\boxed{\text{t1}}$ **Theorem 3.2.** *Let assumptions (H_0) , (H_1) and (H'_1) be satisfied. Let $u_0 \not\equiv 0$, and let u be the corresponding solution of problem (1.1).*

(i) *Suppose that u_0 has compact support and $c_0 < n - 1$. Then*

$$\boxed{\text{e89}} \quad (3.4) \quad \lim_{t \rightarrow \infty} u(x, t) = 0 \quad \text{uniformly in } \mathbb{H}^n.$$

(ii) *Suppose that $c_0 > n - 1$. Then*

$$\boxed{\text{e90}} \quad (3.5) \quad \lim_{t \rightarrow \infty} u(x, t) = 1 \quad \text{uniformly on compact subsets of } \mathbb{H}^n.$$

Observe that a function f of KPP type cannot satisfy condition (1.11). If (1.11) holds, we have the following result.

$\boxed{\text{tblup}}$ **Theorem 3.3.** *Let assumptions (H_0) and (1.11) be satisfied, and let u_0 be sufficiently small. Then equality (3.4) holds for the corresponding solution of problem (1.1).*

Remark 3.1. By “ u_0 sufficiently small” in the above statement we mean $u_0(x) \leq w(\rho(x))$ for any $x \in \mathbb{H}^n$, the function $w = w(\rho)$ being defined in (4.19).

Results analogous to those of Theorem 3.2 hold, if f is of Allen-Cahn type and suitable assumptions on the size of u_0 are made. In fact, the following holds.

$\boxed{\text{t6}}$ **Theorem 3.4.** *Let assumptions (H_0) and (H_2) be satisfied. Let u be the solution of problem (1.1).*

(i) *If $\sup_{\mathbb{H}^n} u_0 < a$, then equality (3.4) holds.*

(ii) *Suppose that u_0 is suitably large and $c_0 > n - 1$. Then equality (3.5) holds.*

Remark 3.2. By “ u_0 suitably large” in the above statement we mean $u_0 \geq v_0$ in \mathbb{H}^n , the function v_0 being defined in (4.20).

The conclusion of Theorem 3.4-(i) holds true, if (H_2) is replaced by the weaker assumption:

$$(H_4) \quad \begin{cases} \text{there exists } a \in (0, 1) \text{ such that } f(u) \leq 0 \text{ for any } u \in [0, a], \\ f(\bar{u}) > 0 \text{ for some } \bar{u} \in (a, 1). \end{cases}$$

In fact, set

$$\boxed{\text{mueta}} \quad (3.6) \quad \sigma \equiv \sigma(\eta) := \sup_{u \in (\eta, 1)} \frac{f(u)}{u - \eta}$$

for any fixed $\eta \in [0, a)$. Then we have the following result, which is the counterpart of [3, Theorem 6.1].

t7 **Theorem 3.5.** *Let assumptions (H_0) and (H_4) be satisfied. Let u be a solution to problem (1.1). Let $\eta \in [0, a)$ be fixed and either of the following assumptions be satisfied:*

- (i) $\sigma(\eta) \leq \lambda_1$;
- (ii) $\lambda_1 < \sigma(\eta) < \infty$ and there exists $t_0 > 0$ such that

$$\text{e63} \quad (3.7) \quad \int_{\mathbb{H}^n} [u_0(y) - \eta]_+ d\mu_y \leq \frac{t_0^{\frac{n}{2}}}{C_n(1+t_0)^{\frac{n-3}{2}}} e^{[\lambda_1 - \sigma(\eta)]t_0} (a - \eta),$$

where $C_n > 0$ is the constant in inequality (2.15).

Then

$$\text{e120} \quad (3.8) \quad \limsup_{t \rightarrow \infty} u(x, t) \leq \eta \quad \text{uniformly in } \mathbb{H}^n.$$

Moreover, if $f(u) < 0$ for any $u \in (0, \eta]$, then equality (3.4) holds.

3.2. Speed of propagation and asymptotical symmetry. Let us now address the case of propagation. Concerning the spreading speed, we have the following result.

t3 **Theorem 3.6.** *Let assumptions (H_0) , and either (H_1) - (H'_1) or (H_2) be satisfied. Let $u_0 \not\equiv 0$ have compact support, and let u_0 be suitably large if (H_2) holds. Moreover, assume $c_0 > n - 1$.*

- (i) Let $c > c_0 - (n - 1)$. Then for every $y \in \mathbb{H}^n$

$$\text{e20} \quad (3.9) \quad \lim_{t \rightarrow \infty} \sup_{d(x, y) > ct} u(x, t) = 0.$$

- (ii) Let $0 < c < c_0 - (n - 1)$. Then for every $y \in \mathbb{H}^n$

$$\text{e20a} \quad (3.10) \quad \lim_{t \rightarrow \infty} \inf_{d(x, y) < ct} u(x, t) = 1.$$

Let us also prove some geometrical properties of the level sets of the solution of problem (1.1), which are the counterpart of [13, Theorem] in \mathbb{R}^n . This is the content of the following

t5 **Theorem 3.7.** *Let the assumptions of Theorem 3.6 be satisfied. Then for any $a \in (0, 1)$ and $t \in \mathbb{R}_+$ sufficiently large the following holds:*

- (i) the level set

$$\text{levelset} \quad (3.11) \quad \Gamma_a(u; t) := \{x \in \mathbb{H}^n \mid u(x, t) = a\} \quad (t \in \mathbb{R}_+)$$

is a smooth $(n - 1)$ -dimensional submanifold of \mathbb{H}^n ;

- (ii) every geodesic orthogonal to $\Gamma_a(u; t)$ intersects the convex hull of the support of u_0 .

Since the size of $\Gamma_a(u; t)$ tends to infinity as $t \rightarrow \infty$, while that of $\text{supp } u_0$ remains unchanged, the above theorem implies, roughly, that the shape of the expanding front $\Gamma_a(u; t)$ converges to that of a sphere, at least in the C^1 sense. Furthermore, one can derive from the above theorem the following corollary.

Corollary 3.8. *Let the assumptions of Theorem 3.6 be satisfied. Fix a point $P \in \text{supp } u_0$, and let $\rho_{\min}(t)$ (respectively $\rho_{\max}(t)$) denote the minimal (respectively maximal) geodesic distance from P to $\Gamma_a(u; t)$. Then there exists a constant $M > 0$ such that*

$$\rho_{\max}(t) - \rho_{\min}(t) \leq M \quad \text{for all } t \geq 0.$$

how

3.3. Horospheric waves. Let us first state the following definition.

defhw

Definition 3.1. By a horospheric wave solution of problem (1.1) we mean any solution whose level sets $\Gamma_a(u; t)$ ($a \in (0, 1)$, $t \in \mathbb{R}_+$) are horospheres of the form (1.17):

$$\Gamma_a(u; t) = \{x \in \mathbb{U}^n \mid x_n = K_a e^{-\kappa t}\} \quad (K_a > 0, \kappa \in \mathbb{R})$$

in the Poincaré half-space model \mathbb{U}^n .

Concerning existence of horospheric waves, the following result will be proven.

pexhw

Theorem 3.9. *Let assumptions (H_0) and either (H_1) or (H_2) be satisfied. Suppose $c_0 > n - 1$. Then there exists a horospheric wave, which in the Poincaré half-space model \mathbb{U}^n takes the form (1.18):*

$$u(x_1, \dots, x_n, t) := q^*(-\log x_n - c^* t) \quad ((x_1, \dots, x_n, t) \in \mathbb{U}^n \times \mathbb{R}_+).$$

Observe that the existence of the profile function q^* is ensured by Proposition 3.1-(i). The quantity $c^* := c_0 - (n - 1)$ is called the speed of the horospheric wave.

As already remarked, the above expression of the horospheric wave in \mathbb{U}^n is formally analogous to that of a planar wave in \mathbb{R}^n . This analogy suggests to express solutions of the problem

e77r

$$(3.12) \quad \begin{cases} \frac{\partial u}{\partial t} = \Delta_H u + f(u) & \text{in } \mathbb{U}^n \times \mathbb{R}_+ \\ u = u_0 & \text{in } \mathbb{U}^n \times \{0\} \end{cases}$$

in a moving frame, so that horospheric waves appear as stationary states - namely, to introduce the new coordinate $e^{c^* t} x_n$ ($t \in \mathbb{R}_+$; see Section 6).

Remarkably, horospheric waves provide upper and lower bounds for large times of any solution of problem (3.12), as plane waves in \mathbb{R}^n do for solutions of problem (1.3). To address this point, let us think of the function f as defined on \mathbb{R} and satisfying the following assumption (which implies both (H_0) and (H_2)):

$$(H_5) \quad \begin{cases} (i) f \in C^1(\mathbb{R}), f(0) = f(1) = 0, \\ f'(0) < 0, f'(1) < 0, \int_0^1 f(u) du > 0; \\ (ii) \text{ there exists } a \in (0, 1) \text{ such that } f(u) > 0 \text{ if } u \in (-\infty, 0) \cup (a, 1), \\ f(u) < 0 \text{ if } u \in (0, a) \cup (1, \infty). \end{cases}$$

Then the following result can be proven.

t9

Theorem 3.10. *Let assumption (H_5) be satisfied, and let u be any solution of problem (3.12). Suppose that*

cinz

$$(3.13) \quad \limsup_{x_n \rightarrow 0^+} \sup_{x' \in \mathbb{R}^{n-1}} u_0(x', x_n) < a, \quad \liminf_{x_n \rightarrow \infty} \inf_{x' \in \mathbb{R}^{n-1}} u_0(x', x_n) > a.$$

Then there exist $k_*, k^* \in \mathbb{R}_+$ such that

e79r6

$$(3.14) \quad q^*(-\log(k_* x_n) - c^* t) \leq u(x', x_n, t) \leq q^*(-\log(k^* x_n) - c^* t)$$

for every $(x', x_n) \in \mathbb{U}^n$, $t \in \mathbb{R}_+$.

Remark 3.3. For each given direction $\theta \in S^{n-1}$ in the Poincaré disk model we can choose the direction of the x_n -axis along this θ direction, and express the solution in the half-space model \mathbb{U}^n with respect to this choice of the x_n -axis. By this choice,

we can properly define “the ω -limit set in the direction θ ” of any solution u of problem (3.12).

By assumptions (A) and (H_5) -(i), using standard comparison and compactness results it is easily seen that, for any $\theta \in S^{n-1}$, the ω -limit set of u in the direction θ is nonempty (e.g., see [16] for details). If $\tilde{u} = \tilde{u}(x', x_n, t)$ is any point of this set, by inequality (3.14) we have

$$q^*(-\log(k_* x_n) - c^* t) \leq \tilde{u}(x', x_n, t) \leq q^*(-\log(k^* x_n) - c^* t) \quad ((x', x_n) \in \mathbb{U}^n, t \in \mathbb{R}_+).$$

However, it is an open problem whether every ω -limit point \tilde{u} itself is a horospheric wave (see [6, Theorem 3.1] for the analogous result concerning plane waves in \mathbb{R}^n).

4. EXTINCTION VERSUS PROPAGATION: PROOFS

Let us first prove Theorem 3.2.

Proof of Theorem 3.2. (i) By assumption, there exists $R > 0$ such that $\text{supp } u_0 \subseteq \mathfrak{B}_R$. Hence there exists a smooth function $\tilde{u}_0 : \mathbb{R}_+ \rightarrow [0, 1]$, $\tilde{u}_0 = \tilde{u}_0(\rho)$, $\tilde{u}'_0 \leq 0$ in \mathbb{R}_+ , $\tilde{u}_0(\rho) = 0$ for any $\rho \geq R$, such that

$$\boxed{\text{e21}} \quad (4.1) \quad u_0(x) \leq \tilde{u}_0(\rho(x)) \quad \text{for any } x \in \mathbb{H}^n.$$

Let \tilde{u} be the solution of the problem

$$\boxed{\text{e24}} \quad (4.2) \quad \begin{cases} \frac{\partial u}{\partial t} = \frac{\partial^2 u}{\partial \rho^2} + (n-1) \coth \rho \frac{\partial u}{\partial \rho} + f(u) & \text{in } \mathbb{R}_+ \times \mathbb{R}_+ \\ \frac{\partial u}{\partial \rho} = 0 & \text{in } \{0\} \times \mathbb{R}_+ \\ u = \tilde{u}_0 & \text{in } \mathbb{R}_+ \times \{0\}. \end{cases}$$

Then $\tilde{u}(\rho(x), t)$ solves problem (1.1) with Cauchy data $\tilde{u}_0(\rho(x))$ (see (2.12)). By (4.1) and comparison results (see [17]) we obtain

$$\boxed{\text{e25}} \quad (4.3) \quad 0 \leq u(x, t) \leq \tilde{u}(\rho(x), t) \quad \text{for any } (x, t) \in \mathbb{H}^n \times \mathbb{R}_+.$$

On the other hand, consider the problem

$$\boxed{\text{e22}} \quad (4.4) \quad \begin{cases} \frac{\partial w}{\partial t} = \frac{\partial^2 w}{\partial \rho^2} + (n-1) \frac{\partial w}{\partial \rho} + f(w) & \text{in } \mathbb{R}_+ \times \mathbb{R}_+ \\ \frac{\partial w}{\partial \rho} = 0 & \text{in } \{0\} \times \mathbb{R}_+ \\ w = \tilde{u}_0 & \text{in } \mathbb{R}_+ \times \{0\}. \end{cases}$$

We shall prove the following

Claim 1: Let $c_0 < n - 1$. Then the solution w of problem (4.4) satisfies

$$\boxed{\text{e23}} \quad (4.5) \quad \sup_{\rho \in \mathbb{R}_+} w(\rho, t) \rightarrow 0 \quad \text{as } t \rightarrow \infty,$$

$$\boxed{\text{e26}} \quad (4.6) \quad \frac{\partial w}{\partial \rho} \leq 0 \quad \text{in } \mathbb{R}_+ \times \mathbb{R}_+.$$

From the above claim the conclusion easily follows. In fact, since $\coth \rho \geq 1$ and $\frac{\partial w}{\partial \rho} \leq 0$, w is a supersolution of problem (4.2). Therefore, by comparison results

$$0 \leq \tilde{u}(\rho, t) \leq w(\rho, t) \quad \text{in } \mathbb{R}_+ \times \mathbb{R}_+,$$

whence by (4.3) and (4.5)

$$0 \leq \sup_{x \in \mathbb{H}^n} u(x, t) \leq \sup_{\rho \in \mathbb{R}_+} \tilde{u}(\rho, t) \rightarrow 0 \quad \text{as } t \rightarrow \infty.$$

It remains to prove Claim 1. To this purpose observe that, since by assumption $c_0 < n - 1$, by Proposition 3.1-(iii) the ordinary differential equation

$$q'' + (n - 1)q' + f(q) = 0$$

has a solution $q = q(\xi)$ in \mathbb{R}_+ such that

$$q(0) = 1, \quad q' < 0 \quad \text{in } \mathbb{R}_+, \quad q(\xi) \rightarrow 0 \quad \text{as } \xi \rightarrow \infty.$$

Set

$$\boxed{\text{e261}} \quad (4.7) \quad \varphi(\xi) := \begin{cases} 1 & \text{if } \xi < R \\ q(\xi - R) & \text{if } \xi \geq R. \end{cases}$$

Let v be the solution of the problem

$$\boxed{\text{e262}} \quad (4.8) \quad \begin{cases} \frac{\partial v}{\partial t} = \frac{\partial^2 v}{\partial \xi^2} + (n - 1) \frac{\partial v}{\partial \xi} + f(v) & \text{in } \mathbb{R} \times \mathbb{R}_+ \\ v = \varphi & \text{in } \mathbb{R} \times \{0\}. \end{cases}$$

The proof of [3, Theorem 5.1] shows that

$$\boxed{\text{e57a}} \quad (4.9) \quad \frac{\partial v}{\partial \xi} \leq 0 \quad \text{in } \mathbb{R} \times \mathbb{R}_+ \quad \text{and} \quad \lim_{t \rightarrow \infty} v(\xi, t) = 0$$

for any $\xi \in \mathbb{R}$. Since $\tilde{u}_0 \leq \varphi$ in \mathbb{R}_+ , by the inequality in (4.9) v is a supersolution of problem (4.4). Therefore $w \leq v$ in $\mathbb{R}_+ \times \mathbb{R}_+$. This fact and (4.9) yield (4.5).

To prove (4.6), consider the problem

$$\begin{cases} \frac{\partial z}{\partial t} = \frac{\partial^2 z}{\partial \rho^2} + (n - 1) \frac{\partial z}{\partial \rho} + f'(w)z & \text{in } \mathbb{R}_+ \times \mathbb{R}_+ \\ z = 0 & \text{in } \{0\} \times \mathbb{R}_+ \\ z = 0 & \text{in } \mathbb{R}_+ \times \{0\}, \end{cases}$$

where w is the solution of problem (4.4). Since by assumption $\tilde{u}'_0 \leq 0$, the function $\frac{\partial w}{\partial \rho}$ is a subsolution of the above problem, whereas $z \equiv 0$ is a solution. Hence by comparison (4.6) follows. This completes the proof of Claim 1, thus of (i).

(ii) Under the present assumptions inequality (1.15) holds, λ_1 denoting the infimum of the L^2 -spectrum of the operator $-\Delta_H$ in \mathbb{H}^n . Consider the eigenvalue problem

$$\boxed{\text{e29}} \quad (4.10) \quad \begin{cases} \Delta_H \varphi + [f'(0) + \mu]\varphi = 0 & \text{in } \mathfrak{B}_R \\ \varphi = 0 & \text{on } \partial \mathfrak{B}_R \quad (R > 0). \end{cases}$$

Denote by $\mu_1 = \mu_1(\mathfrak{B}_R)$ the first eigenvalue of (4.10) and by $\varphi_1 = \varphi_1(\rho) > 0$ the corresponding eigenfunction. Clearly,

$$\mu_1(\mathfrak{B}_R) = \lambda_1(\mathfrak{B}_R) - f'(0)$$

(see (2.10)), where $\lambda_1(\mathfrak{B}_r)$ denotes the first eigenvalue of $-\Delta_H$ in \mathfrak{B}_r with Dirichlet zero boundary conditions. Hence by (1.15) and (2.9) there exists $R_0 > 0$ such that $\mu_1(\mathfrak{B}_R) < 0$ for any $R > R_0$.

Set

$$\boxed{\text{e31}} \quad (4.11) \quad w_\varepsilon(\rho) := \begin{cases} \varepsilon \varphi_1(\rho) & \text{in } \mathfrak{B}_R \\ 0 & \text{otherwise,} \end{cases}$$

with $R > R_0$ fixed. Since $\mu_1(\mathfrak{B}_R) < 0$, it is easily seen that there exists $\varepsilon_0 > 0$ such that for any $\varepsilon \in (0, \varepsilon_0)$ the function w_ε is a subsolution of the equation

$$\boxed{\text{e33}} \quad (4.12) \quad \Delta_H v + f(v) = 0 \quad \text{in } \mathbb{H}^n.$$

On the other hand, every solution $u = u(x, t)$ of problem (1.1) with Cauchy data $u_0 \not\equiv 0$ satisfies $u(\cdot, \bar{t}) > 0$ in \mathbb{H}^n for any $\bar{t} \in \mathbb{R}_+$ (this is a standard consequence of the strong maximum principle; see the form (2.11) of the operator Δ_H in \mathbb{D}^n). Choose $\varepsilon \in (0, \varepsilon_0)$ so small that

$$w_\varepsilon(\rho(x)) \leq u(x, \bar{t}) \quad \text{for any } x \in \mathbb{H}^n.$$

Then by comparison arguments

$$\boxed{\text{e34}} \quad (4.13) \quad u_\varepsilon(\cdot, t) \leq u(\cdot, t + \bar{t}) \leq 1 \quad \text{in } \mathbb{H}^n$$

for any $t \in \mathbb{R}_+$, where u_ε denotes the solution of the problem

$$\boxed{\text{e32}} \quad (4.14) \quad \begin{cases} \frac{\partial u}{\partial t} = \Delta_H u + f(u) & \text{in } \mathbb{H}^n \times \mathbb{R}_+ \\ u = w_\varepsilon & \text{in } \mathbb{H}^n \times \{0\}. \end{cases}$$

By standard arguments,

- (a) the function $t \mapsto u_\varepsilon(x, t)$ is nondecreasing in \mathbb{R}_+ for any $x \in \mathbb{H}^n$;
- (b) its pointwise limit

$$u_\infty(x) := \lim_{t \rightarrow \infty} u_\varepsilon(x, t) \quad (x \in \mathbb{H}^n)$$

is a solution of equation (4.12);

- (c) the convergence $u_\varepsilon(\cdot, t) \rightarrow u_\infty$ as $t \rightarrow \infty$ is uniform on compact subsets of \mathbb{H}^n .

Therefore the conclusion will follow from inequality (4.13), if we prove that

$$\boxed{\text{e37}} \quad (4.15) \quad u_\infty \equiv 1 \quad \text{in } \mathbb{H}^n.$$

To this purpose, choose $\varepsilon \in (0, \varepsilon_0)$ so small that

$$\boxed{\text{e373}} \quad (4.16) \quad 0 \leq w_\varepsilon(x) < u_\infty(x) \quad \text{for any } x \in \mathbb{H}^n.$$

We shall prove the following

Claim 2: There holds

$$B := \{y \in \mathbb{H}^n \mid w_\varepsilon(\tau_y(x)) < u_\infty(x) \text{ for any } x \in \mathbb{H}^n\} = \mathbb{H}^n,$$

where $\tau_y : \mathbb{H}^n \rightarrow \mathbb{H}^n$ is the continuous family of isometries defined in (2.6).

From the above claim equality (4.15) follows easily. In fact, since $B = \mathbb{H}^n$, we obtain in particular

$$w_\varepsilon(\tau_x(x)) = w_\varepsilon(0) < u_\infty(x) \quad \text{for any } x \in \mathbb{H}^n.$$

This implies

$$\boxed{\text{e68}} \quad (4.17) \quad \xi(t) \leq u_\infty(x) \quad \text{for any } x \in \mathbb{H}^n, t \in \mathbb{R}_+,$$

where ξ is the solution of the problem

$$\begin{cases} \xi' = f(\xi) \\ \xi(0) = w_\varepsilon(0). \end{cases}$$

Since $\xi(t) \rightarrow 1$ as $t \rightarrow \infty$, inequality (4.17) implies (4.15).

To prove Claim 2, observe preliminarily that $0 \in B$. In fact, by (4.16)

$$w_\varepsilon(\tau_0(x)) = w_\varepsilon(x) < u_\infty(x) \quad \text{for any } x \in \mathbb{H}^n.$$

By the continuity of the map $y \rightarrow \tau_y(x)$ for any fixed $x \in \mathbb{H}^n$ (see (2.6)), the set B is open. Moreover, B is also closed. In fact, let $\{y_n\} \subseteq B$, $y \in \mathbb{H}^n$ such that $d(y_n, y) \rightarrow 0$. Then by the continuity of w_ε there holds

$$\boxed{\text{e666}} \quad (4.18) \quad w_\varepsilon(\tau_{y_n}(x)) \leq u_\infty(x) \quad \text{for any } x \in \mathbb{H}^n.$$

However, if it were

$$w_\varepsilon(\tau_y(\bar{x})) = u_\infty(\bar{x}) \quad \text{for some } \bar{x} \in \mathbb{H}^n,$$

we would have a contradiction with the strong maximum principle, since u_∞ a solution of equation (4.12) and $w_\varepsilon \circ \tau_y$ a subsolution of the same equation (for, w_ε is a subsolution and Δ_H commutes with isometries of \mathbb{H}^n). Hence the inequality in (4.18) is strict, which proves that $y \in B$, thus B is closed.

Since the set B is open, closed and nonempty, Claim 2 follows. This completes the proof of claim (ii), thus the conclusion follows. \square

Remark 4.1. The arguments used in the proof of Theorem 3.2-(ii) show that $v \equiv 1$ is the unique nontrivial solution of equation (4.12) such that $0 \leq v \leq 1$, if (1.15) holds.

Let us recall the following result (see [4, Lemma A.1]).

$\boxed{\text{1e:gst}}$ **Lemma 4.1.** *For any $k > 0$ there exists a unique radial ground state $\phi = \phi(\rho)$ of equation (1.16) such that $\phi(0) = k$. Moreover, $\lim_{\rho \rightarrow \infty} \phi(\rho) = 0$.*

Proof of Theorem 3.3. By assumption (1.11) there exist $L > 0$ and $\sigma \in (0, 1)$ such that $f(u) \leq Lu^p$ for any $u \in (0, \sigma)$. Set

$$\boxed{\text{e3bu}} \quad (4.19) \quad w := c\phi \quad \text{in } \mathbb{H}^n,$$

where ϕ is the ground state mentioned in Lemma 4.1 and

$$0 < c < \frac{1}{\|\phi\|_\infty} \min \left\{ \sigma, \left(\frac{\lambda_1}{L} \right)^{\frac{1}{p-1}} \right\}.$$

Define also

$$\bar{u}(x, t) := e^{-\lambda_1 t} \zeta(t) w(\rho(x)) \quad \text{for any } (x, t) \in \mathbb{H}^n \times \bar{\mathbb{R}}_+,$$

where

$$\zeta(t) := \left\{ 1 - \frac{L}{\lambda_1} \|w\|_\infty^{p-1} [1 - e^{-(p-1)\lambda_1 t}] \right\}^{-\frac{1}{p-1}} \quad (t \in \bar{\mathbb{R}}_+).$$

Assuming that $u_0(x) \leq w(\rho(x))$ for any $x \in \mathbb{H}^n$, it is easily seen that \bar{u} is a super-solution to problem (1.1). In fact,

(a) for any $(x, t) \in \mathbb{H}^n \times \bar{\mathbb{R}}_+$ there holds

$$\bar{u}(x, t) \leq w(\rho(x)) \leq c\|\phi\|_\infty < \sigma,$$

hence

$$f(\bar{u}) \leq L\bar{u}^p \quad \text{in } \mathbb{H}^n \times \bar{\mathbb{R}}_+;$$

(b) by (a) and the definition of \bar{u} there holds

$$\begin{aligned} & \frac{\partial \bar{u}}{\partial t} - \Delta_H \bar{u} - f(\bar{u}) \geq \\ & \geq \frac{\partial \bar{u}}{\partial t} - \Delta_H \bar{u} - L\bar{u}^p \geq e^{-\lambda_1 t} w [\zeta' - L\|w\|_\infty^{p-1} e^{-(p-1)\lambda_1 t} \zeta^p] = 0; \end{aligned}$$

(c) by assumption, $u_0(x) \leq w(\rho(x)) = \bar{u}(x, 0)$.

Then by comparison results we have $0 \leq u \leq \bar{u}$ in $\mathbb{H}^n \times \bar{\mathbb{R}}_+$. Since

$$\sup_{\mathbb{H}^n} \bar{u} \leq \|w\|_\infty e^{-\lambda_1 t} \rightarrow 0 \quad \text{as } t \rightarrow \infty,$$

the conclusion follows. \square

Let us now prove Theorem 3.4. To prove claim (ii) of this theorem a preliminary result is needed.

Let $c_0 > n - 1$. By Proposition 3.1-(ii), for any $c \in (0, c_0 - (n - 1))$ there exists $\gamma_c \in (0, 1)$ such that: for any $\eta \in (\gamma_c, 1)$ there exist $b = b(c, \eta) > 0$ and a solution q of the equation

$$q'' + (c + n - 1)q' + f(q) = 0$$

such that

$$q(0) = \eta, \quad q'(0) = 0, \quad q(b) = 0, \quad q' < 0 \quad \text{in } (0, b].$$

For any fixed $R > 0$ define

$$\boxed{\text{e43a}} \quad (4.20) \quad v_0(\rho) := \begin{cases} \eta & \text{if } \rho \leq R, \\ q(\rho - R) & \text{if } R < \rho \leq R + b, \\ 0 & \text{if } R + b < \rho. \end{cases}$$

Concerning the solution of the parabolic problem

$$\boxed{\text{e43}} \quad (4.21) \quad \begin{cases} \frac{\partial v}{\partial t} = \Delta_H v + f(v) & \text{in } \mathbb{H}^n \times \mathbb{R}_+ \\ v = v_0(\rho) & \text{in } \mathbb{H}^n \times \{0\}, \end{cases}$$

the following holds (see [3, Lemma 5.1]).

$\boxed{12}$ **Lemma 4.2.** *Let assumptions (H_0) and either (H_1) or (H_2) be satisfied. Let $c_0 > n - 1$, $c \in (0, c_0 - (n - 1))$, $\gamma_c \in (0, 1)$ and $\eta \in (\gamma_c, 1)$ as above. Then for any $R > 0$ such that*

$$\boxed{\text{e42}} \quad (4.22) \quad \coth R < 1 + \frac{c}{n - 1}$$

the solution v of problem (4.21) satisfies the inequality

$$\boxed{\text{e45}} \quad (4.23) \quad v(x, t) \geq \eta \quad \text{for any } (x, t) \in \mathbb{H}^n \times \mathbb{R}_+ \text{ such that } \rho(x) \leq R + kt,$$

where

$$\boxed{\text{e454}} \quad (4.24) \quad k \equiv k(c) := c + (n-1)(1 - \coth R) > 0.$$

Therefore,

$$\boxed{\text{e44}} \quad (4.25) \quad \lim_{t \rightarrow \infty} v(x, t) = 1 \quad \text{uniformly in compact subsets of } \mathbb{H}^n.$$

Proof. Following [3], define

$$W(x, t) := v_0(\rho(x) - kt) \quad ((x, t) \in \mathbb{H}^n \times \mathbb{R}_+)$$

with k given by (4.24). It is easily seen that:

$$(i) \quad \frac{\partial W}{\partial t} - \Delta_H W - f(W) = -f(\eta)$$

if $\rho(x) \leq R + kt$;

$$(ii) \quad \frac{\partial W}{\partial t} - \Delta_H W - f(W) = q'(\rho(x) - kt - R) [c - k + (n-1)(1 - \coth \rho(x))]$$

if $R + kt < \rho(x) \leq R + b + kt$;

$$(iii) \quad \frac{\partial W}{\partial t} - \Delta_H W - f(W) = 0$$

if $\rho(x) > R + b + kt$. Observe that

$$q'(\rho(x) - kt - R) < 0,$$

$$c - k + (n-1)(1 - \coth \rho(x)) \geq c - k + (n-1)(1 - \coth R) = 0$$

if $R + kt < \rho(x) \leq R + b + kt$. Moreover, under the present assumptions there holds $f(\eta) > 0$ if η is sufficiently close to 1.

Arguing as in [3], it follows $v \geq W$ in $\mathbb{H}^n \times \mathbb{R}_+$. Since by definition $W(x, t) = \eta$ if $\rho(x) \leq R + kt$, inequality (4.23) follows. Moreover, since η is arbitrarily close to 1, equality (4.25) follows from (4.23). This completes the proof. \square

Now we can prove Theorem 3.4.

Proof of Theorem 3.4. (i) Let $\zeta = \zeta(t)$ be the solution of the problem

$$\begin{cases} \zeta' = f(\zeta) & \text{in } \mathbb{R}_+ \\ \zeta(0) = \sup_{\mathbb{H}^n} u_0. \end{cases}$$

Clearly, ζ is a supersolution of problem (1.1), thus by comparison results $0 \leq u(x, t) \leq \zeta(t)$ for any $(x, t) \in \mathbb{H}^n \times \mathbb{R}_+$. Since by assumption $\zeta(0) < a$ and (H_2) -(i) holds, we have $\zeta(t) \rightarrow 0$ as $t \rightarrow \infty$. Hence claim (i) follows.

(ii) Let $u_0(\rho) \geq v_0(\rho)$ in \mathbb{H}^n , where v_0 is the function defined in (4.20). By comparison results we have $u \geq v$ in $\mathbb{H}^n \times \mathbb{R}_+$, v being the solution of problem (4.21). Then by Lemma 4.2 the conclusion follows. \square

We conclude this section by proving Theorem 3.5.

Proof of Theorem 3.5. Let w be the solution of the problem

$$\boxed{\text{e74}} \quad (4.26) \quad \begin{cases} \frac{\partial w}{\partial t} = \Delta_H w + \sigma(\eta)w & \text{in } \mathbb{H}^n \times \mathbb{R}_+ \\ w = (u_0 - \eta)_+ & \text{in } \mathbb{H}^n \times \{0\}. \end{cases}$$

Define

$$\boxed{\text{e74a}} \quad (4.27) \quad v(x, t) := w(x, t) + \eta \quad ((x, t) \in \mathbb{H}^n \times \mathbb{R}_+).$$

Since $w \geq 0$, there holds $v \geq \eta$, whence by the definition of $\sigma(\eta)$

$$f(v) \leq \sigma(\eta)(v - \eta) = \sigma(\eta)w \quad \text{in } \mathbb{H}^n \times \mathbb{R}_+.$$

Then we have

$$\frac{\partial v}{\partial t} - \Delta_H v - f(v) \geq \frac{\partial w}{\partial t} - \Delta_H w - \sigma(\eta)w = 0 \quad \text{in } \mathbb{H}^n \times \mathbb{R}_+.$$

Moreover, there holds

$$u_0(x) \leq v(x, 0) \quad \text{for any } x \in \mathbb{H}^n.$$

Then by comparison results

$$\boxed{\text{e75}} \quad (4.28) \quad u(x, t) \leq v(x, t) \quad \text{for any } (x, t) \in \mathbb{H}^n \times \mathbb{R}_+.$$

On the other hand, from (4.26), (4.27) and the estimate (2.15) we obtain

$$\boxed{\text{e76}} \quad (4.29) \quad \begin{aligned} v(x, t) &= \eta + e^{\sigma(\eta)t} \int_{\mathbb{H}^n} G(x, y, t) [u_0(y) - \eta]_+ d\mu_y \leq \\ &\leq \eta + C_n \frac{(1+t)^{\frac{n-3}{2}}}{t^{\frac{n}{2}}} e^{[\sigma(\eta) - \lambda_1]t} \int_{\mathbb{H}^n} [u_0(y) - \eta]_+ d\mu_y \end{aligned}$$

for any $(x, t) \in \mathbb{H}^n \times \mathbb{R}_+$. Inequality (4.29) and either assumption (i), (ii) imply that there exists $t_0 > 0$ such that $v(x, t_0) \leq a$, thus by inequality (4.28) we obtain $u(x, t_0) \leq a$ for any $x \in \mathbb{H}^n$. Since by assumption $f(a) \leq 0$, by comparison we obtain that $u \leq a$ in $\mathbb{H}^n \times [t_0, \infty)$, thus $f(u) \leq 0$ in $\mathbb{H}^n \times [t_0, \infty)$ by assumption (H₄).

Now let z solve the problem

$$\begin{cases} \frac{\partial z}{\partial t} = \Delta_H z & \text{in } \mathbb{H}^n \times (t_0, \infty) \\ z = v & \text{in } \mathbb{H}^n \times \{t_0\}. \end{cases}$$

Since $f(u) \leq 0$ in $\mathbb{H}^n \times [t_0, \infty)$ and inequality (4.28) holds, by comparison results we have

$$\boxed{\text{e78}} \quad (4.30) \quad u(x, t) \leq z(x, t), \quad \text{in } \mathbb{H}^n \times [t_0, \infty).$$

On the other hand,

$$\boxed{\text{e79}} \quad (4.31) \quad \begin{aligned} z(x, t) &= \int_{\mathbb{H}^n} G(x, y, t - t_0) v(y, t_0) d\mu_y = \eta + \int_{\mathbb{H}^n} G(x, y, t - t_0) w(y, t_0) d\mu_y = \\ &= \eta + e^{\sigma(\eta)t_0} \int_{\mathbb{H}^n} G(x, y, t) [u_0(y) - \eta]_+ d\mu_y \leq \\ &\leq \eta + C_n e^{\sigma(\eta)t_0} \frac{(1+t)^{\frac{n-3}{2}}}{t^{\frac{n}{2}}} e^{-\lambda_1 t} \int_{\mathbb{H}^n} [u_0(y) - \eta]_+ d\mu_y \end{aligned}$$

for any $t \geq t_0$ (here use of (2.15) and (3.7) has been made). From inequalities (4.30)-(4.31) and either assumption (i), (ii) we obtain plainly (3.8).

To conclude the proof, let $f(u) < 0$ in $(0, \eta]$. Then by the continuity of f there exists $\varepsilon \in (0, a - \eta)$ such that $f(u) < 0$ for any $u \in (0, \eta + \varepsilon]$. On the other hand, by (4.30)-(4.31) there exists $t_\varepsilon > 0$ such that

$$\boxed{\text{e82}} \quad (4.32) \quad u(x, t) \leq z(x, t) < \eta + \frac{\varepsilon}{2} \quad \text{for any } (x, t) \in \mathbb{H}^n \times [t_\varepsilon, \infty).$$

Let ζ solve the problem

$$\begin{cases} \zeta' = f(\zeta) & \text{in } (t_\varepsilon, \infty) \\ \zeta(t_\varepsilon) = \eta + \frac{\varepsilon}{2}. \end{cases}$$

Then by inequality (4.32) ζ is a supersolution, whereas u is a subsolution of the problem

$$\begin{cases} \frac{\partial y}{\partial t} = \Delta_H y + f(y) & \text{in } \mathbb{H}^n \times (t_\varepsilon, \infty) \\ y = z & \text{in } \mathbb{H}^n \times \{t_\varepsilon\}. \end{cases}$$

Hence by comparison results

$$0 \leq u(x, t) \leq \zeta(t) \quad \text{for any } (x, t) \in \mathbb{H}^n \times [t_\varepsilon, \infty).$$

Since $\zeta(t) \rightarrow 0$ as $t \rightarrow \infty$, letting $t \rightarrow \infty$ in the above inequality we obtain equality (3.4). This completes the proof. \square

5. SPEED OF PROPAGATION AND ASYMPTOTICAL SYMMETRY: PROOFS

Let us prove Theorem 3.6.

Proof of Theorem 3.6. (i) Choose \tilde{u}_0 as in the proof of Theorem 3.2, and let \tilde{u} be the corresponding solution of (4.2). Then inequality (4.3) holds.

Since by assumption $c + n - 1 > c_0$, there exists a solution $q = q(\xi)$ of the ordinary differential equation

$$q'' + (c + n - 1)q' + f(q) = 0$$

in \mathbb{R}_+ , such that

$$q(0) = 1, \quad q' < 0 \quad \text{in } \mathbb{R}_+, \quad q(\xi) \rightarrow 0 \quad \text{as } \xi \rightarrow \infty$$

(see Proposition 3.1-(iii)). Let $\varphi = \varphi(\xi)$ be the function defined in (4.7), so that $\tilde{u}_0 \leq \varphi$ in \mathbb{R}_+ , and let v be the solution of the problem

$$\begin{cases} \frac{\partial v}{\partial t} = \frac{\partial^2 v}{\partial \xi^2} + (c + n - 1) \frac{\partial v}{\partial \xi} + f(v) & \text{in } \mathbb{R} \times \mathbb{R}_+ \\ v = \varphi & \text{in } \mathbb{R} \times \{0\}. \end{cases}$$

As in the proof of Theorem 3.2, the function v has the properties (4.9), namely

$$\frac{\partial v}{\partial \xi} \leq 0 \quad \text{in } \mathbb{R} \times \mathbb{R}_+, \quad \lim_{t \rightarrow \infty} v(\xi, t) = 0.$$

Then the function

$$w(\rho, t) := v(\rho - ct, t) \quad (\rho \geq 0, t \geq 0).$$

satisfies the problem

$$\boxed{999} \quad (5.1) \quad \begin{cases} \frac{\partial w}{\partial t} = \frac{\partial^2 w}{\partial \rho^2} + (n-1) \frac{\partial w}{\partial \rho} + f(w) & \text{in } \mathbb{R}_+ \times \mathbb{R}_+ \\ w = \varphi & \text{in } \mathbb{R}_+ \times \{0\}, \end{cases}$$

and there holds

$$\boxed{9999} \quad (5.2) \quad \frac{\partial w}{\partial \rho} \leq 0 \quad \text{in } \mathbb{R}_+ \times \mathbb{R}_+.$$

Since $\coth \rho \geq 1$, by (5.1)-(5.2) w is a supersolution of problem (4.2).

Then by comparison results and inequality (4.3) we have

$$u(x, t) \leq \tilde{u}(\rho(x), t) \leq w(\rho(x), t) = v(\rho(x) - ct, t) \quad \text{in } \mathbb{H}^n \times \mathbb{R}_+.$$

Let $y \in \mathbb{H}^n$. Since $\xi \mapsto v(\xi, t)$ is nonincreasing for any $t > 0$, and $d(x, y) > ct$ implies $\rho(x) - ct > -\rho(y)$, by (4.9) we obtain

$$\begin{aligned} \sup_{\{x \in \mathbb{H}^n \mid d(x, y) > ct\}} u(x, t) &\leq \sup_{\{x \in \mathbb{H}^n \mid d(x, y) > ct\}} v(\rho(x) - ct, t) \leq \\ &\leq v(-\rho(y), t) \rightarrow 0 \quad \text{as } t \rightarrow \infty. \end{aligned}$$

Hence the claim follows.

(ii) Under the present assumptions equality (3.5) holds, namely

$$\lim_{t \rightarrow \infty} u(x, t) = 1 \quad \text{uniformly on compact subsets of } \mathbb{H}^n.$$

Therefore, for any compact $K \subseteq \mathbb{H}^n$ and for any $\eta \in (0, 1)$ there exists $h = h(K, \eta) > 0$ such that

$$\boxed{e353} \quad (5.3) \quad u(x, t) \geq \eta \quad \text{for any } x \in K, t \geq h.$$

Let $c \in (0, c_0 - (n-1))$. Fix $\bar{c} \in (c, c_0 - (n-1))$ and $\bar{R} > 0$ such that

$$\boxed{\text{cotg}} \quad (5.4) \quad \coth \bar{R} < 1 + \frac{\bar{c} - c}{n-1},$$

thus in particular

$$\coth \bar{R} < 1 + \frac{\bar{c}}{n-1}.$$

Let $\gamma_{\bar{c}} \in (0, 1)$, $\bar{\eta} \in (\gamma_{\bar{c}}, 1)$, $\bar{b} = b(\bar{c}, \bar{\eta}) > 0$, \bar{q} , \bar{v} , \bar{v}_0 denote the quantities used in the proof of Lemma 4.2, with c replaced by \bar{c} and R by \bar{R} . Then by Lemma 4.2 we have

$$\boxed{e4545} \quad (5.5) \quad \bar{v}(x, t) \geq \bar{\eta} \quad \text{if } \rho(x) \leq \bar{R} + \bar{k}t,$$

where \bar{v} denotes the solution of problem (4.21) with Cauchy data \bar{v}_0 and

$$\bar{k} \equiv k(\bar{c}) := \bar{c} + (n-1)(1 - \coth \bar{R}) > 0$$

(see (4.23)-(4.24)).

On the other hand, observe that the definition of \bar{v}_0 (see (4.20)) and the inequality $\bar{q}' < 0$ in $(0, \bar{b}]$ imply

$$\bar{v}_0(\rho(x)) \leq \bar{\eta} \quad \text{if } \rho(x) \leq \bar{R} + \bar{b}, \quad \bar{v}_0(\rho(x)) = 0 \quad \text{otherwise.}$$

Therefore, using inequality (5.3) in the compact $K = \{x \in \mathbb{H}^n \mid \rho(x) \leq \bar{R} + \bar{b}\}$ we obtain

$$u(x, h) \geq \bar{v}_0(\rho(x)) \quad \text{for any } x \in \mathbb{H}^n,$$

whence by comparison results

$$\boxed{e555} \quad (5.6) \quad u(x, t+h) \geq \bar{v}(x, t) \quad \text{for any } (x, t) \in \mathbb{H}^n \times \mathbb{R}_+.$$

From (5.5)-(5.6) we obtain

$$u(x, t+h) \geq \bar{\eta} \quad \text{for any } (x, t) \in \mathbb{H}^n \times \mathbb{R}_+ \text{ such that } \rho(x) \leq \bar{R} + \bar{k}t \quad (t \in \mathbb{R}_+),$$

namely

$$u(x, t) \geq \bar{\eta} \quad \text{for any } (x, t) \in \mathbb{H}^n \times \mathbb{R}_+ \text{ such that } \rho(x) \leq \bar{R} + \bar{k}(t-h) \quad (t > h).$$

Let $y \in \mathbb{H}^n$. Since inequality (5.4) implies $c < \bar{k}$, we have

$$\{x \in \mathbb{H}^n \mid d(x, y) < ct\} \subseteq \{x \in \mathbb{H}^n \mid \rho(x) < \bar{R} + \bar{k}(t-h)\}$$

for any $t \in \mathbb{R}_+$ sufficiently large. Therefore, by the above remarks

$$\inf_{d(x,y) < ct} u(x, t) \geq \inf_{\rho(x) \leq \bar{R} + \bar{k}(t-h)} u(x, t) \geq \bar{\eta}.$$

To summarize, we proved that for any $\bar{\eta} \in (\gamma_{\bar{c}}, 1)$ there exists $\tau \in \mathbb{R}_+$ such that

$$\inf_{d(x,y) < ct} u(x, t) \geq \bar{\eta} \quad \text{for any } t > \tau.$$

Since $u \leq 1$ in $\mathbb{H}^n \times \mathbb{R}_+$, equality (3.5) follows. This completes the proof. \square

Proof of Theorem 3.7. Denote by ω the convex hull of the support of u_0 . Take any hyperplane $\pi \subseteq \mathbb{H}^n$ such that $\pi \cap \omega = \emptyset$. Let $\mathbb{H}_\omega^n \subseteq \mathbb{H}^n$ be the half-space containing ω . Define

$$\tilde{u}(x) := u(R_\pi(x)) \quad (x \in \mathbb{H}^n),$$

where R_π denotes the reflection with respect to π . Observe that both u and \tilde{u} satisfy the equation

$$\frac{\partial u}{\partial t} = \Delta_H u + f(u) \quad \text{in } \mathbb{H}_\omega^n \times \mathbb{R}_+,$$

whereas

$$u = u_0 \geq 0 = \tilde{u} \quad \text{in } \mathbb{H}_\omega^n \times \{0\}, \quad u = \tilde{u} \quad \text{in } \pi \times \mathbb{R}_+.$$

Then by the strong maximum principle we get

$$u > \tilde{u} \quad \text{in } \mathbb{H}_\omega^n \times \mathbb{R}_+,$$

and by the Hopf Boundary Lemma we obtain

$$\frac{\partial u}{\partial \nu} > \frac{\partial \tilde{u}}{\partial \nu} = -\frac{\partial u}{\partial \nu} \quad \text{in } \pi \times \mathbb{R}_+,$$

where ν is the vector field orthogonal to π pointing towards \mathbb{H}_ω^n . Hence

$$\boxed{\text{e53}} \quad (5.7) \quad \langle \nabla_H u, \nu \rangle_H > 0 \quad \text{in } \pi \times \mathbb{R}_+.$$

(i) Let us now prove the first claim. Since $\text{supp } u_0$ is compact and (3.5) holds, we have $\Gamma_a(u; t) \cap \omega = \emptyset$ for any $a \in (0, 1)$ and $t \in \mathbb{R}_+$ sufficiently large. Let $x_0 \in \Gamma_a(u; t)$. Then we can find a hyperplane $P \subseteq \mathbb{H}^n$ such that $x_0 \in P$ and $P \cap \omega = \emptyset$ (see Remark 2.1-(ii)). Using the above remarks with $\pi = P$, by (5.7) we obtain $\nabla_H u(x_0, t) \neq 0$. Since x_0 is arbitrary, the smoothness of $\Gamma_a(u; t)$ follows.

(ii) To prove the second claim we argue by contradiction. Let there exist $x_0 \in \Gamma_a(u; t)$ and an infinite geodesic γ orthogonal to $\Gamma_a(u; t)$ at x_0 , which does not intersect ω . Then we can choose a hyperplane $Q \subseteq \mathbb{H}^n$ such that $\gamma \subseteq Q$, $Q \cap \omega = \emptyset$ (see Remark 2.1-(iii)). Using the above remarks with $\pi = Q$, by (5.7) we obtain $\langle \nabla_H u(x_0, t), \tau \rangle_H \neq 0$, where τ is a tangent vector to $\Gamma_a(u; t)$ in x_0 . However, this contradicts the very definition of $\Gamma_a(u; t)$. Hence the conclusion follows. \square

6. HOROSPHERIC WAVES: PROOFS

hopf

Let us first prove Theorem 3.9.

Proof of Theorem 3.9. Since the function u defined in (1.18) only depends on (x_n, t) , plainly we have

$$\begin{aligned} \frac{\partial u}{\partial t} - \Delta_H u - f(u) &= -c^* (q^*)' - [(q^*)'' + (q^*)'] + (2-n)(q^*)' - f(q^*) = \\ &= -(q^*)'' + (1-n-c^*)(q^*)' - f(q^*) = 0 \end{aligned}$$

(here the expression (2.17) of Δ_H in \mathbb{U}^n has been used). Therefore u satisfies equation (1.12). Moreover, its level set is

$$\Gamma_a(u; t) := \left\{ x \in \mathbb{U}^n \mid x_n = e^{-z-c^*t} \right\} \quad (a \in (0, 1); t \in \mathbb{R}_+),$$

where $z := (q^*)^{-1}(a)$ (recall that q^* is decreasing in \mathbb{R} by Proposition 3.1-(i)). Then for any $a \in (0, 1)$ and $t \in \mathbb{R}_+$ $\Gamma_a(u; t)$ is a horosphere with centre at infinity. This completes the proof. \square

To prove Theorem 3.10 we need some preliminary results. As anticipated in Subsection 3.3, the formal analogy between horospheric waves in \mathbb{H}^n and planar waves in \mathbb{R}^n suggests to introduce the new coordinate $e^{c^*t}x_n$. However, at variance from the case of translations in \mathbb{R}^n , the change of coordinates

$$\chi_t : \mathbb{U}^n \rightarrow \mathbb{U}^n, \quad \chi_t(x', x_n) := (x', e^{c^*t}x_n) \quad (x' \in \mathbb{R}^{n-1}, x_n > 0, t \in \mathbb{R}_+)$$

is not an isometry of \mathbb{U}^n (hereafter, any point $x \in \mathbb{U}^n$ is denoted by $x \equiv (x', x_n)$ with $x' \equiv (x_1, \dots, x_{n-1}) \in \mathbb{R}^{n-1}, x_n \in \mathbb{R}_+$). Instead, the map

$$\tau_t : \mathbb{U}^n \rightarrow \mathbb{U}^n, \quad \tau_t(x', x_n) := e^{c^*t}(x', x_n) \equiv (X', X_n) \quad (t \in \mathbb{R}_+)$$

is an isometry (see (2.16)). Therefore we define

$$w := u \circ \tau_t^{-1} : \mathbb{U}^n \times \mathbb{R}_+ \rightarrow \mathbb{R}, \quad w(X', X_n, t) := u(e^{-c^*t}X', e^{-c^*t}X_n, t) \quad (t \in \mathbb{R}_+),$$

where $u = u(x', x_n, t)$ is any solution of problem (3.12). An elementary calculation shows that w solves the problem

$$\text{e778} \quad (6.1) \quad \begin{cases} \frac{\partial w}{\partial t} = X_n^2 \Delta w - (c_0 - 1) \frac{\partial w}{\partial X_n} X_n - c^* \sum_{i=1}^{n-1} \frac{\partial w}{\partial X_i} X_i + f(w) & \text{in } \mathbb{U}^n \times \mathbb{R}_+ \\ w(X', X_n, 0) = w_0(X', X_n) := u_0(X', X_n) \end{cases}$$

((X', X_n) $\in \mathbb{U}^n$; here the expression (2.17) of Δ_H in \mathbb{U}^n has been used).

Defining $z := -\log X_n$ and

$$v : \mathbb{R}^n \times \mathbb{R}_+ \rightarrow \mathbb{R}, \quad v(X', z, t) := w(X', e^{-z}, t),$$

problem (6.1) reads

$$\text{e777} \quad (6.2) \quad \begin{cases} \frac{\partial v}{\partial t} = \frac{\partial^2 v}{\partial z^2} + c_0 \frac{\partial v}{\partial z} + e^{-2z} \sum_{i=1}^{n-1} \frac{\partial^2 v}{\partial X_i^2} - c^* \sum_{i=1}^{n-1} \frac{\partial v}{\partial X_i} X_i + f(v) & \text{in } \mathbb{R}^n \times \mathbb{R}_+ \\ v(X', z, 0) = v_0(X', z) := w_0(X', e^{-z}) \end{cases}$$

((X', z) $\in \mathbb{R}^n$; recall that by definition $c_0 = c^* + n - 1$).

If v does not depend on $X' \in \mathbb{R}^{n-1}$, by (6.2) it satisfies the equation

$$\boxed{\text{e81r}} \quad (6.3) \quad \frac{\partial v}{\partial t} = \frac{\partial^2 v}{\partial z^2} + c_0 \frac{\partial v}{\partial z} + f(v) \quad \text{in } \mathbb{R} \times \mathbb{R}_+.$$

Concerning (6.3), let us recall the following lemma ([9]; see also [16, Lemma 4.2]).

$\boxed{13}$ **Lemma 6.1.** *Let assumption (H_5) be satisfied. Then for any $\delta_1 \in (0, a)$ and any $\delta_2 \in (0, 1 - a)$ there exist $\beta > 0$ and $C \geq 1$ (only depending on δ_1, δ_2 and f) such that the functions $v^\pm : \mathbb{R} \times \mathbb{R}_+ \rightarrow \mathbb{R}$,*

$$\boxed{\text{defpiu}} \quad (6.4) \quad \begin{aligned} v^+(z, t) &:= q^*(z - C\delta_1(1 - e^{-\beta t})) + \delta_1 e^{-\beta t}, \\ v^-(z, t) &:= q^*(z + C\delta_2(1 - e^{-\beta t})) - \delta_2 e^{-\beta t} \end{aligned}$$

are a supersolution and a subsolution of equation (6.3), respectively.

By a suitable modification of the proof of [16, Lemma 4.1], we can prove the following result.

$\boxed{\text{t10}}$ **Proposition 6.2.** *Let assumption (H_5) be satisfied. Let $v = v(X', z, t)$ be any solution of problem (6.2).*

(i) *Suppose that*

$$\boxed{\text{cin}} \quad (6.5) \quad \limsup_{z \rightarrow \infty} \sup_{X' \in \mathbb{R}^{n-1}} v_0(X', z) < a.$$

Then there exists $z^* \in \mathbb{R}$ such that

$$\boxed{\text{e80r}} \quad (6.6) \quad \limsup_{t \rightarrow \infty} \sup_{X' \in \mathbb{R}^{n-1}} v(X', z, t) \leq q^*(z - z^*) \quad \text{uniformly for } z \in \mathbb{R}.$$

(ii) *Similarly, if*

$$\boxed{\text{cin1}} \quad (6.7) \quad \liminf_{z \rightarrow -\infty} \inf_{X' \in \mathbb{R}^{n-1}} v_0(X', z) > a,$$

there exists $z_* \in \mathbb{R}$ such that

$$\boxed{\text{e79z}} \quad (6.8) \quad \liminf_{t \rightarrow \infty} \inf_{X' \in \mathbb{R}^{n-1}} v(X', z, t) \geq q^*(z - z_*) \quad \text{uniformly for } z \in \mathbb{R}.$$

Proof. We only prove claim (i), the proof of (ii) being similar. To this purpose, we shall prove the following

Claim: There exist $\tau \in \mathbb{R}_+$, $z_0 \in \mathbb{R}$ and $\delta_1 \in (0, a)$ such that

$$\boxed{\text{e83r}} \quad (6.9) \quad \sup_{X' \in \mathbb{R}^{n-1}} v(X', z, \tau) \leq v^+(z - z_0, 0) = q^*(z - z_0 - C\delta_1) + \delta_1 \quad \text{for any } z \in \mathbb{R}.$$

By the Claim the conclusion follows. In fact, inequality (6.9) and Lemma 6.1 yield by comparison results

$$\sup_{X' \in \mathbb{R}^{n-1}} v(X', z, t) \leq v^+(z - z_0, t - \tau) \quad \text{for any } z \in \mathbb{R}, t \geq \tau.$$

Letting $t \rightarrow \infty$ in the above inequality gives

$$\limsup_{t \rightarrow \infty} \sup_{X' \in \mathbb{R}^{n-1}} v(X', z, t) \leq q^*(z - z_0 - C\delta_1)$$

(see (6.4)), namely inequality (6.6) with $z^* := z_0 + C\delta_1$.

It remains to prove the Claim. This will be done in three steps.

(i) By inequality (6.5) we can choose $\delta_1 \in (0, a)$ and $\beta \in (0, \delta_1)$ such that

$$\boxed{\text{e83rs}} \quad (6.10) \quad \limsup_{z \rightarrow \infty} \sup_{X' \in \mathbb{R}^{n-1}} v_0(X', z) < \beta < \delta_1.$$

Let us show that for every fixed $\tau \in \mathbb{R}_+$

$$\boxed{\text{e85r}} \quad (6.11) \quad \limsup_{z \rightarrow \infty} \sup_{X' \in \mathbb{R}^{n-1}} v(X', z, \tau) < \delta_1.$$

To this purpose, choose first $M \geq 1$ such that

$$\boxed{\text{e85rs}} \quad (6.12) \quad \sup_{X' \in \mathbb{R}^{n-1}} v_0(X', z) \leq \beta + Me^{-c_0 z} \quad \text{for any } z \in \mathbb{R}.$$

This is possible, since by inequality (6.10) there exists $\bar{z} > 0$ such that

$$\sup_{X' \in \mathbb{R}^{n-1}} v_0(X', z) \leq \beta \quad \text{for any } z > \bar{z};$$

moreover,

$$\sup_{X' \in \mathbb{R}^{n-1}} v_0(X', z) \leq 1 \leq e^{-c_0(z-\bar{z})} \quad \text{for any } z \leq \bar{z}.$$

Setting $M := e^{c_0 \bar{z}}$ we obtain (6.12).

Further, let us observe that the function

$$\bar{v}(z, t) := \beta + Me^{-c_0(z-\alpha t)} \quad (z \in \mathbb{R}, t \in \mathbb{R}_+)$$

is a supersolution of the first equation in (6.2) for any $\alpha > 0$ sufficiently large. In fact, it is easily checked that

$$\boxed{\text{e85rst}} \quad (6.13) \quad \frac{\partial \bar{v}}{\partial t} - \frac{\partial^2 \bar{v}}{\partial z^2} - c_0 \frac{\partial \bar{v}}{\partial z} - f(\bar{v}) = \alpha Me^{-c_0(z-\alpha t)} - f(\beta + Me^{-c_0(z-\alpha t)}) \geq 0$$

if $\alpha \geq \frac{1}{a-\beta} \sup_{s \in [a, 1]} f(s)$, since

$$f(\bar{v}) \geq 0 \quad \Leftrightarrow \quad 0 \leq z - \alpha t \leq \frac{1}{c_0} \left| \log \frac{a-\beta}{M} \right|$$

(see assumption (H_5)).

By (6.12)-(6.13) the function \bar{v} (with α, β, M as above) is a supersolution of problem (6.2). Therefore by comparison we obtain

$$\sup_{X' \in \mathbb{R}^{n-1}} v(X', z, \tau) \leq \beta + Me^{-c_0(z-\alpha\tau)}$$

for any $z \in \mathbb{R}$ and any fixed $\tau \in \mathbb{R}_+$. Hence inequality (6.11) follows.

(ii) Since by assumption (H_5) there holds $f(s) < 0$ if $s > 1$, the function $\tilde{v} := 1 + \frac{\delta_1}{2}$ is a supersolution of problem (6.2). Hence by comparison

$$\boxed{\text{e84r}} \quad (6.14) \quad \sup_{X' \in \mathbb{R}^{n-1}} v(X', z, \tau) \leq 1 + \frac{\delta_1}{2} \quad \text{for any } z \in \mathbb{R}, \tau \in \mathbb{R}_+.$$

(iii) Fix any $\tau \in \mathbb{R}_+$. Set $z_1 := C\delta_1 + (q^*)^{-1} \left(1 - \frac{\delta_1}{2} \right)$. Then by (6.14) we have

$$\sup_{X' \in \mathbb{R}^{n-1}} v(X', z, \tau) \leq 1 + \frac{\delta_1}{2} \leq q^*(z - C\delta_1) + \delta_1 \quad \text{for any } z \geq z_1.$$

On the other hand, by (6.11) there exists $z_2 > 0$ such that

$$\sup_{X' \in \mathbb{R}^{n-1}} v(X', z, \tau) \leq \delta_1 \quad \text{for any } z > z_2.$$

Defining $z_0 := \max\{z_2 - z_1, 0\}$ we obtain inequality (6.9), thus the Claim follows. This completes the proof. \square

Now we can prove Theorem 3.10.

Proof of Theorem 3.10. Under the change of variables $(x', x_n) \rightarrow (X', z)$, assumption (3.13) corresponds to assumptions (6.5)-(6.7) of Proposition 6.2. Then by inequalities (6.6)-(6.8) we obtain

$$q^*(z - z_*) \leq v(X', z, t) \leq q^*(z - z^*) \text{ for every } (X', z) \in \mathbb{R}^n, t \in \mathbb{R}_+.$$

Since by definition $z = -\log x_n - c^*t$ and $v(X', z, t) = u(x', x_n, t)$, setting $k^* := e^{z^*}$, $k_* := e^{z_*}$ inequality (3.14) follows. This completes the proof. \square

REFERENCES

- A** [1] L. A. Ahlfors, *Möbius Transformations in Several Variables*, Lecture Notes, School of Mathematics, University of Minnesota (1981).
- AW2** [2] D. G. Aronson & H. F. Weinberger, *Nonlinear diffusion in population genetics, combustion, and nerve pulse propagation*, in "Partial Differential Equations and Related Topics", Lecture Notes in Mathematics **446**, pp. 6-49 (Springer, 1975).
- AW** [3] D. G. Aronson & H. F. Weinberger, *Multidimensional nonlinear diffusion arising in population genetics*, Adv. Math. **30** (1978), 33-76.
- BPT** [4] C. Bandle, M. A. Pozio & A. Tesei, *The Fujita exponent for the Cauchy problem in the hyperbolic space*, J. Differential Equations **251** (2011), 2143-2163.
- BP** [5] R. Benedetti & C. Petronio, *Lectures on Hyperbolic Geometry* (Springer, 1992).
- BH** [6] H. Berestycki & F. Hamel, *Generalized travelling waves for reaction diffusion equations*, in "Perspectives in Nonlinear Partial Differential Equations. In Honor of H. Brezis", Contemporary Mathematics **446**, pp. 101-123 (Amer. Math. Soc., 2007).
- Dav** [7] E. B. Davies, *Heat Kernel and Spectral Theory* (Cambridge University Press, 1989).
- Fen** [8] W. Fenchel, *Elementary Geometry in Hyperbolic Space* (de Gruyter, 1989).
- FML** [9] P.C. Fife & J. B. McLeod, *The approach of solutions of nonlinear diffusion equations to travelling front solutions*, Arch. Rat. Mech. Anal. **65** (1977), 335-361.
- Fu66** [10] H. Fujita, *On the blowing up of solutions of the Cauchy problem for $u_t = \Delta u + u^{1+\alpha}$* , J. Fac. Sci. Tokyo Sect. IA Math. **13** (1966), 109-124.
- Grig** [11] A. Grigoryan, *Analytic and geometric background of recurrence and non-explosion of the Brownian motion on Riemannian manifolds*, Bull. Amer. Math. Soc. **36** (1999), 135-249.
- Iv** [12] B. Iversen, *Hyperbolic Geometry* (London Mathematical Society, 1992).
- Jon** [13] C. Jones, *Asymptotic behaviour of a reaction-diffusion equation in higher space dimensions*, Rocky Mount. J. Math. **13** (1983), 355-364.
- KP** [14] S. Kumaresan & J. Prajapat, *Serrin's result for hyperbolic space and sphere*, Duke Math. J. **91** (1998), 17-28.
- KP2** [15] S. Kumaresan & J. Prajapat, *Analogue of Gidas-Ni Nirenberg result in hyperbolic and spherical spaces*, Rend. Ist. Mat. Univ. Trieste **30** (1998), 107-112.
- MN** [16] H. Matano & M. Nara, *Large time behavior of disturbed planar fronts in the Allen-Cahn equation*, J. Differential Equations **251** (2011), 3522-3557.
- PPT** [17] M. A. Pozio, F. Punzo & A. Tesei, *Uniqueness and nonuniqueness of solutions to parabolic problems with singular coefficients*, Discr. Cont. Dyn. Syst. **30** (2011), 891-916.

DEPARTMENT OF MATHEMATICS, GRADUATE SCHOOL OF MATHEMATICAL SCIENCES, UNIVERSITY OF TOKYO, 3-8-1 KOMABA, MEGURO-KU, TOKYO 153-8902, JAPAN
E-mail address: matano@ms.u-tokyo.ac.jp

DIPARTIMENTO DI MATEMATICA "G. CASTELNUOVO", UNIVERSITÀ DI ROMA "LA SAPIENZA", P.LE A. MORO 5, I-00185 ROMA, ITALY
E-mail address: punzo@mat.uniroma1.it

DIPARTIMENTO DI MATEMATICA "G. CASTELNUOVO", UNIVERSITÀ DI ROMA "LA SAPIENZA", P.LE A. MORO 5, I-00185 ROMA, ITALY
E-mail address: tesei@mat.uniroma1.it