

AMENABILITY AND SUBEXPONENTIAL SPECTRAL GROWTH RATE OF DIRICHLET FORMS ON VON NEUMANN ALGEBRAS

FABIO CIPRIANI, JEAN-LUC SAUVAGEOT

ABSTRACT. In this work we apply Noncommutative Potential Theory to characterize (relative) amenability and the (relative) Haagerup Property (H) of von Neumann algebras in terms of the spectral growth of Dirichlet forms. Examples deal with (inclusions of) countable discrete groups and free orthogonal compact quantum groups.

1. INTRODUCTION AND DESCRIPTION OF THE RESULTS.

Classical results relate the metric properties of conditionally negative definite functions on a countable discrete group Γ to its approximation properties. For example, there exists a *proper*, conditionally negative definite function ℓ on Γ if and only if there exists a sequence $\varphi_n \in c_0(\Gamma)$ of normalized, positive definite functions, vanishing at infinity and converging pointwise to the constant function 1.

In a celebrated work [Haa2], U. Haagerup proved that the length function of a free group \mathbb{F}_n with $n \in \{2, \dots, \infty\}$ generators is negative definite, thus establishing for free groups the above approximation property. Since then the property is referred to as Haagerup Approximation Property (H) or Gromov a-T-menability (see [CCJJV]).

In addition, if for a conditionally negative definite function ℓ on a countable discrete group Γ , the series $\sum_{g \in \Gamma} e^{-t\ell(g)}$ converges for all $t > 0$, then there exists a sequence $\varphi_n \in l^2(\Gamma)$ of normalized, positive definite functions, converging pointwise to the constant function 1 ([GK Thm 5.3]). This latter property is just one of the several equivalent appearances of *amenability*, a property introduced by J. von Neumann in 1929 [vN] in order to explain the Banach-Tarski paradox in Euclidean spaces \mathbb{R}^n exactly when $n \geq 3$.

In this note we are going to discuss extensions of the above results concerning amenability for σ -finite von Neumann algebras N .

The direction along which we are going to look for substitutes of the above summability condition related to amenability, is that of Noncommutative Potential Theory.

This is suggested by a recent result by Caspers-Skalski [CaSk] asserting that N has the (suitably formulated) Haagerup Approximation Property (H) if and only if there exists a Dirichlet form $(\mathcal{E}, \mathcal{F})$ on the standard Hilbert space $L^2(N)$, having *discrete spectrum*.

The link between the properness condition for a conditionally negative definite function ℓ on a countable discrete groups Γ and the generalized one on von Neumann algebras, relies on the fact that, when the von Neumann algebra $N = L(\Gamma)$ is the one generated by the left regular representation of Γ , the quadratic form $\mathcal{E}_\ell[a] = \sum_{g \in \Gamma} \ell(g)|a(g)|^2$ on the standard space $L^2(L(\Gamma), \tau) \simeq l^2(\Gamma)$ is a Dirichlet form if and only if the function ℓ is conditionally negative

¹This work has been supported by GREFI-GENCO INDAM Italy-CNRS France and MIUR PRIN 2012 Project No 2012TC7588-003

Date: December 3rd, 2016, revised May 10th 2017, final revision October 8th 2017.

1991 Mathematics Subject Classification. 46L57, 46L87, 46L54, 43A07.

Key words and phrases. von Neumann algebra, Amenability, Haagerup Property (H), Dirichlet form spectral growth, countable discrete group.

definite and its spectrum is discrete if and only if ℓ is proper. Moreover, on a countable, finitely generated, discrete group Γ with polynomial growth, there exist a conditionally negative definite functions ℓ , having polynomial growth and growth dimensions arbitrarily close to the homogeneous dimension of Γ (see [CS5]).

This point of view thus suggests that a condition providing amenability of a von Neumann algebra with faithful normal state (N, ω) could be the *subexponential spectral growth* of a Dirichlet form $(\mathcal{E}, \mathcal{F})$ on the standard space $L^2(N, \omega)$, i.e. the discreteness of the spectrum of $(\mathcal{E}, \mathcal{F})$ and the summability of the series $\sum_{k \geq 0} e^{-t\lambda_k}$ for all $t > 0$, where $\lambda_0, \lambda_1, \dots$ are the eigenvalues of $(\mathcal{E}, \mathcal{F})$.

The second fundamental fact that will allow to use Dirichlet forms to investigate the amenability of a von Neumann algebra, is the possibility to express this property in terms of Connes' *correspondences*: N is amenable if and only if the identity or standard N - N -correspondence $L^2(N)$ is weakly contained in the coarse or Hilbert-Schmidt N - N -correspondence $L^2(N) \otimes L^2(N)$ (see [Po1]).

In the second part of the work we provide a condition guaranteeing the *relative amenability* of an inclusion $B \subseteq N$ of finite von Neumann algebras introduced by Popa [Po1,2], in terms of the existence of a Dirichlet form $(\mathcal{E}, \mathcal{F})$ on $L^2(N)$ having *relative subexponential spectral growth*. Also this result is based on the possibility to express the relative amenability of a von Neumann algebra N with respect to a subalgebra $B \subseteq N$ in terms of the weak containment of the identity correspondence $L^2(N)$ in the relative tensor product correspondence $L^2(N) \otimes_B L^2(N)$ introduced by Sauvageot [S1], [Po2].

Using a suitable Dirichlet form constructed in [CFK], whose construction uses tools developed by M. Brannan in [Bra], we apply the above result to prove amenability of the von Neumann algebra of the free orthogonal quantum group O_2^+ and Haagerup Property (H) of the free orthogonal quantum groups O_N^+ for $N \geq 3$ (see also the recent [DFSW]), results firstly obtained by M. Brannan [Bra].

A detailed discussion of the relative Haagerup Property (H) for inclusions of countable discrete groups in terms of conditionally negative definite functions is presented.

The paper is organized as follows: in Section 2 we provide the necessary tools on noncommutative potential theory on von Neumann algebra as Dirichlet forms, Markovian semigroups and resolvents.

In Section 3 we first recall some equivalent constructions of the coarse or Hilbert-Schmidt correspondence of a von Neumann algebra N and some connections between the modular theories of N , of its opposite N^o , and of their spatial tensor product $N \overline{\otimes} N^o$. Then we introduce the spectral growth rate of a Dirichlet form and we prove the first main result of the work about the amenability of von Neumann algebra admitting a Dirichlet form with subexponential spectral growth rate. This part terminates with an application to the amenability of countable discrete groups and with a partially alternative approach to the proof of a result of M. Brannan [Bra] about the amenability of the free orthogonal quantum group O_2^+ .

Section 4 starts recalling some fundamental tool of the basic construction $\langle N, B \rangle$ for inclusions $B \subseteq N$ of finite von Neumann algebras, needed to prove the second main result of the work concerning the amenability of N with respect to its subalgebra B . To formulate the criterion, we introduce the spectral growth rate of a B -invariant Dirichlet form on the standard space $L^2(N)$ relatively to the subalgebra B , using the compact ideal space $\mathcal{J}(\langle N, B \rangle)$ of $\langle N, B \rangle$ (cf. [PO1,2]). The section terminates discussing relative amenability for two natural subalgebras $B_{\min} \subseteq N$ and $B_{\max} \subseteq N$ associated to any Dirichlet form.

In Section 5 we extend the spectral characterization of the Haagerup Property (H) of von Neumann algebras with countable decomposable center due to M. Caspers and A. Skalski [CaSk] to the Relative Haagerup Property (H) for inclusions of finite von Neumann algebras $B \subseteq N$ formulated by S. Popa [Po 1,2].

In Section 6 we discuss the relative Haagerup Property (H) for inclusions $H < G$ of countable discrete groups in terms of the existence of an H -invariant conditionally negative definite function on G which is proper on the homogeneous space G/H and in terms of quasi-normality of H in G .

The content of the present work has been the subject of talks given in Rome II (March 2015), Paris (GREFI-GENCO April 2015), Berkeley (UC Seminars September 2015), Krakov (September 2015), Varese (May 2016).

2. DIRICHLET FORMS ON σ -FINITE VON NEUMANN ALGEBRAS

Recall that a von Neumann algebra N is σ -finite, or countably decomposable, if any collection of mutually orthogonal projections is at most countable and that this property is equivalent to the existence of a normal, faithful state. This is the case, for example, if N acts faithfully on a separable Hilbert space.

Let us consider on a σ -finite von Neumann algebra N a fixed faithful, normal state $\omega \in N_{*+}$. Let us denote by $(N, L^2(N, \omega), L^2_+(N, \omega), J_\omega)$ the standard form of N and by $\xi_\omega \in L^2_+(N, \omega)$ the cyclic vector representing the state (see [Haa1]).

For a real vector $\xi = J_\omega \xi \in L^2(N, \omega)$, let us denote by $\xi \wedge \xi_\omega$ the Hilbert projection of the vector ξ onto the closed and convex set $C_\omega := \{\eta \in L^2(N, \omega) : \eta = J_\omega \eta, \xi_\omega - \eta \in L^2_+(N, \omega)\}$.

We recall here the definition of Dirichlet form and Markovian semigroup (see [C1]) on a generic standard form of a σ -finite von Neumann algebra. For a definition particularized to the Haagerup standard form see [GL1].

Definition 2.1 (Dirichlet forms on σ -finite von Neumann algebras). A densely defined, non-negative and lower semicontinuous quadratic form $\mathcal{E} : L^2(N, \omega) \rightarrow [0, +\infty]$ is said to be:

i) *real* if

$$(2.1) \quad \mathcal{E}[J_\omega(\xi)] = \mathcal{E}[\xi] \quad \xi \in L^2(N, \omega);$$

ii) a *Dirichlet form* if it is real and *Markovian* in the sense that

$$(2.2) \quad \mathcal{E}[\xi \wedge \xi_\omega] \leq \mathcal{E}[\xi] \quad \xi = J_\omega \xi \in L^2(N, \omega);$$

iii) a *completely Dirichlet form* if all the canonical extensions \mathcal{E}_n to $L^2(\mathbb{M}_n(N), \omega \otimes \text{tr}_n)$

$$(2.3) \quad \mathcal{E}_n[[\xi_{i,j}]_{i,j=1}^n] := \sum_{i,j=1}^n \mathcal{E}[\xi_{i,j}] \quad [\xi_{i,j}]_{i,j=1}^n \in L^2(\mathbb{M}_n(N), \omega \otimes \text{tr}_n),$$

are Dirichlet forms.

By the self-polarity of the standard cone $L^2_+(N, \omega)$, any real vector $\xi = J_\omega \xi \in L^2(N, \omega)$ decomposes uniquely as a difference $\xi = \xi_+ - \xi_-$ of two positive, orthogonal vectors $\xi_\pm \in L^2_+(N, \omega)$ (the positive part ξ_+ being just the Hilbert projection of ξ onto the positive cone). The modulus of ξ is then defined as the sum of the positive and negative parts $|\xi| := \xi_+ + \xi_-$. Notice that, in general, the contraction property

$$\mathcal{E}[|\xi|] \leq \mathcal{E}[\xi] \quad \xi = J_\omega \xi \in L^2(N, \omega)$$

is a consequence of Markovianity and that it is actually equivalent to it when $\mathcal{E}[\xi_\omega] = 0$.

The domain of the Dirichlet form is defined as the (dense) subspace of $L^2(N, \omega)$ where the quadratic form is finite: $\mathcal{F} := \{\xi \in L^2(N, \omega) : \mathcal{E}[\xi] < +\infty\}$. We will denote by $(L, D(L))$ the densely defined, self-adjoint, nonnegative operator on $L^2(N, \omega)$ associated with the closed quadratic form $(\mathcal{E}, \mathcal{F})$

$$\mathcal{F} = D(\sqrt{L}) \quad \text{and} \quad \mathcal{E}[\xi] = \|\sqrt{L}\xi\|^2 \quad \xi \in D(\sqrt{L}) = \mathcal{F}.$$

Definition 2.2 (Markovian semigroups on standard forms of von Neumann algebras).

a) A bounded operator T on $L^2(N, \omega)$ is said to be

i) *real* if it commutes with the modular conjugation: $TJ_\omega = J_\omega T$,

ii) *positive* if it leaves globally invariant the positive cone: $T(L^2_+(N, \omega)) \subseteq L^2_+(N, \omega)$,

iii) *Markovian* if it is real and it leaves globally invariant the closed, convex set C_ω :

$$T(C_\omega) \subseteq C_\omega,$$

iv) *completely positive, resp. completely Markovian*, if it is real and all of its matrix amplifications $T^{(n)}$ to $L^2(\mathbb{M}_n(N), \omega \otimes \text{tr}_n) \simeq L^2(N, \omega) \otimes L^2(\mathbb{M}_n(\mathbb{C}), \text{tr}_n)$ defined by

$$T^{(n)}[[\xi_{i,j}]_{i,j=1}^n] := \sum_{i,j=1}^n [T\xi_{i,j}]_{i,j=1}^n \quad [\xi_{i,j}]_{i,j=1}^n \in L^2(\mathbb{M}_n(N), \omega \otimes \text{tr}_n),$$

are positive, resp. Markovian;

b) A strongly continuous, uniformly bounded, self-adjoint semigroup $\{T_t : t > 0\}$ on $L^2(N, \omega)$ is said to be real (resp. positive, Markovian, completely positive, completely Markovian) if the operators T_t are real (resp. positive, Markovian, completely positive, completely Markovian) for all $t > 0$.

In literature, property in item iii) above is sometime termed *submarkovian*, while *markovian* is meant positivity preserving *and unital*. Our choice is only dictated by a willing of simplicity.

Notice that T is (completely) Markovian iff it is (completely) positive and $T\xi_\omega \leq \xi_\omega$.

Notice that if N is abelian, then positive (resp. Markovian) operators are automatically completely positive (resp. completely Markovian).

Dirichlet forms are in one-to-one correspondence with Markovian semigroups (see [C1]) through the relations

$$T_t = e^{-tL} \quad t \geq 0$$

where $(L, D(L))$ is the self-adjoint operator associated to the quadratic form $(\mathcal{E}, \mathcal{F})$.

Dirichlet forms and Markovian semigroups are also in correspondence with a class of semigroups on the von Neumann algebra. To state this fundamental relation, let us consider the *symmetric embedding* i_ω determined by the cyclic vector ξ_ω

$$i_\omega : N \rightarrow L^2(N, \omega) \quad i_\omega(x) := \Delta_\omega^{\frac{1}{4}} x \xi_\omega \quad x \in N.$$

Here, Δ_ω is the modular operator associated with the faithful normal state ω (see [T]). We will denote by $\{\sigma_t^\omega : t \in \mathbb{R}\}$ the modular automorphisms group associated to ω and by $N_{\sigma^\omega} \subseteq N$ the subalgebra of elements which are analytic with respect to it. Then (see [C1]) (completely) Dirichlet forms $(\mathcal{E}, \mathcal{F})$ and (completely) Markovian semigroups $\{T_t : t > 0\}$ on $L^2(N, \omega)$ are in one-to-one correspondence with those weakly*-continuous, (completely) positive and contractive semigroups $\{S_t : t > 0\}$ on the von Neumann algebra N which are *modular ω -symmetric* in the sense that

$$(2.4) \quad \omega(S_t(x)\sigma_{-i/2}^\omega(y)) = \omega(\sigma_{-i/2}^\omega(x)S_t(y)) \quad x, y \in N_{\sigma^\omega}, \quad t > 0,$$

through the relation

$$i_\omega(S_t(x)) = T_t(i_\omega(x)) \quad x \in N, \quad t > 0.$$

Relation (2.4) is called *modular symmetry* and it is equivalent to

$$(2.5) \quad (J_\omega y \xi_\omega | S_t(x) \xi_\omega) = (J_\omega S_t(y) \xi_\omega | x \xi_\omega) \quad x, y \in N, \quad t > 0.$$

Remark 2.3. In case ω is a trace, the symmetric embedding reduces to $i_\omega(x) = x \xi_\omega$ while the modular symmetry simplifies to $\omega(S_t(x)y) = \omega(xS_t(y))$ for $x, y \in N$ and $t > 0$.

*To shorten notations, in the forthcoming part of the paper
"Dirichlet form" will always mean "completely Dirichlet form" and
"Markovian semigroup" will always mean "completely Markovian semigroup".*

Whenever no confusion can arise, the modular conjugation J_ω will be sometime denoted J .

2.1. Examples of Dirichlet forms. Instances of the notions introduced above may be found in various frameworks. We just recall here some examples of different origins. One may consult the fundamental works [BeDe], [FOT] for the commutative case and [C2], [C3] for surveys in the noncommutative setting.

a) The archetypical Dirichlet form on the Euclidean space \mathbb{R}^n or, more generally, on any Riemannian manifold V , endowed with its Riemannian measure m , is the Dirichlet integral

$$\mathcal{E}[a] = \int_V |\nabla a|^2 dm \quad a \in L^2(V, m).$$

In this case the trace on $L^\infty(V, m)$ is given by the integral with respect to the measure m and the form domain is the Sobolev space $H^1(V) \subset L^2(V, m)$. The associated Markovian semigroup is the familiar heat semigroup of the Riemannian manifold. Interesting variations of the above Dirichlet integral are the Dirichlet forms of type

$$\mathcal{E}[a] := \int_{\mathbb{R}^n} |\nabla a|^2 d\mu \quad a \in L^2(\mathbb{R}^n, \mu),$$

that for suitable choices of positive Radon measures μ , are ground state representations of Hamiltonian operators in Quantum Mechanics.

b) Dirichlet forms are a fundamental tool to introduce differential calculus and study Markovian stochastic processes on fractal sets (see [Ki], [CS3], [CGIS 1,2]).

c) On a countable discrete group Γ , any conditionally negative definite function ℓ gives rise to a Dirichlet form

$$\mathcal{E}_\ell[\xi] := \sum_{s \in \Gamma} |\xi(s)|^2 \ell(s),$$

on the Hilbert space $l^2(\Gamma)$, considered as the standard Hilbert space of the left von Neumann algebra $L(\Gamma)$ generated by the left regular representation of Γ (see [CS1], [C2]). The associated Markovian semigroup is simply given by the multiplication operator

$$T_t(a)(s) = e^{-t\ell(s)} a(s) \quad t > 0, \quad s \in G, \quad a \in l^2(\Gamma).$$

d) On noncommutative tori A_θ , $\theta \in [0, 1]$ (see [Co2]), which are C^* -algebras generated by two unitaries u and v , satisfying the relation

$$vu = e^{2i\pi\theta} uv,$$

the *heat semigroup* $\{T_t : t \geq 0\}$ defined by

$$T_t(u^n v^m) = e^{-t(n^2 + m^2)} u^n v^m \quad (n, m) \in \mathbb{Z}^2,$$

is a τ -symmetric Markovian semigroup on the von Neumann algebra N_θ generated by the G.N.S. representation of the faithful, tracial state $\tau : A_\theta \rightarrow \mathbb{C}$ characterized by

$$\tau(u^n v^m) = \delta_{n,0} \delta_{m,0} \quad n, m \in \mathbb{Z}.$$

e) There exists a general interplay between Dirichlet forms and differential calculus on tracial C^* -algebras (A, τ) (see [S 2,3], [CS1]) and this provides a source of Dirichlet forms on von Neumann algebras (generated by A in the G.N.S. representation of the trace). In fact, denoting by N the von Neumann algebra generated by the G.N.S. representation of the trace, if $(\partial, D(\partial))$ is a densely defined closable derivation from $L^2(N, \tau)$ to Hilbert A -bimodule \mathcal{H} , then the closure of the quadratic form

$$\mathcal{E}[a] := \|\partial a\|_{\mathcal{H}}^2 \quad a \in \mathcal{F} := D(\partial)$$

is a Dirichlet form on $L^2(N, \tau)$. Viceversa, any Dirichlet form on $L^2(N, \tau)$ whose domain is dense in A arises in this way from an essentially unique derivation on A canonically associated with it (see [CS1]). Examples of this differential calculus can be found in all the situations illustrated above as well as in the geometric framework of Riemannian foliations (see [S4]) and also in the framework of Voiculescu's Free Probability theory (see [V1]). There, the Dirichlet form associated to Voiculescu's derivation presents several aspects connected to Noncommutative Hilbert Transform, Free Fischer Information and Free Entropy.

3. AMENABILITY OF σ -FINITE VON NEUMANN ALGEBRAS

In this section we relate a certain characteristic of the spectrum of a Dirichlet form to the amenability of the von Neumann algebra. Recall that a von Neumann algebra N is said to be *amenable* if, for every normal dual Banach N -bimodule X , the derivations $\delta : N \rightarrow X$ are all inner, i.e. they have the form

$$\delta(x) = x\xi - \xi x \quad x \in N$$

form some vector $\xi \in X$. It is a remarkable fact, and the byproduct of a tour de force, that this property is equivalent to several others of apparently completely different nature, such as *hyperfiniteness*, *injectivity*, *semi-discreteness*, *Schwartz property P*, *Tomiyama property E*. We refer to [Co2 Ch. V] for a review on these connections. Among the main examples of amenable von Neumann algebras, we recall: the von Neumann algebra of a locally compact amenable group, the crossed product of an abelian von Neumann algebra by an amenable locally compact group, the commutant von Neumann algebra of any continuous unitary representation of a connected locally compact group, the von Neumann algebra generated by any representation of a nuclear C^* -algebra.

3.1. Standard form of the spatial tensor product of von Neumann algebras. Here we summarize some well known properties of the standard form of the spatial tensor product of two von Neumann algebras in terms of Hilbert-Schmidt operators (details may be found in [T]), mainly with the intention to make precise, in the next section, some properties of the symmetric embedding of a product state. More precisely we shall use the following facts:

3.1. *let $N \subseteq \mathcal{B}(\mathcal{H})$ be a von Neumann algebra. A vector $\xi \in H$ is cyclic for the commutant N' if and only if it is separating for N ;*

3.2. *let $N_k \subseteq \mathcal{B}(H_k)$ $k = 1, 2$ be von Neumann algebras. If the vectors $\xi_k \in H_k$, $k = 1, 2$ are cyclic for N_k , then the vector $\xi_1 \otimes \xi_2 \in H_1 \otimes H_2$ is cyclic for the spatial tensor product $N_1 \overline{\otimes} N_2$;*

3.3. *Let N_k $k = 1, 2$ be von Neumann algebras and $L^2(N_k)$ their standard forms. If the vectors $\xi_k \in L^2_+(N_k)$ $k = 1, 2$ are cyclic for N_k (hence separating) then the vector $\xi_1 \otimes \xi_2 \in L^2(N_1) \otimes L^2(N_2)$ is cyclic and separating for the spatial tensor product $N_1 \overline{\otimes} N_2$.*

3.2. Symmetric embedding of tensor product of von Neumann algebras. Here we recall the definition and a property of the symmetric embedding of a von Neumann algebra in its standard Hilbert space. Let N be a σ -finite von Neumann algebra and $\omega \in N_{*,+}$ a faithful, normal state.

In the standard form $(N, L^2(N, \omega), L^2_+(N, \omega))$, we denote by $\xi_\omega \in L^2_+(N, \omega)$ the cyclic vector representing the state ω and by J_ω and Δ_ω its modular conjugation and modular operator, respectively.

The symmetric embedding $i_\omega : N \rightarrow L^2(N, \omega)$, defined by $i_\omega(x) := \Delta_\omega^{\frac{1}{4}} x \xi_\omega$ for $x \in N$, is a completely positive contraction with dense range, which is also continuous between the weak*-topology of N and the weak topology of $L^2(N, \omega)$. It is also an order isomorphism of completely ordered sets between $\{x = x^* \in N : 0 \leq x \leq 1_N\}$ and $\{\xi = J_\omega \xi \in L^2(N, \omega) : 0 \leq \xi \leq \xi_\omega\}$ (see [Ara], [Co1], [Haa1] and [BR]). We shall make use of the following properties:

3.4. *Let N_k $k = 1, 2$ be von Neumann algebras and $L^2(N_k)$ their standard forms. Consider the cyclic (hence separating) vectors $\xi_k \in L^2(N_k)$ $k = 1, 2$ and the cyclic and separating vector $\xi_1 \otimes \xi_2 \in H_1 \otimes H_2$ for the spatial tensor product $N_1 \overline{\otimes} N_2$.*

Let J_k, Δ_k be the modular conjugation and the modular operator associated to $\xi_k \in H_k$ $k = 1, 2$ and $J_{\xi_1 \otimes \xi_2}, \Delta_{\xi_1 \otimes \xi_2}$ be the modular conjugation and the modular operator associated to $\xi_1 \otimes \xi_2$. Then the following identifications hold true

- $J_{\xi_1 \otimes \xi_2} = J_{\xi_1} \otimes J_{\xi_2}$;
- $N_1 \xi_1 \odot N_2 \xi_2 \subseteq H_1 \otimes H_2$ is a core for the closed operator $\Delta_{\xi_1 \otimes \xi_2}^{\frac{1}{2}}$;
- $\Delta_{\xi_1 \otimes \xi_2}^{\frac{1}{2}}(\eta_1 \otimes \eta_2) = \Delta_{\xi_1}^{\frac{1}{2}}(\eta_1) \otimes \Delta_{\xi_2}^{\frac{1}{2}}(\eta_2)$ for $\eta_k \in N_k \xi_k$ and $k = 1, 2$.

We will denote by N° the opposite algebra of N : it coincides with N as a vector space but the product is taken in the reverse order $x^\circ y^\circ := (yx)^\circ$ for $x^\circ, y^\circ \in N^\circ$. As customary, we adopt the convention that elements $y \in N$, when regarded as elements of the opposite algebra are denoted by $y^\circ \in N^\circ$.

A linear functional ω on N , when considered as a linear functional on the opposite algebra N° is denoted by ω° and called the *opposite* of ω . As N and N° share the same positive cone, if ω is positive on N so is ω° on N° and if ω is normal so does its opposite.

By the properties of standard forms of von Neumann algebras, it follows that for the standard form $(N^\circ, L^2(N^\circ, \omega^\circ), L^2_+(N^\circ, \omega^\circ))$ of N° one has the following identifications

$$L^2(N^\circ, \omega^\circ) = L^2(N, \omega), \quad L^2_+(N^\circ, \omega^\circ) = L^2_+(N, \omega), \quad J_\omega = J_{\omega^\circ}, \quad \Delta_{\omega^\circ} = \Delta_\omega^{-1}, \quad \xi_{\omega^\circ} = \xi_\omega.$$

Using the isomorphism between N° and the commutant N' , given by $N^\circ \ni y^\circ \rightarrow J_\omega y^* J_\omega \in N'$, we can regard $L^2(N, \omega)$ not only as a left N -module but also as a left N° -module, hence as a right N -module and finally as a N - N -bimodule

$$y^\circ \xi := J_\omega y^* J_\omega \xi, \quad \xi y := J_\omega y^* J_\omega \xi, \quad x \xi y := x J_\omega y^* J_\omega \xi \quad x, y \in N, \xi \in L^2(N, \omega).$$

The symmetric embeddings associated to ω and ω° are related by

$$\begin{aligned} i_{\omega^\circ}(y^\circ) &= \Delta_{\omega^\circ}^{\frac{1}{4}}(\xi_\omega y) = \Delta_{\omega^\circ}^{\frac{1}{4}} J_\omega y^* J_\omega \xi_\omega = \Delta_\omega^{-\frac{1}{4}} \Delta_\omega^{\frac{1}{2}} y \xi_\omega = \Delta_\omega^{\frac{1}{4}} y \xi_\omega = i_\omega(y). \\ J_\omega(i_\omega(y^*)) &= J_\omega \Delta_\omega^{\frac{1}{4}}(y^* \xi_\omega) = J_\omega \Delta_\omega^{\frac{1}{4}} J_\omega \Delta_\omega^{\frac{1}{2}}(y \xi_\omega) = \Delta_\omega^{\frac{1}{4}}(y \xi_\omega) = i_\omega(y) = i_{\omega^\circ}(y^\circ). \end{aligned}$$

3.3. Coarse correspondence. Recall that a Hilbert-Schmidt operator T is a bounded operator on $L^2(N, \omega)$ such that $\text{Trace}_{L^2(N, \omega)}(T^*T) < +\infty$. It may be represented as

$$T\xi := \sum_{k=0}^{\infty} \mu_k(\eta_k|\xi)\xi_k \quad \xi \in L^2(N, \omega)$$

in terms of suitable orthonormal systems $\{\eta_k : k \in \mathbb{N}\}$, $\{\xi_k : k \in \mathbb{N}\} \subset L^2(N, \omega)$ and a sequence $\{\mu_k : k \in \mathbb{N}\} \subset \mathbb{C}$ such that $\sum_{k=0}^{\infty} |\mu_k|^2 < +\infty$. The set of Hilbert-Schmidt operators $HS(L^2(N, \omega))$ is a Hilbert space under the scalar product $(T_1|T_2) := \text{Trace}_{L^2(N, \omega)}(T_1^*T_2)$.

Lemma 3.5. *The binormal representations $\pi_{\text{co}}^1, \pi_{\text{co}}^2, \pi_{\text{co}}^3$ of $N \otimes_{\max} N^\circ$, characterized by*

$$\begin{aligned} \pi_{\text{co}}^1 : N \otimes_{\max} N^\circ &\rightarrow \mathcal{B}(\text{HS}(L^2(N, \tau))) \\ \pi_{\text{co}}^1(x \otimes y^\circ)(T) &:= xTy \quad x, y \in N, \quad T \in \text{HS}(L^2(N, \tau)), \\ \pi_{\text{co}}^2 : N \otimes_{\max} N^\circ &\rightarrow \mathcal{B}(L^2(N, \tau) \otimes \overline{L^2(N, \tau)}) \\ \pi_{\text{co}}^2(x \otimes y^\circ)(\xi \otimes \bar{\eta}) &:= x\xi \otimes \bar{\eta}y \quad x, y \in N, \quad \xi, \eta \in L^2(N, \tau) \\ \pi_{\text{co}}^3 : N \otimes_{\max} N^\circ &\rightarrow \mathcal{B}(L^2(N, \tau) \otimes L^2(N, \tau)) \\ \pi_{\text{co}}^3(x \otimes y^\circ)(\xi \otimes \eta) &:= x\xi \otimes \eta y \quad x, y \in N, \quad \xi, \eta \in L^2(N, \tau), \end{aligned}$$

are unitarely equivalent by

$$\begin{aligned} U : L^2(N, \tau) \otimes \overline{L^2(N, \tau)} &\rightarrow L^2(N, \tau) \otimes L^2(N, \tau) \quad U(\xi \otimes \bar{\eta}) := \xi \otimes J_\omega \eta \\ V : L^2(N, \tau) \otimes \overline{L^2(N, \tau)} &\rightarrow \text{HS}(L^2(N, \tau)) \quad V(\xi \otimes \bar{\eta})(\zeta) := (\eta|\zeta)\xi. \end{aligned}$$

They give rise by weak closure

$$(\pi_{\text{co}}^3(N \otimes_{\max} N^\circ))'' = N \overline{\otimes} N^\circ$$

of the spatial tensor product of N by its opposite N° .

Lemma 3.6. *The normal extension of the coarse representation π_{co} of the C^* -algebra $N \otimes_{\max} N^\circ$ to the von Neumann tensor product $N \overline{\otimes} N^\circ$ is the standard representation of $N \overline{\otimes} N^\circ$ (and it will still denoted by the same symbol).*

The standard positive cone in the various equivalent representations is determined as

- $\text{HS}(L^2(N, \omega))_+$, the set of all nonnegative Hilbert-Schmidt operators on $L^2(N, \omega)$;
- $(L^2(N, \tau) \otimes \overline{L^2(N, \omega)})_+$, generated by the vectors $\xi \otimes \bar{\xi}$ with $\xi \in L^2(N, \omega)$;
- $(L^2(N, \tau) \otimes L^2(N, \omega))_+$, generated by the vectors $\xi \otimes J_\omega \xi$ with $\xi \in L^2(N, \omega)$.

The standard Hilbert space and the positive cone of $N \overline{\otimes} N^\circ$ will be denoted also by

$$L^2(N \overline{\otimes} N^\circ, \omega \otimes \omega^\circ), \quad L^2_+(N \overline{\otimes} N^\circ, \omega \otimes \omega^\circ).$$

Lemma 3.7. *Let $T : L^2(N, \omega) \rightarrow L^2(N, \omega)$ be a bounded operator and consider on the involutive algebra $N \odot N^\circ$, the linear functional determined by*

$$\Theta_T : N \odot N^\circ \rightarrow \mathbb{C} \quad \Theta_T(x \otimes y^\circ) := (i_\omega(y^*)|Ti_\omega(x)) \quad x \otimes y^\circ \in N \odot N^\circ.$$

Then Θ_T is a positive linear functional on $N \odot N^\circ$ if and only if T is completely positive (c.f. Definition 2.2 iv)).

Proof. i) The positive cone of $N \odot N^\circ$ is generated by elements of type $\nu^* \nu = \sum_{j,k=1}^n x_j^* x_k \otimes (y_k y_j^*)^\circ$ where $\nu = \sum_{k=1}^n x_k \otimes y_k^\circ \in N \odot N^\circ$. The result then follows by the identity

$$\Theta_T(\nu^* \nu) = \sum_{j,k=1}^n \Theta_T(x_j^* x_k \otimes (y_k y_j^*)^\circ) = \sum_{j,k=1}^n (i_\omega(y_j y_k^*) | T i_\omega(x_j^* x_k)),$$

the completely positivity of the symmetric embedding $i_\omega : N \rightarrow L^2(N, \tau)$ and the positivity of $[x_j^* x_k]_{j,k=1}^n$ and $[y_j y_k^*]_{j,k=1}^n$ in $\mathbb{M}_n(N)$. \square

Lemma 3.8. *Let $T : L^2(N, \omega) \rightarrow L^2(N, \omega)$ be a completely positive operator and consider the positive linear functional Θ_T on $N \odot N^\circ$. Then, among the properties*

- a) Θ_T is a state on $N \odot N^\circ$
- b) T is a contraction
- c) $T\xi_\omega = \xi_\omega$

we have that the following relations

- i) a) and b) imply c) and $\|T\| = 1$
- ii) c) implies a) and b).

Proof. i) By a) and b) we have $1 = \Theta_T(1_N \otimes 1_{N^\circ}) = (\xi_\omega | T\xi_\omega) \leq \|\xi_\omega\| \cdot \|T\xi_\omega\| \leq \|\xi_\omega\|^2 \cdot \|T\| = 1$ that implies $\|T\| = \|T\xi_\omega\| = 1$ and $(\xi_\omega | T\xi_\omega) = \|\xi_\omega\| \cdot \|T\xi_\omega\|$ which provide $T\xi_\omega = \xi_\omega$. ii) The proof that c) implies a) is immediate while the proof that c) implies b) can be found in [C1]. \square

3.4. Spectral growth rate. In the following definition, the notion of growth rate of a finitely generated, countable discrete group is extended to σ -finite von Neumann algebras having the Haagerup Property (H), i.e. von Neumann algebras admitting Dirichlet forms with discrete spectrum. The idea for this generalization results from [CS5] (see discussion in Example 3.11 below).

Definition 3.9. (Spectral growth rate of Dirichlet forms). Let (N, ω) be a σ -finite, von Neumann algebra with a fixed faithful, normal state on it. To avoid trivialities we assume N to be *infinite dimensional*.

Let $(\mathcal{E}, \mathcal{F})$ be a Dirichlet form on $L^2(N, \omega)$ and let $(L, D(L))$ be the associated nonnegative, self-adjoint operator. Assume that its spectrum $\sigma(L) = \{\lambda_k \geq 0 : k \in \mathbb{N}\}$ is *discrete*, i.e. its points are isolated eigenvalues of finite multiplicity (repeated in non decreasing order according to their multiplicities).

Then let us set

$$\Lambda_n := \{k \in \mathbb{N} : \lambda_k \in [0, n]\}, \quad \beta_n := \#\Lambda_n, \quad n \in \mathbb{N}$$

and define the *spectral growth rate* of $(\mathcal{E}, \mathcal{F})$ as

$$\Omega(\mathcal{E}, \mathcal{F}) := \limsup_{n \in \mathbb{N}} \sqrt[n]{\beta_n}.$$

The Dirichlet form $(\mathcal{E}, \mathcal{F})$ is said to have

- *exponential growth* if $(\mathcal{E}, \mathcal{F})$ has discrete spectrum and $\Omega(\mathcal{E}, \mathcal{F}) > 1$
- *subexponential growth* if $(\mathcal{E}, \mathcal{F})$ has discrete spectrum and $\Omega(\mathcal{E}, \mathcal{F}) = 1$
- *polynomial growth* if $(\mathcal{E}, \mathcal{F})$ has discrete spectrum and, for some $c, d > 0$, $\beta_n \leq c \cdot n^d$ for all $n \in \mathbb{N}$
- *intermediate growth* if it has subexponential growth but not polynomial growth.

Lemma 3.10. *Setting $\gamma_0 = \beta_0$ and*

$$\gamma_n := \beta_n - \beta_{n-1} = \#\{k \in \mathbb{N} : \lambda_k \in (n-1, n]\}, \quad n \in \mathbb{N}^*,$$

and

$$\Omega'(\mathcal{E}, \mathcal{F}) := \limsup_{n \in \mathbb{N}^*} \sqrt[n]{\gamma_n}$$

we have

$$\Omega(\mathcal{E}, \mathcal{F}) = \Omega'(\mathcal{E}, \mathcal{F}) \geq 1.$$

Proof. On one hand, by definition, we have $\Omega(\mathcal{E}, \mathcal{F}) \geq \Omega'(\mathcal{E}, \mathcal{F})$. On the other hand, since, by assumption, N is infinite dimensional and $\sigma(L)$ is discrete, we have $\Omega(\mathcal{E}, \mathcal{F}) \geq \Omega'(\mathcal{E}, \mathcal{F}) \geq 1$. Consider now the following identity involving analytic functions in a neighborhood of $0 \in \mathbb{C}$

$$\sum_{n=0}^{\infty} \beta_n z^n = (1-z)^{-1} \sum_{n=0}^{\infty} \gamma_n z^n$$

and notice that the radius of convergence of the series on the left-hand side is $R = 1/\Omega(\mathcal{E}, \mathcal{F})$, while the radius of convergence of the series on the right-hand side is $R' = 1/\Omega'(\mathcal{E}, \mathcal{F})$ so that $R \leq R' \leq 1$. Since $(1-z)^{-1}$ is analytic in the open unit disk centered in $z = 0$, the above identity implies that $R \geq R'$ so that $\Omega(\mathcal{E}, \mathcal{F}) \leq \Omega'(\mathcal{E}, \mathcal{F})$. \square

Example 3.11. (Spectral growth rate on countable discrete groups).

i) On a countable discrete group Γ , if there exists a proper, c.n.d. function ℓ , then the associated Dirichlet form $(\mathcal{E}_\ell, \mathcal{F}_\ell)$ has discrete spectrum $\sigma(L) = \{\ell(g) \in [0, +\infty) : g \in \Gamma\}$.

ii) On a finitely generated, countable discrete group Γ , if the length ℓ_S corresponding to a finite system of generators $S \subseteq \Gamma$ is negative definite, then the spectral growth rate $\Omega(\mathcal{E}_{\ell_S}, \mathcal{F}_{\ell_S})$ of the corresponding Dirichlet form coincides with growth rate of (Γ, S) (see [deH Ch. VI]).

iii) Moreover, if (Γ, S) has polynomial growth, it has been shown in [CS5] that there exists on Γ a proper, c.n.d. function ℓ with polynomial growth. The associated Dirichlet form $(\mathcal{E}_\ell, \mathcal{F}_\ell)$ will have polynomial spectral growth rate.

Remark 3.12. By a well known bound (see [R Theorem 3.37])

$$1 \leq \liminf_n \frac{\beta_{n+1}}{\beta_n} \leq \limsup_{n \in \mathbb{N}} \sqrt[n]{\beta_n},$$

if the spectral growth rate is subexponential, then $\liminf_n \frac{\beta_{n+1}}{\beta_n} = 1$ so that there exists a subsequence of $\{\frac{\beta_{n+1}}{\beta_n}\}_{n \in \mathbb{N}}$ converging to 1. In other words, the sequence of spectral subspaces $\{E_n\}_{n \in \mathbb{N}}$ corresponding to the interval $[0, n] \subset [0, +\infty)$ admits a subsequence such that

$$\lim_k \frac{\dim E_{n_k+1}}{\dim E_{n_k}} = 1.$$

Subexponential growth can be equivalently stated in terms of the nuclearity of the completely Markovian semigroup $\{e^{-tL} : t > 0\}$ on $L^2(N, \omega)$:

Lemma 3.13. *The Dirichlet form $(\mathcal{E}, \mathcal{F})$ has discrete spectrum and subexponential spectral growth if and only if the Markovian semigroup $\{e^{-tL} : t > 0\}$ on $L^2(N, \omega)$ is nuclear, or trace-class, in the sense that:*

$$\text{Trace}(e^{-tL}) = \sum_{k \in \mathbb{N}} e^{-t\lambda_k} < +\infty \quad t > 0.$$

Proof. Since

$$\gamma_0 + \sum_{n \in \mathbb{N}^*} \gamma_n e^{-tn} \leq \sum_{k \in \mathbb{N}} e^{-t\lambda_k} \leq \gamma_0 + e^t \sum_{n \in \mathbb{N}^*} \gamma_n e^{-tn} \quad t > 0,$$

the series $\sum_{k \in \mathbb{N}} e^{-t\lambda_k}$ and $\sum_{n \in \mathbb{N}^*} \gamma_n e^{-tn}$ converge or diverge simultaneously. They obviously converge for all $t > 0$ if and only if $\Omega'(\mathcal{E}, \mathcal{F}) \leq 1$. \square

Example 3.14. If on a countable discrete group Γ , there exists a c.n.d. function ℓ , such that $\sum_{g \in \Gamma} e^{-t\ell(g)} < +\infty$ for all $t > 0$, then ℓ is proper, the spectrum of the associated Dirichlet form $(\mathcal{E}_\ell, \mathcal{F}_\ell)$ coincides with $\{\ell(g) \in [0, +\infty) : g \in \Gamma\}$ and it is thus discrete with subexponential growth.

The following is the main result of this section.

Theorem 3.15. *Let (N, ω) be a σ -finite von Neumann algebra endowed with a normal, faithful state on it. If there exists a Dirichlet form $(\mathcal{E}, \mathcal{F})$ on $L^2(N, \omega)$ having subexponential spectral growth, then N is amenable.*

Proof. Recall that N is amenable if and only if the identity or standard bimodule ${}_N L^2(N)_N$ is weakly contained in the coarse or Hilbert-Schmidt bimodule \mathcal{H}_{co} (see [Po1]). Consider the completely positive semigroup $\{T_t := e^{-tL} : t > 0\}$ and assume, for simplicity, that the cyclic vector is invariant: $T_t \xi_\omega = \xi_\omega$ for all $t > 0$. Recall (cf. Lemma 3.7) that the complete positivity of T_t provides a binormal state on $N \otimes_{\max} N^\circ$ characterized by

$$\Phi_t : N \otimes_{\max} N^\circ \rightarrow \mathbb{C} \quad \Phi_t(x \otimes y^\circ) := (i_\omega(y^*) | T_t i_\omega(x)).$$

To compute this state, we consider the spectral representation $T_t = \sum_{k \geq 0} e^{-t\lambda_k} P_k$ (converging strongly) in terms of the rank-one projections P_k on $L^2(N, \omega)$ associated to each eigenvalue λ_k (repeated according to their multiplicity). Notice that by Markovianity, the semigroup commutes with the modular conjugation J_ω so that each eigenvector ξ_k may be assumed to be real: $\xi_k = J_\omega \xi_k$. We then have

$$\begin{aligned} \Phi_t(x \otimes y^\circ) &= (i_\omega(y^*) | T_t i_\omega(x)) \\ &= \sum_{k=0}^{\infty} e^{-t\lambda_k} (i_\omega(y^*) | P_k(i_\omega(x))) \\ &= \sum_{k=0}^{\infty} e^{-t\lambda_k} (i_\omega(y^*) | (\xi_k | i_\omega(x)) \xi_k) \\ &= \sum_{k=0}^{\infty} e^{-t\lambda_k} (\xi_k | i_\omega(x)) (i_\omega(y^*) | \xi_k). \end{aligned}$$

As the series $Z_t := \sum_{k=0}^{\infty} e^{-t\lambda_k} \xi_k \otimes \xi_k$ is norm convergent for all $t > 0$ by the nuclearity of the semigroup, since J_ω is an antiunitary operator on $L^2(N)$, using properties in item 3.4 above

we have

$$\begin{aligned}
\Phi_t(x \otimes y^\circ) &= \sum_{k=0}^{\infty} e^{-t\lambda_k} (\xi_k | i_\omega(x)) (J_\omega \xi_k | J_\omega i_\omega(y^*)) \\
&= \sum_{k=0}^{\infty} e^{-t\lambda_k} (\xi_k | i_\omega(x)) (\xi_k | i_{\omega^\circ}(y^\circ)) \\
&= \sum_{k=0}^{\infty} e^{-t\lambda_k} (\xi_k \otimes \xi_k | i_\omega(x) \otimes i_{\omega^\circ}(y^\circ))_{L^2(N, \omega) \otimes L^2(N, \omega)} \\
&= \left(\sum_{k=0}^{\infty} e^{-t\lambda_k} \xi_k \otimes \xi_k | i_\omega(x) \otimes i_{\omega^\circ}(y^\circ) \right)_{L^2(N, \omega) \otimes L^2(N, \omega)} \\
&= \left(Z_t | i_{\omega \otimes \omega^\circ}(x \otimes y^\circ) \right)_{L^2(N \overline{\otimes} N^\circ, \omega \otimes \omega^\circ)}.
\end{aligned}$$

Since the symmetric embeddings of von Neumann algebras are continuous when $N \overline{\otimes} N^\circ$ is endowed with the weak*-topology and $L^2(N \overline{\otimes} N^\circ, \omega \otimes \omega^\circ)$ is endowed with the weak topology, by continuity we have

$$\Phi_t(z) = \left(Z_t | i_{\omega \otimes \omega^\circ}(z) \right)_{L^2(N \overline{\otimes} N^\circ, \omega \otimes \omega^\circ)} \quad z \in N \overline{\otimes} N^\circ.$$

In other words, the linear functional Φ_t extends as a σ -weakly continuous linear functional on the spatial tensor product $N \overline{\otimes} N^\circ$. Φ_t being positive by Lemma 3.7, there exist a unique positive element $\Omega_t \in L^2_+(N \overline{\otimes} N^\circ, \omega \otimes \omega^\circ)$ (see [Haa1]) such that

$$\Phi_t(z) = (i_\omega(y^*) | T_t i_\omega(x)) = \left(\Omega_t | \pi_{\text{co}}(z) \Omega_t \right)_{L^2(N \overline{\otimes} N^\circ, \omega \otimes \omega^\circ)} \quad z \in N \overline{\otimes} N^\circ$$

and the GNS representation of $N \otimes_{\max} N^\circ$ associated to Φ_t coincides with a sub-representation of π_{co} . In other words, the $N - N$ -correspondence \mathcal{H}_t associated to the completely positive map T_t is contained in the coarse $N - N$ -correspondence \mathcal{H}_{co} for all $t > 0$. Since the semigroup $\{T_t : t > 0\}$ is strongly continuous on $L^2(N, \omega)$, for all $x \otimes y^\circ \in N \otimes_{\max} N^\circ$ we have

$$\begin{aligned}
\lim_{t \downarrow 0} \left(\Omega_t | \pi_{\text{co}}(x \otimes y^\circ) \Omega_t \right)_{L^2(N, \omega) \otimes L^2(N, \omega)} &= (i_\omega(y^*) | i_\omega(x))_{L^2(N, \omega)} \\
&= (\Delta_\omega^{\frac{1}{4}} y^* \xi_\omega | \Delta_\omega^{\frac{1}{4}} x \xi_\omega) \\
&= (\Delta_\omega^{\frac{1}{2}} y^* \xi_\omega | x \xi_\omega) \\
&= (J_\omega y \xi_\omega | x \xi_\omega) \\
&= (J_\omega y J_\omega \xi_\omega | x \xi_\omega) \\
&= (\xi_\omega | J_\omega y^* J_\omega x \xi_\omega) \\
&= (\xi_\omega | x J_\omega y^* J_\omega \xi_\omega) \\
&= (\xi_\omega | x \xi_\omega y) \\
&= (\xi_\omega | \pi_{\text{id}}(x \otimes y^\circ) \xi_\omega)
\end{aligned}$$

and by continuity

$$\lim_{t \downarrow 0} \left(\Omega_t | \pi_{\text{co}}(z) \Omega_t \right)_{L^2(N, \omega) \otimes L^2(N, \omega)} = (\xi_\omega | \pi_{\text{id}}(z) \xi_\omega) \quad z \in N \otimes_{\max} N^\circ.$$

This proves that the identity correspondence \mathcal{H}_{id} is weakly contained in the coarse correspondence \mathcal{H}_{co} and thus N is amenable at least if the semigroup leaves the cyclic vector invariant. To deal with the general case, remark first that, by strong continuity, we have that $\lim_{t \downarrow 0} (\xi_\omega | T_t \xi_\omega) = \|\xi_\omega\|^2 = 1$ and there exist $t_0 > 0$ such that $(\xi_\omega | T_t \xi_\omega) > 0$ for all $0 < t < t_0$. Applying the argument above to the binormal states

$$\tilde{\Phi}_t(x \otimes y^\circ) := \frac{1}{(\xi_\omega | T_t \xi_\omega)} (i_\omega(y^*) | T_t i_\omega(x)) \quad x \otimes y^\circ \in N \otimes_{\max} N^\circ, \quad 0 < t < t_0$$

we get the amenability of N even in the general situation. \square

Remark 3.16. i) The above result implies that if the von Neumann algebra N is not amenable, then any Dirichlet form $(\mathcal{E}, \mathcal{F})$ with respect to any normal, faithful state ω has exponential growth rate $\Omega(\mathcal{E}, \mathcal{F}) > 1$, i.e. its sequence of eigenvalues has exponentially growing distribution. ii) Conversely, it is an open question whether there exist amenable von Neumann algebras on which every Dirichlet form has exponential growth. The analogy with discrete groups suggests that the the answer is likely positive.

The following one is a generalization of a result of Guentner-Kaminker [GK].

Corollary 3.17. *Let Γ be a countable discrete group, $\lambda : \Gamma \rightarrow \mathcal{B}(l^2(\Gamma))$ be its left regular representation, $L(\Gamma)$ its associated von Neumann algebra and τ its trace state. If there exists a Dirichlet form $(\mathcal{E}, \mathcal{F})$ on $L^2(L(\Gamma), \tau)$ having subexponential spectral growth, then the group Γ is amenable.*

Proof. Under the assumptions, the group von Neumann algebra $L(\Gamma)$ is amenable by the above theorem. Hence by a well known result of A. Connes, the group Γ is amenable. \square

Example 3.18. (Free orthogonal quantum groups) On the von Neumann algebra $L^\infty(O_2^+, \tau)$ of the free orthogonal quantum group O_2^+ with respect to its Haar state τ , it has been constructed in [CFK] a Dirichlet form with an explicitly computed discrete spectrum of polynomial growth (and spectral dimension $d := \limsup_n \frac{\ln \beta_n}{\ln n} = 3$). Applying the theorem above one obtains a proof of the amenability of $L^\infty(O_2^+, \tau)$, a result which has been proved by M. Brannan [Bra].

4. RELATIVE AMENABILITY OF INCLUSIONS OF FINITE VON NEUMANN ALGEBRAS

In this section we extend the previous result to the *relative amenability* of inclusions of finite von Neumann algebras $B \subseteq N$, as defined by S. Popa [Po 1,2]. This extension is based on the properties of the relative tensor product of Hilbert bimodules and on the properties of the *basic construction*, which we will presently recall (see [Chr], [GHJ], [J], [SiSm]).

4.1. Basic construction of finite inclusions. Let N be a von Neumann algebra admitting a normal faithful trace state τ and $1_N \in B \subseteq N$ a von Neumann subalgebra with the same identity (see [Chr], [J1], [Po1], [PiPo], [SiSm]).

Recall that the relative tensor product $L^2(N, \tau) \otimes_B L^2(N, \tau)$ over B of the N - B -bimodule ${}_N L^2(N, \tau)_B$ by the B - N -bimodule ${}_B L^2(N, \tau)_N$, constructed in [S1], is isomorphic, as an N - N -bimodule, to the N - N -correspondence \mathcal{H}_B associated to the conditional expectation $E_B : N \rightarrow N$ from N onto B . The latter being generated by the GNS construction applied to the binormal state

$$\Phi_B : N \otimes_{\max} N^\circ \rightarrow \mathbb{C} \quad \Phi_B(x \otimes y^\circ) := \tau(E_B(x)y).$$

According to S. Popa (see [Po1,2]), the inclusion $B \subseteq N$ is said to be *relatively amenable* if the standard bimodule ${}_N L^2(N, \tau)_N$ is weakly included in the relative coarse bimodule ${}_N L^2(N, \tau) \otimes_B L^2(N; \tau)_N$.

Let e_B be the oerthogonal projection in $\mathcal{B}(L^2(N, \tau))$ from $L^2(N, \tau)$ onto $L^2(B, \tau)$ and consider the *basic construction* $\langle N, B \rangle$, i.e. the von Neumann algebra in $\mathcal{B}(L^2(N, \tau))$ generated by N and the projection e_B . For example, if $B = \mathbb{C}1_N$ then $\langle N, B \rangle = \mathcal{B}(L^2(N, \tau))$ and when $B = N$ then $\langle N, B \rangle = N$.

Denoting by $\xi_\tau \in L^2(N, \tau)$ the cyclic vector representing τ one has

$$e_B(x\xi_\tau) = E_B(x)\xi_\tau, \quad e_B x e_B = E_B(x)e_B \quad x \in N.$$

It can be shown that an element $x \in N$ commutes with the projection e_B if and only if $x \in B$. Moreover, $\text{span}(N e_B N)$ is weakly*-dense in $\langle N, B \rangle$ and $e_B \langle N, B \rangle e_B = B e_B$. It can be shown that

$$\langle N, B \rangle = (J B J)' \subseteq \mathcal{B}(L^2(N, \tau))$$

so that $\langle N, B \rangle$ is semifinite since B is finite. In particular, there exists a unique normal, semifinite faithful trace Tr characterized by

$$\text{Tr}(x e_B y) = \tau(xy) \quad x, y \in N.$$

and there exists also a unique $N - N$ -bimodule map Φ from $\text{span}(N e_B N)$ into N satisfying

$$\Phi(x e_B y) = xy \quad x, y \in N, \quad \text{Tr} = \tau \circ \Phi.$$

The map Φ extends to a contraction between the N - N -bimodules $L^1(\langle N, B \rangle, \text{Tr})$ and $L^1(N, \tau)$ and satisfies

$$e_B X = e_B \Phi(e_B X) \quad X \in \langle N, B \rangle.$$

Moreover, $\Phi(e_B X) \in L^2(\langle N, B \rangle, \text{Tr})$ for all $X \in \langle N, B \rangle$. These properties enable us to prove that the identity correspondence $L^2(\langle N, B \rangle, \text{Tr})$ of the algebra $\langle N, B \rangle$ reduces to the relative correspondence \mathcal{H}_B when restricted to the subalgebra $N \subseteq \langle N, B \rangle$.

The following proposition is well known, we give the proof for sake of completeness.

Proposition 4.1. *The N - N -correspondences \mathcal{H}_B and $L^2(\langle N, B \rangle, \text{Tr})$ are isomorphic. In particular, the binormal state is given by*

$$\Phi_B(x \otimes y^\circ) = (e_B | x e_B y)_{L^2(\langle N, B \rangle, \text{Tr})} \quad x, y \in N$$

so that the cyclic vector representing the state Φ_B is $e_B \in L^2(\langle N, B \rangle, \text{Tr})$.

Proof. Let us consider the map Ψ defined on the domain

$$D(\Psi) := \text{span}\{[x \otimes y^\circ]_{\mathcal{H}_B} : x, y \in N\}$$

by

$$\Psi : D(\Psi) \rightarrow L^2(\langle N, B \rangle, \text{Tr}) \quad \Psi([x \otimes y^\circ]_{\mathcal{H}_B}) := x e_B y \quad x, y \in N.$$

Here $[x \otimes y^\circ]_{\mathcal{H}_B}$ denotes the element of \mathcal{H}_B image of the elementary tensor product $x \otimes y^\circ$, in the GNS construction of the state Φ_B . The map is well defined because $\|e_B\|^2 = \text{Tr}(e_B) = \tau(\Phi(e_B)) = \tau(1_N) = 1$ and $\|x e_B y\|_2 \leq \|x\| \cdot \|y\| \cdot \|e_B\|_2 = \|x\| \cdot \|y\|$. By the definition of the

Hilbert space \mathcal{H}_B , the map Ψ is densely defined. For $x, y \in N$ we have

$$\begin{aligned}
\|\Psi([x \otimes y^\circ]_{\mathcal{H}_B})\|_2^2 &= \|xe_B y\|_2^2 \\
&= \text{Tr}((xe_B y)^*(xe_B y)) \\
&= \text{Tr}(y^* e_B x^* x e_B y) \\
&= \text{Tr}(y^* e_B e_B x^* x e_B e_B y) \\
&= \text{Tr}((e_B x^* x e_B)(e_B y y^* e_B)) \\
&= \text{Tr}(E_B(x^* x) e_B E_B(y y^*) e_B) \\
&= \text{Tr}(E_B(x^* x) e_B E_B(y y^*)) \\
&= \tau(E_B(x^* x) E_B(y y^*)) \\
&= \tau(E_B(E_B(x^* x) y y^*)) \\
&= \tau(E_B(x^* x) y y^*) \\
&= \Phi_B(x^* x \otimes (y y^*)^\circ) \\
&= \Phi_B(x^* x \otimes (y^*)^\circ y^\circ) \\
&= \Phi_B((x^* \otimes (y^*)^\circ)(x \otimes y^\circ)) \\
&= \Phi_B((x \otimes y^\circ)^*(x \otimes y^\circ)) \\
&= \|[x \otimes y^\circ]_{\mathcal{H}_B}\|_{\mathcal{H}_B}^2.
\end{aligned}$$

By polarization, for all $\{x_j\}_{j=1}^n \subset N$ we have also

$$(\Psi([x_j \otimes y_j^\circ]_{\mathcal{H}_B}) | \Psi([x_k \otimes y_k^\circ]_{\mathcal{H}_B}))_2 = ([x_j \otimes y_j^\circ]_{\mathcal{H}_B} | [x_k \otimes y_k^\circ]_{\mathcal{H}_B})_{\mathcal{H}_B} \quad j, k = 1, \dots, n.$$

Consider now $\nu = \sum_{k=1}^n [x_k \otimes y_k^\circ]_{\mathcal{H}_B} \in D(\Psi)$ so that $\nu^* \nu = \sum_{j,k=1}^n [x_j^* x_k \otimes (y_k y_j^*)^\circ]_{\mathcal{H}_B} \in D(\Psi)$ and then

$$\|\Psi(\nu)\|_2^2 = \sum_{j,k=1}^n ([x_j \otimes y_j^\circ]_{\mathcal{H}_B} | [x_k \otimes y_k^\circ]_{\mathcal{H}_B})_{\mathcal{H}_B} = (\nu | \nu)_{\mathcal{H}_B} = \|\nu\|_{\mathcal{H}_B}^2.$$

Hence the map Ψ extends to an isometry from \mathcal{H}_B into $L^2(\langle N, B \rangle, \text{Tr})$ which is clearly an $N - N$ -bimodule map. Since $\text{Im}(\Psi) = \text{span}(N e_B N)$ is weakly*-dense in $\langle N, B \rangle$, it is also dense in $L^2(\langle N, B \rangle, \text{Tr})$. By the isometric property we have that $\text{Im}(\Psi)$ is closed so that Ψ is a surjective isometry. Finally, for $x, y \in N$ we compute

$$\begin{aligned}
(e_B | x e_B y)_{L^2(\text{Tr})} &= \text{Tr}(e_B x e_B y) \\
&= \text{Tr}(E_B(x) e_B y) \\
&= \tau(\Phi(E_B(x) e_B y)) \\
&= \tau(E_B(x) y) \\
&= \Phi_B(x \otimes y^\circ).
\end{aligned}$$

□

Definition 4.2. (*B-invariant Dirichlet forms*). Let N be a von Neumann algebra admitting a normal faithful tracial state τ and $1_N \in B \subseteq N$ a von Neumann subalgebra.

A Dirichlet form $(\mathcal{E}, \mathcal{F})$ on $L^2(N, \tau)$ is said to be a *B-invariant* if

$$b\mathcal{F} \subseteq \mathcal{F}, \quad \mathcal{E}(b\xi | \xi) = \mathcal{E}(\xi | b^*\xi) \quad b \in B, \quad \xi \in \mathcal{F}$$

and

$$\mathcal{F}b \subseteq \mathcal{F}, \quad \mathcal{E}(\xi b | \xi) = \mathcal{E}(\xi b^* | \xi) \quad b \in B, \quad \xi \in \mathcal{F}.$$

Since, by definition, a Dirichlet form is J -real, the above two properties are in fact equivalent.

In terms of the associated nonnegative, self-adjoint operator $(L, D(L))$, B -invariance means that the resolvent family $\{(\lambda + L)^{-1} : \lambda > 0\}$ is B -bimodular for some and hence all $\lambda > 0$

$$\begin{aligned} (\lambda + L)^{-1}(b\xi) &= b((\lambda + L)^{-1}\xi) \\ (\lambda + L)^{-1}(\xi b) &= ((\lambda + L)^{-1}\xi)b \quad \xi \in L^2(N, \tau), b \in B, \end{aligned}$$

or that, alternatively, the semigroup $\{e^{-tL} : t > 0\}$ is for some and hence all $t > 0$ a B -bimodular map

$$\begin{aligned} e^{-tL}(b\xi) &= b(e^{-tL}\xi) \\ e^{-tL}(\xi b) &= (e^{-tL}\xi)b \quad \xi \in L^2(N, \tau), b \in B. \end{aligned}$$

Since the Markovianity of the Dirichlet form implies that the semigroup and the resolvent commute with the modular conjugation J , we have that the B -invariance of the Dirichlet form provides that the semigroup and the resolvent belong to the relative commutant of B in the basic construction $\langle N, B \rangle$:

$$(\lambda + L)^{-1}, e^{-tL} \in (JBJ)' \cap B' = \langle N, B \rangle \cap B' \quad t > 0, \quad \lambda > 0.$$

Definition 4.3. (Relative discrete spectrum) We say that $(\mathcal{E}, \mathcal{F})$ or $(L, D(L))$ have *discrete spectrum relative to the inclusion* $B \subseteq N$ if the Markovian semigroup, or equivalently the resolvent, belongs to the *compact ideal space* $\mathcal{J}(\langle N, B \rangle)$ ([J], [Po2], [SiSm]) of the basic construction, generated by projections in $\langle N, B \rangle$ having finite trace:

$$\begin{aligned} e^{-tL} &\in \mathcal{J}(\langle N, B \rangle) \quad \text{for some and hence all } t > 0, \\ (\lambda + L)^{-1} &\in \mathcal{J}(\langle N, B \rangle) \quad \text{for some and hence all } \lambda > 0, \end{aligned}$$

Another way to state it is that the spectrum of $(L, D(L))$ is a discrete subset of \mathbb{R}_+ and that each eigenprojection has finite Tr trace.

Definition 4.4. (Relative spectral growth rate of Dirichlet forms) Let N be a von Neumann algebra admitting a normal faithful tracial state τ and $1_N \in B \subseteq N$ a von Neumann subalgebra. A Dirichlet form $(\mathcal{E}, \mathcal{F})$ on $L^2(N, \tau)$ which is B -invariant is said to be have

- *exponential spectral growth relative to* $B \subseteq N$ if $\text{Tr}(e^{-tL}) = +\infty$ for some $t > 0$;
- *subexponential spectral growth relative to* $B \subseteq N$ if $\text{Tr}(e^{-tL}) < +\infty$ for all $t > 0$.

Notice that, if $\text{Tr}(e^{-tL}) < +\infty$ for some $t > 0$, then $e^{-tL} \in \mathcal{J}(\langle N, B \rangle)$ so that $(L, D(L))$ has discrete spectrum relative to the inclusion $B \subseteq N$. This applies, in particular, to B -invariant Dirichlet forms $(\mathcal{E}, \mathcal{F})$ with subexponential spectral growth relative to $B \subseteq N$ which thus have necessarily discrete spectrum relative to the inclusion $B \subseteq N$.

Remark 4.5. Let \mathbb{E}^L be the spectral measure of the self-adjoint operator $(L, D(L))$. If the Dirichlet form is B -invariant then \mathbb{E}^L takes its values in the class of projections of the von Neumann algebra $\langle N, B \rangle$ and we can consider the positive measure $\nu_B^L := \text{Tr} \circ \mathbb{E}^L$ on $[0, +\infty)$, supported by the spectrum $\sigma(L)$. In the framework of quantum statistical mechanics, where the operator L may represent the total energy observable, the measure ν_B^L acquires the meaning of "density of states" in the sense that $\nu_B^L(\Omega)$ measures the number (relatively to B) of allowed energy levels located in a measurable subset $\Omega \subset \sigma(L)$. In this case the subspace $L^2(B, \tau) \subset L^2(N, \tau)$ may represent the manifold of ground states corresponding to the minimal allowable energy level (see also Example 4.7 below). The measure $\mu_B^L(d\lambda) := \lambda \nu_B^L(d\lambda)$ has then the meaning of "spectral energy density" in the sense that $\mu_B^L(\Omega)$ measures the energy of the system in a situation where all the allowed energy levels in $\Omega \subset \sigma(L)$ are occupied.

The subexponential spectral growth condition (relatively to $B \subseteq N$) can be rephrased in terms of the Laplace Transform $\hat{\nu}_B^L$ saying that its abscissa of convergence vanishes. In this situation $\hat{\nu}_B^L(\beta) = \text{Tr}(e^{-\beta L})$ is called the *partition function* of the system, it is defined for all $\beta > 0$ and the variable β is interpreted as the *inverse temperature*. The subexponential spectral growth condition also allows to consider the so called *Gibbs* normal states, defined by $\Phi_\beta(A) := \frac{\text{Tr}(Ae^{-\beta L})}{\text{Tr}(e^{-\beta L})}$, on the von Neumann (observable) algebra $\langle N, B \rangle$, for any fixed value of the inverse temperature $\beta > 0$. These states possess properties by which they can be regarded as equilibria of the system at the fixed value of the inverse temperature. Finally, notice that the subexponential spectral growth condition is equivalent to the requirement that the mean energy $\Phi_\beta(L) := \frac{\text{Tr}(Le^{-\beta L})}{\text{Tr}(e^{-\beta L})}$ of the system is finite for any $\beta > 0$ (see [BR]).

The following is the main result of this section.

Theorem 4.6. *Let N be a von Neumann algebra admitting a normal faithful tracial state τ and $1_N \in B \subseteq N$ a von Neumann subalgebra.*

If there exists a B -invariant Dirichlet form $(\mathcal{E}, \mathcal{F})$ on $L^2(N, \tau)$ having subexponential spectral growth relatively to $B \subseteq N$, then the inclusion $B \subseteq N$ is amenable.

Proof. Let us check first the following identity

$$(T^*|xe_B y)_{L^2(\text{Tr})} = (i_\tau(y^*)|T(i_\tau(x)))_{L^2(\tau)} \quad T \in \langle N, B \rangle \cap L^2(\langle N, B \rangle, \text{Tr}), \quad x, y \in N.$$

As $\text{span}(Ne_B N)$ is weakly* dense in $\langle N, B \rangle$, it is enough to prove the identity for $T \in Ne_B N$. If $T = ue_B v$ for some $u, v \in N$ we have

$$e_B y T x e_B = e_B y u e_B v x e_B = (e_B y u e_B)(e_B v x e_B) = E_B(yu) e_B E_B(vx) e_B$$

and then

$$\begin{aligned} (T^*|xe_B y)_{L^2(\text{Tr})} &= \text{Tr}(T x e_B y) \\ &= \text{Tr}(e_B y T x e_B) \\ &= \tau(\Phi(e_B y T x e_B)) \\ &= \tau(\Phi(E_B(yu) e_B E_B(vx) e_B)) \\ &= \tau(E_B(yu) \Phi(e_B E_B(vx) e_B)) \\ &= \tau(E_B(yu) \Phi(E_B(vx) e_B)) \\ &= \tau(E_B(yu) E_B(vx)) \\ &= \tau(E_B(yu E_B(vx))) \\ &= \tau(yu E_B(vx)) \\ &= (u^* y^* \xi_\tau | E_B(vx) \xi_\tau)_{L^2(\tau)} \\ &= (i_\tau(y^*) | u E_B(vx) \xi_\tau)_{L^2(\tau)} \\ &= (i_\tau(y^*) | u e_B(v(x \xi_\tau)))_{L^2(\tau)} \\ &= (i_\tau(y^*) | u e_B v(i_\tau(x)))_{L^2(\tau)} \\ &= (i_\tau(y^*) | T(i_\tau(x)))_{L^2(\tau)} \end{aligned}$$

so that the identity holds true. Under the hypothesis of subexponential spectral growth, we have that $T_t := e^{-tL} \in L^2(\langle N, b \rangle, \text{Tr}) \cap L^1(\langle N, b \rangle, \text{Tr})$ for all $t > 0$. Applying the above identity, we have that the binormal states

$$\Phi_t : N \otimes_{\max} N^\circ \rightarrow \mathbb{C} \quad \Phi_t(x \otimes y^\circ) := \frac{1}{(\xi_\tau | T_t \xi_\tau)_{L^2(N, \tau)}} (i_\tau(y^*) | T_t(i_\tau(x)))_{L^2(N, \tau)},$$

well defined, by strong continuity of the semigroup, for t sufficiently close to zero, may be represented for $t > 0$ as $\Phi_t(x \otimes y^\circ) = \frac{1}{(\xi_\tau | T_t \xi_\tau)_{L^2(N, \tau)}} (T_t | x e_B y)_{L^2(\langle N, b \rangle, \text{Tr})}$.

By the identity above, Φ_t extends as a normal state on the von Neumann algebra generated by the left and right representations of N in $L^2(\langle N, B \rangle, \text{Tr})$. The N - N -correspondence \mathcal{H}_t generated by Φ_t is thus a sub-correspondence of a multiple of the N - N -correspondence $L^2(\langle N, B \rangle, \text{Tr})$. Since the semigroup $\{T_t : t > 0\}$ strongly converges to the identity operator on $L^2(N, \tau)$, we obtain that the trivial correspondence from N to N is weakly contained in the relative correspondence \mathcal{H}_B . \square

Example 4.7. (Minimal and maximal inclusions of a Dirichlet form) Let N be a von Neumann algebra and τ a normal, faithful, tracial state and let $(\mathcal{E}, \mathcal{F})$ be a Dirichlet form on $L^2(N, \tau)$ with associated self-adjoint operator $(L, D(L))$.

Assume that $\inf \sigma(L) = 0$ and that this is an eigenvalue (not necessarily of finite multiplicity). The spectral projection P_0 onto the eigenspace corresponding to the Borel subset $\{0\} \subset [0, +\infty)$ can be represented as the strong limit $P_0 = \lim_{t \rightarrow +\infty} e^{-tL}$. Hence P_0 is a completely Markovian projection, so that there exists a von Neumann subalgebra $B_{\min} \subseteq N$ such that $P_0 = e_{B_{\min}}$. Obviously the associated Markovian semigroup is B_{\min} -bimodular and the Dirichlet form is B_{\min} -invariant.

Alternatively, one can consider the inclusion $B_{\max} \subset N$ where $B_{\max} := \{T_t : t > 0\}' \cap N$ is the relative commutant of the Markovian semigroup in N . Notice that by the Spectral Theorem $B_{\max} = \{T_t\}' \cap N$ for all $t > 0$. Obviously the associated Markovian semigroup is B_{\max} -bimodular and the Dirichlet form is B_{\max} -invariant.

Proposition 4.8. *Let (N, τ) be a finite von Neumann algebra with faithful, normal trace. Let $(\mathcal{E}, \mathcal{F})$ a Dirichlet form on $L^2(N, \tau)$ with generator $(L, D(L))$ having pure point spectrum made by distinct, isolated eigenvalues $\sigma(L) := \{\lambda_0 < \lambda_1 < \lambda_2 < \dots\}$ and assume $\lambda_0 := \inf \sigma(L) = 0$.*

Then $(\mathcal{E}, \mathcal{F})$ has discrete spectrum relative to B_{\min} (resp. B_{\max}) if and only if each eigenspace $E_\lambda \subset L^2(N, \tau)$, $\lambda \in \sigma(L)$, has finite coupling constant $\dim_{B_{\min}}(E_\lambda) < +\infty$ (resp. $\dim_{B_{\max}}(E_\lambda) < +\infty$) relative to B_{\min} (resp. B_{\max}).

Remark that the finite coupling constant $\dim_{B_{\min}}(E_\lambda) < +\infty$ (resp. $\dim_{B_{\max}}(E_\lambda) < +\infty$) relative to B_{\min} (resp. B_{\max}) is well defined for any eigenvalue $\lambda \in \sigma(L)$ because any eigenspace E_λ is obviously a left (and also right) B_{\min} -module (resp. B_{\max} -module). We refer to [GHJ Section 3.2] for the definition and properties of the Murray-von Neumann coupling constant.

Example 4.9. Let $K < \Gamma$ be an inclusion of countable, discrete groups and let $L(K) \subset L(\Gamma)$ be the inclusion of the finite von Neumann algebras generated by K and Γ , respectively. Their standard spaces coincide with $l^2(K)$ and $l^2(\Gamma)$ respectively and the projection $e_{L(K)}$ coincides with the projection from $l^2(\Gamma)$ onto its subspace $l^2(K)$.

Let $\ell : \Gamma \rightarrow [0, +\infty)$ be a c.n.d. function. The Dirichlet form $(\mathcal{E}_\ell, \mathcal{F}_\ell)$ associated to ℓ (introduced in Section 3.3) is $L(K)$ -invariant if and only if ℓ vanishes on K or, equivalently, if ℓ is a right K -invariant function. In this situation we have:

Proposition 4.10. *Let Γ be a countable, discrete group and let $L(\Gamma)$ be its left von Neumann algebras. Let $\ell : \Gamma \rightarrow [0, +\infty)$ be a c.n.d. function and $(\mathcal{E}_\ell, \mathcal{F}_\ell)$ the associated Dirichlet form. Denote by $H := \{s \in \Gamma : \ell(s) = 0\}$ the subgroup where ℓ vanishes. We then have*

i) $B_{\min} = B_{\max} = L(H)$;

ii) If K is a subgroup of G , then $(\mathcal{E}_\ell, \mathcal{F}_\ell)$ is $L(K)$ -invariant if and only if $K < H$. In this case :

ii.a) ℓ is $L(K)$ -biinvariant

ii.b) $(\mathcal{E}_\ell, \mathcal{F}_\ell)$ has discrete spectrum relative to $L(K) \subset L(\Gamma)$ if and only if the function

$$\ell_{G/K} : G/K \rightarrow [0, +\infty) \quad \ell_{G/K}(\tilde{s}) := \ell(s)$$

defined for $\tilde{s} = sK \in G/K$, is proper.

ii.c) If, for any $t > 0$, $\sum_{\tilde{s} \in G/K} e^{-t\ell_{G/K}(\tilde{s})} < +\infty$, then the inclusion $L(K) \subset L(G)$ is amenable.

Proof. i) Let $x = \sum_{t \in \Gamma} x(t)\lambda(t) \in B_{\max} = \{T_t : t > 0\}' \cap L(\Gamma)$. We then have

$$x\delta_e = x(I + L)^{-1}\delta_e = (I + L)^{-1}x\delta_e,$$

which implies $0 = L(x\delta_e) = L(\sum_{t \in \Gamma} x(t)\lambda(t)\delta_e) = L(\sum_{t \in \Gamma} x(t)\delta_t) = \sum_{t \in \Gamma} x(t)\ell(t)\delta_t$. So that $x(t)\ell(t) = 0$ for all $t \in \Gamma$ which in turn implies $x \in L(H) = B_{\min}$. The reverse inclusion is obvious.

ii) follows from the arguments of the example above. For ii) b) just notice that $\lambda(s)e_{L(K)}\lambda(s)^{-1}$ is the orthogonal projection P_{sK} onto the subspace $l^2(sH)$. Hence the eigenspace E_λ corresponding to the eigenvalue $\lambda \in \sigma(L)$ is given by $\bigoplus_{\tilde{s} \in G/K, \ell(s)=\lambda} l^2(sK)$. Hence, L will have discrete spectrum relative to K if and only if each of these sums is finite (i.e. for all λ) and the set of values of ℓ is discrete, i.e. $\ell^{-1}(\{\lambda\})/K$ is finite in G/K , i.e. $\ell_{G/K} : G/K \rightarrow [0, +\infty)$ is proper.

For ii.c), note that in the basic construction for $B = L(K) \subset N = L(G)$, $P_{sK} = \lambda(s)e_B\lambda(s)$ belongs to $\langle N, B \rangle$ and has trace 1. Hence $Tr(e^{-tL}) = \sum_{\tilde{s} \in G/K} e^{t\ell(s)}$ for all $t > 0$. \square

Remark 4.11.

i) If the Dirichlet form $(\mathcal{E}_\ell, \mathcal{F}_\ell)$ has discrete spectrum relative to $L(K)$ then

a) the function $\ell_{G/H} : G/H \rightarrow [0, +\infty)$ is proper and $(\mathcal{E}_\ell, \mathcal{F}_\ell)$ has discrete spectrum relative to $L(H)$;

b) $\ell_{G/K}$ being left K -invariant, hence constant on left K -cosets, and proper, left K -cosets in G/K must be finite sets. In other words, K is quasi-normal in G .

ii) On the other hand, if $(\mathcal{E}_\ell, \mathcal{F}_\ell)$ has discrete spectrum relative to $L(H)$ then the function $\ell_{G/K} : G/K \rightarrow [0, +\infty)$ will be constant onto the right H -coset in G/K . Thus $(\mathcal{E}_\ell, \mathcal{F}_\ell)$ has discrete spectrum relative to $L(K)$ if and only if each right H -coset in G/K is a finite union of K -cosets, which happens if and only if K has finite index in H , i.e. when the homogeneous space H/K is finite.

5. A SPECTRAL APPROACH TO THE RELATIVE HAAGERUP PROPERTY

As already mentioned in the Introduction, in a recent work [CaSk], M. Caspers and A. Skalski characterized von Neumann algebras having Property (H) in terms of the existence of a Dirichlet form with discrete spectrum. In the spirit of the previous section, we extend their result to *relative property (H)*, as defined by S. Popa [Po1,2], for inclusions of von Neumann algebras, using a completely different approach. We will make use of the following well known properties:

5.1. Let (N, τ) be a von Neumann algebra endowed with a normal, faithful trace and let $\varphi : N \rightarrow N$ be a completely positive, normal contraction such that $\tau \circ \varphi \leq \tau$. Then

i) there exists a contraction $T_\varphi \in \mathcal{B}(L^2(N, \tau))$ characterized by

$$T_\varphi(x\xi_\tau) = \varphi(x)\xi_\tau \quad x \in N;$$

ii) there exists a completely positive, normal contraction $\varphi^* : N \rightarrow N$ such that

$$T_{\varphi^*} = (T_\varphi)^*$$

or, more explicitly,

$$(\varphi^*(y)\xi_\tau|x\xi_\tau) = (y\xi_\tau|\varphi(x)\xi_\tau) \quad x, y \in N.$$

Definition 5.2. ([Po1,2]) Let N be a finite von Neumann algebra and $B \subseteq N$ a von Neumann subalgebra. Then N is said to have *Property (H) relative to B* if there exist a normal, faithful tracial state τ on N and a net $\{\varphi_i : i \in I\}$ of normal completely positive, B -bimodular maps on N satisfying the conditions

- i) $\tau \circ \varphi_i \leq \tau$
- ii) $T_{\varphi_i} \in \mathcal{J}(\langle N, B \rangle)$
- iii) $\lim_{i \in I} \|x\xi_\tau - T_{\varphi_i}(x\xi_\tau)\|_2 = 0$ for all $x \in N$.

In this definition $\mathcal{J}(\langle N, B \rangle)$ is the compact ideal space, i.e. the norm closed ideal generated by projections with finite trace in $\langle N, B \rangle$ and T_{φ_i} is the operator defined in item 5.1 above.

By a remark of S. Popa [Po2], the maps φ_i in the definition above can be chosen to be contractions. In the following we shall always assume this property for approximating nets of the identity map of a von Neumann algebra.

Theorem 5.3. *Let N be a finite von Neumann algebra with countably decomposable center and faithful tracial state τ . Let $B \subseteq N$ be a sub-von Neumann algebra such that $L^2(N, \tau)$, as B -module, admits a countable base. Then the following properties are equivalent*

- i) N has *Property (H) relative to B*
- ii) *there exists a B -invariant Dirichlet form $(\mathcal{E}, \mathcal{F})$ on $L^2(N, \tau)$ with discrete spectrum relative to B .*

Proof. Assume that there exists a Dirichlet form $(\mathcal{E}, \mathcal{F})$ on $L^2(N, \tau)$ with discrete spectrum relative to B . Hence, the associated generator $(L, D(L))$ has its resolvent in the compact ideal space: $(\lambda + L)^{-1} \in \mathcal{J}(\langle N, B \rangle)$. Then for all $\lambda > 0$, $S_\lambda := \lambda(\lambda + L)^{-1} \in \mathcal{J}(\langle N, B \rangle)$. Moreover, any S_λ is Markovian on $L^2(N, \tau)$ which implies that there exists a completely positive contraction $\varphi_\lambda : N \rightarrow N$ determined by $S_\lambda(x\xi_\tau) = \varphi_\lambda(x)\xi_\tau$ for $x \in N$. Since the S_λ are self-adjoint on $L^2(N, \tau)$, the φ_λ are symmetric with respect to the trace: $\tau(\varphi_\lambda(x)y) = \tau(x\varphi_\lambda(y))$ for all $x, y \in N$. This implies that

$$\tau(\varphi_\lambda(x)) = \tau(\varphi_\lambda(1_N)x) \leq \tau(x) \quad x \in N_+.$$

Last condition iii) in Definition 5.2 above comes from the strong continuity of the resolvent:

$$\lim_{\lambda \rightarrow +\infty} \|\xi - S_\lambda \xi\|_2 = 0 \quad \xi \in L^2(N, \tau).$$

The theorem is proved in the "if" direction. In the reverse direction, let us suppose that $B \subseteq N$ is an inclusion with relative property (H) and that $L^2(N, \tau)$ is separable as B -module. Let $\{\varphi_n : n \in \mathbb{N}\}$ be a sequence of normal, completely positive, B -bimodular contractions of N , satisfying the conditions of the definition above. By [Po2 Proposition 2.2] such a sequence always exists. Each φ_n extends by $T_n(x\xi_\tau) := \varphi_n(x)\xi_\tau$ to a B -bimodular contraction T_n of $L^2(N, \tau)$, which belongs to the compact ideal space $\mathcal{J}(\langle N, B \rangle)$. It is also completely positive with respect to the standard positive cone $L^2_+(N, \tau)$ and its matrix amplifications. It is easy to check that the maps φ_n^* appearing in item 5.1 above, have the same properties as the φ_n 's in definition above. Replacing each φ_n by $(\varphi_n + \varphi_n^*)/2$, we can suppose, without loss of generality, that the φ_n are symmetric with respect to τ so that the corresponding T_n are completely positive, self-adjoint contractions on $L^2(N, \tau)$.

Let $\{\xi_k \in L^2(N, \tau) : k \in \mathbb{N}\}$ be an orthonormal basis for the left B -module $L^2(N, \tau)$. Recall that this means that $L^2(N, \tau) = \overline{\bigoplus_{k \in \mathbb{N}} B\xi_k}$.

For any $k \in \mathbb{N}$ one has $\lim_{n \rightarrow +\infty} T_n \xi_k = \xi_k$ and hence there exists $n_k \in \mathbb{N}$ such that

$$(\xi_j | (I - T_{n_k}) \xi_j) \leq 2^{-k}, \quad k \in \mathbb{N}, \quad j \in \{0, 1, \dots, k\}.$$

Let us consider the quadratic form $(\mathcal{E}, \mathcal{F})$ on $L^2(n, \tau)$ defined by

$$\mathcal{E}[\xi] := \sum_{k \in \mathbb{N}} (\xi | (I - T_{n_k}) \xi) \quad \xi \in L^2(N, \tau).$$

The domain $\mathcal{F} \subseteq L^2(N, \tau)$ being understood as the subspace where the quadratic form is finite. Note first that \mathcal{E} is densely defined since $b\xi_j \in \mathcal{F}$ for all $b \in B$ and $j \in \mathbb{N}$. Noticing that each $\xi \mapsto (\xi | (I - T_{n_k}) \xi)$ is a bounded symmetric Dirichlet form on $L^2(N, \tau)$ we see that $(\mathcal{E}, \mathcal{F})$ is a lower semicontinuous, hence closed Dirichlet form on $L^2(N, \tau)$. The B -modularity of $(\mathcal{E}, \mathcal{F})$ being obvious, what is left to prove is the relative discrete spectrum property.

The generator $(L, D(L))$ associated to $(\mathcal{E}, \mathcal{F})$, given by $L = \sum_{k \in \mathbb{N}} (I - T_{n_k})$, appears as the increasing limit $L = \lim_{m \uparrow +\infty} L_m$ of the bounded operators

$$L_m := \sum_{k=0}^m (I - T_{n_k}) = (m+1)I - \Theta_m$$

where $\Theta_m := \sum_{k=0}^m T_{n_k}$. The important fact is that Θ_m belongs to the compact ideal space $\mathcal{J}(\langle N, B \rangle)$.

Let q_m be the spectral projection of Θ_m corresponding to the interval $[0, (m+1)/2]$ and $p_m := I - q_m$ the spectral projection corresponding to the interval $((m+1)/2, m+1]$. On one hand we have $\tau(p_m) < +\infty$, which implies $(I + L)^{-1/2} p_m \in \mathcal{J}(\langle N, B \rangle)$. On the other hand by spectral calculus we have

$$q_m (I + L)^{-1} q_m \leq q_m (I + L_m)^{-1} q_m \leq \frac{2}{m+1} q_m$$

which implies

$$\|q_m (I + L)^{-1/2}\| \leq \sqrt{\frac{2}{m+1}}.$$

Finally, we have $(I + L)^{-1/2} = \lim_{m \rightarrow +\infty} (I + L)^{-1/2} p_m$ for the uniform norm, which implies that $(I + L)^{-1/2}$ and $(I + L)^{-1}$ are in $\mathcal{J}(\langle N, B \rangle)$. \square

Remark 5.4. It would be very interesting to generalize the above result to type III (factors) inclusions. The first difficulties rely on a convenient identification of the ideal of relative compact operators.

6. RELATIVE PROPERTY (H) FOR INCLUSIONS OF DISCRETE GROUPS AND CONDITIONALLY NEGATIVE DEFINITE FUNCTIONS

In this section we extend a well known characterization of groups with Property (H) in terms of the existence of a proper c.n.d. function, to inclusions of discrete groups $H < G$.

We denote by λ and ρ the left and right regular representation of G in the Hilbert space $l^2(G)$. If $(\delta_t)_{t \in G}$ is the canonical orthonormal basis of $l^2(G)$, then

$$\lambda(s)\delta_t = \delta_{st} \quad \rho(s)\delta_t = \delta_{ts^{-1}} \quad s, t \in G.$$

The associated inclusion of von Neumann algebras is

$$B = L(H) = \lambda(H)'' \subset \lambda(G)'' = L(G) = N.$$

$L(G)$ has the canonical finite trace $\tau : L(G) \rightarrow \mathbb{C}$ determined by $\tau(\lambda(s)) = \delta_e(s)$ for $s \in G$. The standard space $L^2(L(G), \tau)$ identifies canonically with $l^2(G)$ through the unitary map

determined by $\lambda(s)\xi_\tau \rightarrow \delta_s$ for $s \in G$. The projection e_B of the basic construction is the orthogonal projection from $l^2(G)$ onto its subspace $l^2(H)$ given by the multiplication operator by the characteristic function χ_H of the subset H of G .

Lemma 6.1. (*Basic construction for group inclusions*).

i) The basic construction $\langle N, B \rangle$ is the commutant $\rho(H)'$ of the right regular representation restricted to H .

ii) For $T \in \langle N, B \rangle$, the map $\varphi_T : G \rightarrow \mathbb{C}$ given by $s \mapsto (\delta_s, T\delta_s)$ is right H -invariant :

$$\varphi_T(sh) = \varphi_T(s) \quad s \in G, h \in H.$$

iii) The canonical trace on $\langle N, B \rangle$ is given by the formula

$$Tr(T) = \sum_{\tilde{s} \in G/H} \varphi_T(s) = \sum_{\tilde{s} \in G/H} (\delta_s, T\delta_s) \quad T \in \langle N, B \rangle$$

where it is understood that the map $\tilde{s} \mapsto s$ indicates a section of the projection $G \rightarrow G/H$.

Proof. i) On one side, we have $\langle N, B \rangle = (JBJ)'$. On the other side, it is an elementary fact that $J\lambda(h)J = \rho(h)$, $h \in H$. ii) For $h \in H$ and $T \in \mathcal{B}(l^2(G))$ commuting with $\rho(h)$, we compute

$$\varphi_T(sh) = (\rho(h)^{-1}\delta_s, T\rho(h)^{-1}\delta_s) = (\delta_s, \rho(h)T\rho(h)^{-1}\delta_s) = \varphi_T(s) \quad s \in G, h \in H.$$

iii) Fix an arbitrary section $\sigma : G/H \rightarrow G$ of the homogeneous space G/H . The formula

$$\varphi(T) = \sum_{\gamma \in G/H} (\delta_{\sigma(\gamma)}, T\delta_{\sigma(\gamma)}) \in [0, +\infty] \quad T \in \langle N, B \rangle_+ = \rho(H)'_+$$

defines a normal faithful weight φ on the von Neumann algebra $\langle N, B \rangle$. We claim that this weight is equal to the natural trace Tr .

For $x \in L(G)$ with Fourier expansion $x = \sum_{s \in G} x(s)\lambda(s)$ (the series $\{x(s)\}_{s \in G}$ converges at least in the ℓ^2 -sense), let us compute

$$\begin{aligned} \varphi(x^*e_Bx) &= \sum_{\gamma \in G/H} (\delta_{\sigma(\gamma)}, x^*e_Bx\delta_{\sigma(\gamma)}) \\ &= \sum_{\gamma \in G/H} \|e_Bx\delta_{\sigma(\gamma)}\|^2 \end{aligned}$$

with

$$\begin{aligned} \|e_Bx\delta_{\sigma(\gamma)}\|^2 &= \|e_B \sum_{s \in G} x(s)\delta_{s\sigma(\gamma)}\|^2 \\ &= \|e_B \sum_{s \in G} x(s\sigma(\gamma)^{-1})\delta_s\|^2 \\ &= \left\| \sum_{h \in H} x(h\sigma(\gamma)^{-1})\delta_h \right\|^2 \\ &= \sum_{s \in H\sigma(\gamma)^{-1}} \|x(s)\|^2 \end{aligned}$$

and finally

$$\varphi(x^*e_Bx) = \sum_{\gamma \in G/H} \sum_{s \in H\sigma(\gamma)^{-1}} \|x(s)\|^2 = \sum_{s \in G} \|x(s)\|^2 = \tau(x^*x) = Tr(x^*e_Bx).$$

This proves that the normal weight φ is semifinite and that it coincides with the canonical trace Tr on elements of the form x^*e_Bx , $x \in N$.

The Radon-Nykodim derivative $d\varphi/dTr$ must be equal to 1, hence $\varphi = Tr$. □

Theorem 6.2. *Let G be a countable discrete group and $H < G$ a subgroup. Then the inclusion of von Neumann algebras $L(H) \subset L(G)$ has the relative Property (H) if and only if there exists a conditionally negative type function $\ell : G \rightarrow [0, +\infty)$ such that*

- i) $\ell|_H = 0$
- ii) ℓ is proper on G/H .

Proof. Let us suppose that i) and ii) are satisfied. Then for any $t > 0$, $e^{-t\ell}$ is a positive type function equal to 1 on H and it induces a normal, completely positive, trace preserving contraction ϕ_t on the von Neumann algebra $L(G)$, characterized by

$$\phi_t(\lambda_u) = e^{-t\ell(u)}\lambda_u \quad u \in G.$$

Let (π, H_π) be the orthogonal representation and $c : G \rightarrow H_\pi$ the 1-cocycle associated to the c.n.d. function ℓ . If $\ell(u) = 0$ then $0 = \ell(u) = \|c(u)\|_{H_\pi}^2$ so that $c(u) = 0$ for all $u \in H$. Then, $c(vu) = c(v) + \pi(v)c(u) = c(v)$ for all $v \in G$ and $u \in H$ and $\ell(vu) = \ell(v)$, $v \in G$, $u \in H$. Finally, as $\ell(v^{-1}) = \ell(v)$ for all $v \in G$, one also has $\ell(uv) = \ell(v)$ for all $v \in G$ and $u \in H$.

Consequently we shall have

$$\phi_t(\lambda_v\lambda_u) = e^{-t\ell(vu)}\lambda_v\lambda_u = e^{-t\ell(v)}\lambda_v\lambda_u = \phi_t(\lambda_v)\lambda_u \quad v \in G, u \in H$$

and

$$\phi_t(ab) = \phi_t(a)b \quad a \in L(G), b \in L(H).$$

Similarly

$$\phi_t(ba) = b\phi_t(a) \quad a \in L(G), b \in L(H).$$

This proves that ϕ_t is a $L(H)$ -bimodular map for all $t > 0$.

The self-adjoint operator T_t on $l^2(G)$, induced by ϕ_t on $L(G)$, is just the multiplication operator by the function $e^{-t\ell}$. Its spectrum coincides with the set of values $e^{-t\lambda}$ where λ runs in the range $\ell(G)$ which is, by assumption ii), a discrete subset of $[0, +\infty)$.

The eigenspace E_λ , corresponding to the eigenvalue $e^{-t\lambda}$, is the set of functions in $l^2(G)$ supported by $S_\lambda := \{s \in G : \ell(s) = \lambda\}$. Again by assumption ii), S_λ is a finite union of right H -cosets: $S_\lambda = \bigcup_{i=1}^k u_i H$ for some $u_1, \dots, u_k \in G$. The corresponding spectral projection P_λ is then the sum of projections on those cosets:

$$P_\lambda = \sum_{i=1}^k \text{multiplication by } \chi_{u_i H}.$$

Since, for $B := L(H) \subset L(G) = N$, the projection e_B is just the multiplication operator by the characteristic function χ_H of the subgroup, the projection $\lambda_u e_B \lambda_u^{-1}$, for $u \in G$, is the multiplication operator by the function χ_{uH} . The spectral projection P_λ onto E_λ is then a finite sum of projections of trace one (for the trace Tr), and its trace is equal to the number of right H -cosets in S_λ :

$$\text{Tr}(P_\lambda) = \text{Tr}\left(\sum_{i=1}^k \lambda_{u_i} e_B \lambda_{u_i}^{-1}\right) = \sum_{i=1}^k \text{Tr}(\lambda_{u_i} e_B \lambda_{u_i}^{-1}) = \sum_{i=1}^k \text{Tr}(e_B) = \sum_{i=1}^k 1 = k.$$

This proves that T_t belong to $\mathcal{J}(\langle N, B \rangle)$, which ends the proof in the forward direction.

Conversely, let us suppose that the inclusion $L(H) \subset L(G)$ has the relative property (H). By Theorem 5.3, there exists an $L(H)$ -invariant symmetric Dirichlet form with generator L and discrete spectrum relative to the subalgebra $B = L(H)$.

Let us observe that, for $\varepsilon > 0$, the resolvent maps $(I + \varepsilon L)^{-1}$ are a completely positive, normal contractions of the von Neumann algebra $N = L(G)$ and that the map

$$\omega_\varepsilon : G \rightarrow [0, +\infty) \quad \omega_\varepsilon(s) = (\delta_s, (I + \varepsilon L)^{-1} \delta_s) = \tau(\lambda(s)^*(I + \varepsilon L)^{-1}(\lambda(s)))$$

is positive definite on G and H -right invariant. Notice that ω_ε is symmetric because it is positive definite and real, as L is self-adjoint. Moreover, by the weak*-continuity of the resolvent, one has

$$\lim_{\varepsilon \rightarrow 0} \omega_\varepsilon(s) = 1 \quad s \in G.$$

We claim that ω_ε vanishes at infinity on the quotient space G/H . This holds true because $(I + \varepsilon L)^{-1}$ being in the compact ideal $\mathcal{J}(\langle\langle N, B \rangle\rangle)$, it will be a uniform limit of trace class operators $T_n \in L^1(\langle\langle N, B \rangle\rangle, Tr)$ and that, for such T_n , the function $s \mapsto (\delta_s, T_n \delta_s)$ is summable on G/H (by Lemma 6.1 iii) thus vanishing at infinity.

Let $(F_k)_{k \geq 1}$ be an increasing family of finite subsets of G such that $\cup_k F_k = G$. For any k , let us choose $\varepsilon_k > 0$ such that

$$0 \leq 1 - \omega_{\varepsilon_k}(s) \leq 2^{-k} \quad s \in F_k$$

and consider the function $\ell : G \rightarrow [0, +\infty)$ defined by

$$\ell(s) = \sum_{k=1}^{\infty} (1 - \omega_{\varepsilon_k}(s)) \quad s \in G.$$

By the choice of ε_k , the series converges for any s in any F_k and thus for any $s \in G$ so that ℓ is well defined on G . The function ℓ is c.n.d. as a sum of c.n.d. functions and a right H -invariant function as a sum of right- H -invariant functions. It can thus be considered as a function on G/H .

Moreover, as the functions ω_ε vanish at infinity on G/H , the set $\Gamma_k = \{\tilde{s} \in G/H \mid \omega_{\varepsilon_k}(s) \geq 1/2\}$ is finite and for $\tilde{s} \notin \cup_{k=1}^N \Gamma_k$, one has $\ell(s) \geq N/2$. Hence, for any $N \in \mathbb{N}$, the set $\{\tilde{s} \in G/H \mid \ell(s) \leq N/2\}$ is finite, which proves that ℓ is proper on G/H . \square

Corollary 6.3. *If $H < G$ and $L(H) \subset L(G)$ has relative property (H), then H is a quasi normal subgroup: each orbit of the left action of H on the right-cosets space G/H is finite.*

Proof. The c.n.d. function ℓ constructed in Theorem 6.2 is left H -invariant, as it clearly satisfies $\ell(s^{-1}) = \ell(s)$ for $s \in G$. It is thus constant on the orbits of the left action of H on the right cosets space G/H . A subset of G/H on which the proper function ℓ is constant must be finite. \square

Corollary 6.4. *If $H < G$ is normal, then the inclusion $L(H) \subset L(G)$ has relative property (H) if and only if the quotient group G/H has the Haagerup property.*

Example 6.5. Let us consider the free group \mathbb{F}_2 with generators $a, b \in \mathbb{F}_2$ and the abelian subgroup $H := \{a^k : k \in \mathbb{Z}\}$ isomorphic to the additive group \mathbb{Z} of integer numbers.

H is not quasi-normal in \mathbb{F}_2 , so that the inclusion $H < G$ has not the relative property (H), though both G and H have property (H).

More generally, for countable discrete groups G_1, G_2 , the inclusion $G_1 < G_1 * G_2$ is not quasi-normal as soon as G_1 is infinite and G_2 has at least two elements. Hence, this inclusion has not the relative property H .

REFERENCES

- [Ara] H. Araki, Some properties of modular conjugation operator of von Neumann algebras and a non-commutative Radon-Nikodym theorem with a chain rule, *Pacific J. Math.* **50** (1974), 309-354.
- [BeDe] A. Beurling and J. Deny, Dirichlet spaces, *Proc. Nat. Acad. Sci.* **45** (1959), 208-215.
- [Bra] M. Brannan, Approximation properties for free orthogonal and free unitary quantum groups, *J. Reine Angew. Math.* **672** (2012), 223-251.
- [BR] Bratteli O., Robinson D.W., “Operator algebras and Quantum Statistical Mechanics 1”, Second edition, 505 pages, Springer-Verlag, Berlin, Heidelberg, New York, 1987.
- [CaSk] M. Caspers, A. Skalski, The Haagerup approximation property for von Neumann algebras via quantum Markov semigroups and Dirichlet forms, *Comm. Math. Phys.* **336** no. 3, (2015), 1637–1664.
- [CCJJV] P.-A. Cherix, M. Cowling, P. Jolissaint, P. Julg, A. Valette, “Groups with the Haagerup property. Gromov’s a-T-menability”, Progress in Mathematics, 197, Birkhäuser Verlag, Basel, 2001
- [Chr] E. Christensen, Subalgebras of finite algebras, *Math. Ann.* **243** (1979), 17–29.
- [C1] F. Cipriani, Dirichlet forms and Markovian semigroups on standard forms of von Neumann algebras, *J. Funct. Anal.* **147** (1997), no. 1, 259–300.
- [C2] F. Cipriani, “Dirichlet forms on Noncommutative spaces”, Springer ed. L.N.M. 1954, 2007.
- [C3] F. Cipriani, “Noncommutative potential theory: A survey”, *J. of Geometry and Physics* **105** (2016), 25–59.
- [CFK] F. Cipriani, U. Franz, A. Kula, Symmetries of Lévy processes on compact quantum groups, their Markov semigroups and potential theory, *J. Funct. Anal.* **266** (2014), no. 5, 2789–2844.
- [CS1] F. Cipriani, J.-L. Sauvageot, Derivations as square roots of Dirichlet forms, *J. Funct. Anal.* **201** (2003), no. 1, 78–120.
- [CS3] F. Cipriani, J.-L. Sauvageot, Fredholm modules on P.C.F. self-similar fractals and their conformal geometry, *Comm. Math. Phys.* **286** (2009), no. 2, 541–558.
- [CS5] F. Cipriani, J.-L. Sauvageot, Negative type functions on groups with polynomial growth, in “Noncommutative Analysis, Operator Theory and Applications”, *Operator Theory Advances and Applications* **252** (2016), D. Alpay, F. Cipriani, F. Colombo, D. Guido, I. Sabadini, J.-L. Sauvageot eds., Birkhauser
- [CGIS1] F. Cipriani, D. Guido, T. Isola, J.-L. Sauvageot, “Integrals and Potential of differential 1-forms on the Sierpinski Gasket”, *Adv. in Math.* **239** (2013), 128–163.
- [CGIS2] F. Cipriani, D. Guido, T. Isola, J.-L. Sauvageot, “Spectral triples for the Sierpinski Gasket”, *J. Funct. Anal.*, **266** (2014), 4809–4869.
- [Co1] A. Connes, Caractérisation des espaces vectoriels ordonnés sous-jacents aux algèbres de von Neumann, *Ann. Inst. Fourier (Grenoble)* **24** (1974), 121-155.
- [Co2] A. Connes, “Noncommutative Geometry”, Academic Press, 1994.
- [DFSW] M. Daws, P. Fima, A. Skalski, S. White, The Haagerup property for locally compact quantum groups, *J. Reine Angew. Math.* **711** (2016), 189-229.
- [deH] P. de la Harpe, “Topics in Geometric Group Theory”, Chicago Lectures in Mathematics, The University of Chicago Press, 2000.
- [Dix] J. Dixmier, “Les C*-algèbres et leurs représentations”, Gauthier–Villars, Paris, 1969.
- [FOT] M. Fukushima, Y. Oshima, M. Takeda, “Dirichlet Forms and Symmetric Markov Processes”, de Gruyter Studies in Mathematics 19, 2nd ed. 2010.

- [GHJ] F.M. Goodman, P. de la Harpe, V.F.R. Jones, “Coxeter graphs and towers of algebras”, Mathematical Sciences Research Institute Publications, 14, Springer-Verlag, New York, 1989. x+288 pp..
- [GL1] S. Goldstein, J.M. Lindsay, Beurling-Deny conditions for KMS-symmetric dynamical semigroups, *C. R. Acad. Sci. Paris Sr. I Math.* **317** no. 11, (1993), 1053–1057.
- [GK] Guentner-Kaminker, Exactness and uniform embeddability of discrete groups, *J. London Math. Soc.* **70**, (2004), 703-718.
- [Haa1] U. Haagerup, Standard forms of von Neumann algebras, *Math. Scand.* **37** (1975), 271-283.
- [Haa2] U. Haagerup, An example of a nonnuclear C*-algebra, which has the metric approximation property, *Invent. Math.* **50** (1978), no. 3, 279-293.
- [Ki] J. Kigami, “Analysis on Fractals”, Cambridge Tracts in Mathematics vol. **143**, Cambridge University Press, 2001.
- [J] V.F.R. Jones, Index for subfactors., *Invent. Math.* **72** (1983), 1-26.
- [Po1] S. Popa, Correspondences, *INCREST Bucharest preprint, unpublished* (1986), 95 pages.
- [Po2] S. Popa, On a class of type II_1 factors with Betti numbers invariants, *Annals of Math.* **163** (2006), 809-899.
- [R] W. Rudin, “Principles of Mathematical Analysis”, International Series in Pure and Applied Mathematics, McGraw-Hill Higher Education, 1976.
- [S1] J.-L. Sauvageot, Sur le produit tensoriel relatif d’espaces de Hilbert, *J. Operator Theory* **9** (1983), 237-252.
- [S4] J.-L. Sauvageot, Semi-groupe de la chaleur transverse sur la C*-algèbre d’un feuilletage riemannien, *J. Funct. Anal.* **142** (1996), 511-538.
- [SiSm] A.M. Sinclair, R.R. Smith, “Finite von Neumann algebras and Masas”, London Mathematical Society Lectures Notes Series 351, Cambridge University Press, 2008.
- [T] M. Takesaki, “Theory of Operator Algebras I”, Encyclopedia of Mathematical Physics, 415 pages, Springer-Verlag, Berlin, Heidelberg, New York, 2000.
- [V1] D.V. Voiculescu, The analogues of entropy and of Fisher’s information measure in free probability theory, *Invent. Math.* **132** (1998), 189-227.
- [vN] J. von Neumann, Zur allgemeinen Theorie des Masses *Fund. Math.* **13** (1) (1929), 73-111.

DIPARTIMENTO DI MATEMATICA, POLITECNICO DI MILANO, PIAZZA LEONARDO DA VINCI 32, 20133 MILANO, ITALY.

E-mail address: fabio.cipriani@polimi.it

INSTITUT DE MATHÉMATIQUES, CNRS-UNIVERSITÉ DENIS DIDEROT, F-75205 PARIS CEDEX 13, FRANCE

E-mail address: jean-luc.sauvageot@imj-prg.fr